

Loughborough University Institutional Repository

Removal of antiretroviral drugs stavudine and zidovudine in water under UV254 and UV254/H₂O₂ processes: Quantum yields, kinetics and ecotoxicology assessment

This item was submitted to Loughborough University's Institutional Repository by the/an author.

Citation: RUSSO, D. ...et al., 2018. Removal of antiretroviral drugs stavudine and zidovudine in water under UV254 and UV254/H₂O₂ processes: Quantum yields, kinetics and ecotoxicology assessment. *Journal of Hazardous Materials*, 349, pp. 195-204.

Additional Information:

- This paper was accepted for publication in the journal *Journal of Hazardous Materials* and the definitive published version is available at <https://doi.org/10.1016/j.jhazmat.2018.01.052>

Metadata Record: <https://dspace.lboro.ac.uk/2134/32719>

Version: Accepted for publication

Publisher: © Crown Copyright Published by Elsevier

Rights: This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the published version.

Manuscript Number: HAZMAT-D-17-03351

Title: Removal of antiretroviral drugs stavudine and zidovudine in water under UV254 and UV254/H₂O₂ processes: quantum yields, kinetics and ecotoxicology assessment

Article Type: Research Paper

Keywords: photodegradation; microreactor; mutagenicity and genotoxicity; water reuse; antiretroviral

Corresponding Author: Professor Raffaele Marotta, Ph.D

Corresponding Author's Institution: University

First Author: Danilo Russo

Order of Authors: Danilo Russo; Antonietta Siciliano; Marco Guida; Roberto Andreozzi; Nuno M Reis; Gianluca Li Puma; Raffaele Marotta, Ph.D

Abstract: The concentration of antiretroviral drugs in wastewater treatment plants (WWTP) effluents and surface waters of developed and developing countries, especially in the African region more affected by HIV, has increased significantly in recent years due to their widespread use. The presence of antiretroviral in natural water bodies limits the possibility of reuse of such waters, after traditional disinfection process (i.e. UV, chlorine) for civil and irrigation purposes. The removal of stavudine and zidovudine under UV254 or UV254/H₂O₂ irradiation was investigated in distilled water. The quantum yield of direct photolysis and the kinetic constant of reaction of hydroxyl radical with the antiretrovirals at different pH have been evaluated. A battery of ecotoxicological tests (i.e. inhibition growth, bioluminescence, mutagenic and genotoxic activity) using different living organisms belonging to bacteria (*Aliivibrio fischeri*, *Salmonella typhimurium*), crustacean (*Daphnia magna*) and algae (*Raphidocelis subcapitata*) revealed a marked influence of the UV dose absorbed by the solution during the photolytic processes on the ecotoxic activity.

Novelty

We present for the first time a kinetic and ecotoxicological investigation on the removal with UV₂₅₄-assisted processes of two antiretrovirals (stavudine and zidovudine) from milli-Q water through a microphotoreactor, which allows very fast experimentation with minimal sample volumes. Zidovudine and stavudine are new emerging poor biodegradable microcontaminants detected in STP effluents and surface waters, especially in African countries, due to the highest incidence of HIV-positive people. Recently, the level of zidovudine in Kenian surface waters increased up to three order of magnitude. Moreover, these substances have been demonstrated to exert a carcinogenic activity. For this purpose, the ecotoxicity of solutions was evaluated to assess genotoxicity and mutagenicity.

1
2
3 **Removal of antiretroviral drugs stavudine and zidovudine in water under**
4 **UV₂₅₄ and UV₂₅₄/H₂O₂ processes: quantum yields, kinetics and**
5 **ecotoxicology assessment**
6
7
8
9

10 Danilo Russo^a, Antonietta Siciliano^b, Marco Guida^b, Roberto Andreozzi^a, Nuno M. Reis^d,
11
12 Gianluca Li Puma^{e,‡} and Raffaele Marotta^{a,†}
13
14
15
16
17

18 ^aDipartimento di Ingegneria Chimica, dei Materiali e della Produzione Industriale, Università
19 di Napoli Federico II, p.le V. Tecchio 80, Napoli, Italy.
20
21

22 ^bDipartimento di Biologia, Università di Napoli Federico II, Complesso Universitario Monte
23 Sant'Angelo, via Cinthia 4, Napoli, Italy.
24
25
26

27 ^c Dipartimento di Scienze Chimiche, Università di Napoli Federico II, Complesso
28 Universitario Monte Sant'Angelo, via Cinthia 4, Napoli, Italy.
29
30
31

32 ^dDepartment of Chemical Engineering, University of Bath, Claverton Down, Bath BA2 7AY,
33 UK.
34
35
36

37 ^e Environmental Nanocatalysis & Photoreaction Engineering Department of Chemical
38 Engineering, Loughborough University, Loughborough LE11 3TU, UK.
39
40
41
42

43
44
45 [†] *Corresponding author*: Tel.: +39(0)817682968, fax: +39(0)815936936. E-mail address:
46 rmarotta@unina.it (R. Marotta).
47
48

49
50 [‡] *Corresponding author*: Tel.: +44(0)1509222510, fax: +44(0)1509223923. E-mail address:
51 G.Lipuma@lboro.ac.uk (G. Li Puma).
52
53
54
55
56
57
58
59
60
61
62

ABSTRACT

The concentration of antiretroviral drugs in wastewater treatment plants (WWTP) effluents and surface waters of developed and developing countries, especially in the African region more affected by HIV, has increased significantly in recent years due to their widespread use. The presence of antiretroviral in natural water bodies limits the possibility of reuse of such waters, after traditional disinfection process (i.e. UV, chlorine) for civil and irrigation purposes. The removal of stavudine and zidovudine under UV₂₅₄ or UV₂₅₄/H₂O₂ irradiation was investigated in distilled water. The quantum yield of direct photolysis and the kinetic constant of reaction of hydroxyl radical with the antiretrovirals at different pH have been evaluated. A battery of ecotoxicological tests (i.e. inhibition growth, bioluminescence, mutagenic and genotoxic activity) using different living organisms belonging to bacteria (*Aliivibrio fischeri*, *Salmonella typhimurium*), crustacean (*Daphnia magna*) and algae (*Raphidocelis subcapitata*) revealed a marked influence of the UV dose absorbed by the solution during the photolytic processes on the ecotoxic activity.

Keywords: photodegradation, microreactor, mutagenicity, genotoxicity, water reuse, zidovudine, stavudine, antiretroviral.

1. Introduction

In the last decades, as a result of the widespread availability of pharmaceutical drugs, occurrence, identification, quantification, removal and environmental fate of these emerging contaminants have received significant critical attention [1-3]. Among this new and increasingly growing class of water microcontaminants, the presence of antiretroviral drugs (ARVs) in wastewater and surface water has been the focus of recent research [4-9]. Since their introduction into the market in the early 90s, ARVs have rapidly spread across the world

1 because of their effectiveness in the treatment of the HIV virus [10]. In fact, ARVs inhibit the
2 reverse transcriptase of the HIV virus, repressing viral replication [11]. The most commonly
3 used ARVs include zidovudine (ZDV), stavudine (STV), lamivudine, abacavir and
4 nevirapine which are usually administered as a combination therapy to increase their
5 effectiveness in preventing HIV reproduction [12]. ZDV was the first marketed antiretroviral
6 [12] and is still one of the most widely used. STV is also one of the most common ARVs,
7 despite presenting several side effects, because of its relatively low price [13]. Collectively,
8 ARVs increase the life expectancy of HIV-positive patients, however, significant concerns
9 have been raised about their simultaneous release to the environment [4,5,9]. New concerns
10 are also related to their consumption in the illicit drugs nyaope [14] and whoonga [15]. As a
11 result, STV and ZDV have been often detected in effluents of wastewater treatment plants
12 (WWTPs) and in natural surface water, in Europe and in Africa, at levels of tens of $\text{ng}\cdot\text{L}^{-1}$ up
13 to hundreds of $\text{ng}\cdot\text{L}^{-1}$ (Table 1).

14 In Europe, the main ARVs contamination route of natural waters is through human body
15 excretion and subsequent release in the sewage system [9]. The further presence of ARVs in
16 the effluents of WWTPs and surface water demonstrates the inefficiency of current WWTPs
17 treatment methods. The highest concentrations have been detected in Kenya and South
18 Africa. The levels of ZDV and STV in these African countries, have been shown to be higher
19 in surface water compared to WWTPs effluents, which contrast with the general trend in
20 Europe. The level of ZDV in Kenian surface waters increased up to three order of magnitude
21 during the period 2012 to 2016 [16,18]. Recently, ZDV has also been detected in
22 groundwater [16] which can be probably ascribed to the illicit use and direct spillage in
23 water.

24 It has been reported [7] that ZDV is not completely removed in conventional treatment
25 plants, a conclusion also shown for an aerobic and anaerobic WWTP in Germany [9],

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

although, these authors reported 68% of ZDV removal in different German WWTP with an activated sludge system. Further biological treatment studies performed in synthetic wastewater demonstrated that ZDV is non biodegradable, toxic, and inhibitory to activated sludge bacteria [22]. Higher removals have been reported for STV through activated sludge (> 78%) and biological treatment (> 89%) [7,9]. Even though the reported LC_{50} (Daphnid acute 48 h) are $980 \text{ mg}\cdot\text{L}^{-1}$ and higher than $100 \text{ mg}\cdot\text{L}^{-1}$ for STV and ZDV respectively [23-25], synergistic and mutagenic effects on the aquatic fauna and humans cannot be ruled out. For example, ZDV has been demonstrated to have carcinogenic potential [26]. Advanced oxidation processes (AOPs) have increasingly been proposed as effective tertiary treatments for the removal of biorecalcitrant emerging contaminants [27,28]. Among these, the UV_{254}/H_2O_2 is considered one of the most convenient process since it can be simply applied in existing municipal water treatment plants adopting UV_{254} lamps for water disinfection, such as treatment plants for water reuse and tertiary units in conventional STP [29]. Notably, reclaimed water reuse for irrigation is especially suitable in water stressed areas [30], which often also present the highest incidence of HIV-positive people, such as Central and South Africa. In spite of the apparent effectiveness of AOPs in micropollutants removal, the potential for the formation of highly toxic by-products [31,32] calls for longer treatment times and for the further evaluation of the ecotoxicity of the treated water. In this study the kinetics of ZDV and STV direct photolysis under UV_{254} radiation and in the presence of hydrogen (UV_{254}/H_2O_2) was investigated in order to estimate important photo-kinetic parameters, such as the quantum yields and the second-order kinetic constant of reaction between OH radicals and the compounds, which are necessary for design and retrofitting of water treatment plants. The reaction kinetics were investigated by means of a recent developed methodology which used microcapillary photoreactor systems [33], previously adopted for the investigation of the photolytic kinetics of other micropollutants [34-36]. The

1 use of this new microphotoreactor technology has been shown to be particularly suitable for
2 the study on highly priced, hazardous, or poorly available water contaminants since it allows
3
4 to run the entire experimental campaign using minimal amount of compounds, in this case
5
6 less than 50 mg of ZDV and STV.
7

8
9 The implementation of water reclamation systems and of the environmental risks posed by
10 the effluents, requires comprehensive ecotoxicological assessment on a set of biological tests
11
12 on species at different trophic levels [37]. For this purpose, the three most frequently
13
14 ecotoxicity bioassays in aquatic systems are the assessment on *Aliivibrio fischeri* and
15
16 *Daphnia magna* tests for acute toxicity and the *Raphidocelis subcapitata* test for chronic
17
18 toxicity. Although these target organisms have often been used to assess the impact of
19
20 contaminated water, the main focus of water quality testing should also concern organisms-
21
22 dependent chemical-physical and biological properties of the target molecules. In particular,
23
24 several studies have demonstrated that ARVs differ in genotoxic potency, chromosomal
25
26 damage and aberration types induced *in vitro* and in perinatally exposed mice and infants [38-
27
28 40].
29

30 In consequence, in the present study we investigated the ecotoxicity of untreated and treated
31
32 solutions of ZDV and STV using a battery of ecologically relevant testing species to assess
33
34 the acute and chronic toxicities and genotoxicity and mutagenicity.
35

36 **2. Experimental**

37 *2.1. Materials*

38 Zidovudine (> 99%), stavudine (> 98%), NaOH (≥ 98%), H₂SO₄ (98%), hydrogen peroxide
39
40 (30% in H₂O), acetonitrile (≥ 99.9%), methanol (≥ 99.9%), phosphoric acid (85% in H₂O),
41
42 catalase from *Micrococcus lysodeikticus*, CaCl₂·2H₂O (≥ 99.5%), MgSO₄·7H₂O (≥ 98%),
43
44 NaHCO₃ (≥ 99.5 %), KCl (≥ 99 %), NH₄Cl (≥ 99.9%), MgCl₂·6H₂O (≥ 98%), KH₂PO₄ (≥
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 99%), $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ ($\geq 98\%$), $\text{Na}_2\text{EDTA} \cdot 2\text{H}_2\text{O}$ ($\geq 99.9\%$), H_3BO_3 ($\geq 99\%$), $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (\geq
2 98%), ZnCl_2 ($\geq 99\%$), $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ($\geq 98\%$), $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ ($\geq 98\%$) and $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (\geq
3 98%) were all purchased from Sigma-Aldrich and used as received.
4

5 Reconstitution solution, osmotic adjusting solution (OAS) and diluent (NaCl 2%) were the
6 reagents used for the *Aliivibrio fischeri* toxicity test (Strategic diagnostics Inc. SDI). All the
7 reacting solutions were prepared adding the contaminants and hydrogen peroxide to milliQ
8 water. When necessary, pH was adjusted by using dilute aqueous solutions of NaOH and
9 H_2SO_4 .
10
11
12
13
14
15
16
17
18
19
20
21

22 2.2. Photolytic treatments

23 A microcapillary film (MCF) array photoreactor was used to perform the UV_{254} photolysis
24 and $\text{UV}_{254}/\text{H}_2\text{O}_2$ experiments. A detailed description and a scheme of the reactor can be
25 found elsewhere [33,34]. Briefly, the polymeric film microreactor consists of ten tubular
26 microcapillaries (Lamina Dielectrics Ltd, Billingshurst, West Sussex, UK) with a mean
27 hydraulic diameter of 195 μm and an average optical path length of 152 μm . The MFC was
28 fed by means of a syringe pump (HA Harvard Apparatus PHD Ultra) and wrapped around a
29 germicidal lamp (Germicidal G8T5, Ge Lighting) emitting at 254 nm. The nominal lamp
30 power could be varied from 8.0 to 4.5 W with the use of a switch power supplier. The emitted
31 photon fluxes per unit volume (I_0/V) were estimated by hydrogen peroxide actinometry
32 [41,42] and were $1.92 \cdot 10^{-2} \text{ ein} \cdot \text{s}^{-1} \cdot \text{L}^{-1}$ and $1.27 \cdot 10^{-2} \text{ ein} \cdot \text{s}^{-1} \cdot \text{L}^{-1}$, respectively. The residence
33 time (space time) through the MFC was varied changing the length of the film exposed to the
34 light. Samples were collected at the outlet of the MFC after reaching steady state conditions,
35 and rapidly analyzed by HPLC. Negligible temperature differences ($\sim 25^\circ\text{C}$) between the
36 inlet and the outlet samples were found in all the experiments. All the experimental runs were
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 carried out in duplicate. The entire experimental campaign (> 50 runs) was carried out using
2 about 2500 ml of distilled water.
3

4 In order to produce relatively larger volumes of treated solutions, necessary to run the
5 ecotoxicology tests, experiments were also carried out in a thermostated (25 °C) stirred glass
6 batch annular photoreactor with a volume of $4.8 \cdot 10^{-1}$ L, housing a low pressure mercury lamp
7 emitting at 254 nm (Helios Italquartz, HGL10T5L, 17 W) in the centre axis of the annulus. In
8 both photoreactor devices, ZDV and STV solutions at an initial concentration of $4.5 \text{ mg} \cdot \text{L}^{-1}$
9 and $4.35 \text{ mg} \cdot \text{L}^{-1}$ respectively, were treated without and with the addition of hydrogen
10 peroxide (molar ratio $\text{H}_2\text{O}_2/\text{ARV} = 100$) and with a UV_{254} dose corresponding to treatment
11 times or space times necessary to achieve a 45% and 90% conversion of the antiretrovirals
12 and for space times double those needed to achieve a complete conversion. The UV_{254} dose
13 ($\text{mJ} \cdot \text{cm}^{-2}$) was calculated as the average photon fluence rate multiplied by the treatment time
14 (s). The average photon fluence rate emitted by the UV lamp at 254 nm was $4.7 \text{ mW} \cdot \text{cm}^{-2}$
15 (UVC DELTA OHM radiometer). The experimental device was described elsewhere [36].
16 Catalase enzyme was also added to both treated and untreated solutions to determine the
17 ecotoxicological effects without the interference given by the presence of H_2O_2 residuals.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 2.3. Analytical methods

43 ZDV, STV, and hydrogen peroxide concentrations were measured by HPLC (1100 Agilent)
44 equipped with a Gemini 5u C6-Phenyl (260 x 4.60 mm) (Phenomenex) column. An isocratic
45 method was used for the simultaneous quantification of ZDV and hydrogen peroxide with
46 mobile phase ($0.8 \text{ mL} \cdot \text{min}^{-1}$) made of 93% aqueous phosphoric acid (10 mM) and 7%
47 acetonitrile. Under these analytical conditions the retention times of hydrogen peroxide and
48 ZDV were 4.1 and 13.7 min, respectively. The mobile phase was changed to 80% water and
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

20% methanol for the simultaneous identification of H₂O₂ and STV with retention times of 3.7 and 10 min, respectively. The signals were acquired at 266 nm.

The molar absorption coefficients of STV and ZDV were estimated using a Perkin Elmer UV/VIS spectrometer (mod. Lambda 35).

2.4. *Ecotoxicological methods*

2.4.1. *Toxicity test with Daphnia magna*

The test was conducted in accordance with ISO 6341 [43]. Every test was performed in quadruplicate with four control groups. Briefly, neonates aged less than 24 hours were separated into four groups and exposed to different concentrations of untreated and treated solutions of ZDV and STV. All tests were carried out at constant temperature (20±2 °C) and in darkness and organisms were not fed during the experiments. After 48 h exposure, daphnids that were not able to swim within 15 seconds under gentle agitation were considered to be immobilized.

2.4.2. *Bacteria toxicity test*

The Microtox® SPT procedure [44] was used to evaluate the acute toxicity of the samples using as endpoint the bioluminescence inhibition of the naturally emitted by *A. fischeri* (strain NRRL-B-11177) after a contact time of 30 min with the test sample. The samples were serially diluted to a series of four concentrations, then a volume of 10 µL of reconstituted bacterial reagent was added to dilutions series of samples. The emission of bioluminescence was recorded after 30 min of contact time with the bacteria at 15±2 °C.

2.4.3. Algal growth inhibition test with *Raphidocelis subcapitata*

The growth inhibition test was assessed following the ISO 8692:2012 standard procedure [45]. Exponentially growing algae (10^4 cell·mL⁻¹) were exposed to various concentrations of the test samples in six replicates over a period of 72 h under defined conditions, as described elsewhere [46]. Growth and inhibition were quantified from measurements of the algal biomass density (cell counts) as a function of time. The specific growth rate of *R. subcapitata* in each replicate culture was calculated from the logarithmic increase in cell density in the intervals from 0 to 72 h using the following equation: $\mu = \frac{\ln N_n - \ln N_0}{t_n - t_0}$, where N_0 is the cell concentration at $t = 0$, N_n the final cell concentration after 72 h of exposure, t_0 the time of start measurement, and t_n the time of last measurement (hours from start). The results were expressed as the mean (\pm standard deviation) of the percentage inhibition of the cell growth (% I) of the sample compared with the negative control ($p \leq 0.05$).

2.4.4. Mutagenicity assay with *Salmonella typhimurium*

The Muta-Chromoplate kit was used to evaluate the mutagenicity [47]. The fluctuation tests were performed using *S. typhimurium* strains TA100 and observing the potential *his*- reverse mutation after exposure to mutagens [48].

Bacteria cultures, grown overnight and reaching the exponential growth phase, were exposed for 5 days at 37° C to different samples concentrations, in a liquid medium into 96-well microtiter plates. After this period, the positives samples wells that turned yellow were counted, while the purple wells were considered as negatives. The reversion of mutants (*his*+) exhibited the yellow color due to the acidification of the test medium resulting from the growth of reverse mutants. The number of *his*+ revertant colonies in each sample was determined as a mean value of the three plates. The results were expressed as a mutagenicity ratio (MR), i.e. the ratio of the number of *S. typhimurium* revertants grown in the presence of

1 the tested sample to the number of spontaneously appeared revertants. The sample was
2 considered mutagenic when $MR \geq 2$ [49]. χ -square analysis was used for statistical evaluation
3 of the treated plates versus the control plates.
4
5
6
7
8

9 2.4.5. Genotoxicity assay with *Salmonella typhimurium*

10 The umu test [50] was performed according to standard procedure [51], which was developed
11 for the detection of genotoxic materials that cause DNA cell damage. In this assay, a
12 modified strain of *S. typhimurium* TA1535/pSK 1002 bacteria was used, whereby a β -
13 galactosidase gene was linked to SOS-DNA response. Bacterial cultures were grown
14 overnight at 37 °C and then diluted in TGA medium (Tryptone-Glucose-Ampicillin medium)
15 until the cells entered the logarithmic growth phase. The cells were then exposed to the test
16 samples for 2 hours. The induction of genotoxicity (expressed as β -galactosidase activity)
17 was determined colorimetrically at 420 nm after adding o-nitrophenyl galactopyranoside to
18 the samples. Growth was measured as the absorbance at a wavelength of 600 nm. The result
19 was calculated as an induction ratio, $IR = (1/G) \cdot US$, where G was the growth and US the
20 relative enzyme activity. The sample was considered genotoxic when IR was greater than 1.5.
21
22 The significance of the differences between the mean values of different tests and controls
23 was assessed by Student's test and analysis of variance (ANOVA) with a 0.05 significance
24 level. Moreover, post-hoc analysis were carried out by Tukey's test [52].
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 3. Results and Discussion

50 3.1. Absorbance spectra

51 The absorbance spectra of ZDV (Fig. 1a) and STV (Fig. 1b) at pH 4.0, 6.5 and 8.0 showed
52 invariance in the pH range from 4.0 to 8.0 for ZDV and from 6.5- to 8.0 for STV. Since pH
53 did not affect ZDV and STV degradation kinetics in the pH ranges 4.0-8.0 and 6.0-8.0,
54
55
56
57
58
59
60
61
62
63
64
65

1 respectively, the reaction kinetics were investigated in the slightly acidic to alkaline pH range
 2 from 6.0 to 8.0, which also is more environmentally relevant. The estimated molar absorption
 3 coefficients at 254 nm are summarized in Table 2.
 4
 5

6 7 8 9 10 3.2. ZDV and STV direct photolysis and quantum yield estimation

11 The quantum yield (ϕ_i^{254}) of direct photolysis at 254 nm was determined with different sets
 12 of experimental runs carried out in the MCF varying the ARV initial concentration, the pH
 13 and the lamp power (Table 3). The degradation of the generic ARV by direct photolysis
 14 follows the mass balance (eq. 1):
 15
 16
 17
 18
 19
 20

$$21 \frac{dC_i}{dt} = -\frac{I_0}{V} \cdot \phi_i^{254} \cdot (1 - \exp(-2.3 \cdot l \cdot \varepsilon_i^{254} \cdot C_i)) \quad (1)$$

22 where C_i is the concentration of ARV, I_0/V is the photon flux per unit volume, l is the
 23 average optical length of the reactor (see *Photolytic treatments section*), and ε_i^{254} the molar
 24 absorption coefficient of the ARV species (Table 2). A Matlab optimization routine based on
 25 the Runge-Kutta method was adopted to determine the value of ϕ_i^{254} which minimized the
 26 objective function (optimisation mode):
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36

$$37 \sum_j^m \sum_i^n (y_{j,i} - c_{j,i})^2 \quad (2)$$


38 where c and y are the experimental and calculated concentrations at different reaction times
 39 (n) and experimental runs (m). Table 3 shows the estimated quantum yields and the 97%
 40 interval of confidence of ZDV and STV. The results corroborate with the previous photolytic
 41 decomposition investigation [53], that reported higher sensitivity of ZDV to UVA light
 42 compared to STV, with a fluorescent irradiation source in the wavelength spectra from 320
 43 nm to 400 nm.
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54

55 All the adopted experimental runs used for the kinetic modeling along with the experimental
 56 conditions and the average standard deviation are reported in Table 4 whereas Fig. 2 (a-d)
 57
 58
 59
 60
 61
 62
 63
 64
 65

show representative examples of the comparison between the experimental and the calculated profiles for ARV degradation by direct photolysis. As reported, the model was validated simulating simultaneously different experimental runs not included in the optimization routine (simulation mode, Fig. 2e-f).

3.3. Kinetic modeling of UV₂₅₄/H₂O₂ process

The second order rate constant $k_{OH,i}$ of the reaction between ZDV and STV with hydroxyl radicals was estimated from a set of experimental runs carried out in the MCF at varying pH, lamp power and $C_{H_2O_2}/C_i$ molar ratio (Table 5). They were modeled according to the following simplified reaction scheme:



Hydroxyl radicals formed by direct photolysis of hydrogen peroxide under UV₂₅₄ radiation (r_1) attack the ARV (r_3), the P and S pseudo by-products formed (r_4 - r_5), and the oxidant hydrogen peroxide which generates hydroperoxyl radicals (r_6). The latter recombine according to the termination reaction (r_7) to form H₂O₂. The simultaneous photolysis of ARV is considered in reaction (r_2). Assuming that hydroxyl radicals attack both substrate and the first generation of chemical intermediates with the same rate constant $k_{OH,i}$ and that at any

1 reaction time the sum of the unconverted substrate (C_i) and its by-products ($C_{P,S}$)
 2 concentration is equal to the initial antiretroviral concentration (C_o):
 3
 4

$$5 \quad C_o = C_i + C_{P,S}$$

6
 7
 8 the concentrations of hydroxyl and hydroperoxyl radicals, under the steady state
 9 approximation equal:
 10

$$11 \quad [HO^*]_{ss} = \frac{2F_{H_2O_2}}{k_{OH,i} \cdot C_o + k_H \cdot C_{H_2O_2}} \quad (3)$$

$$12 \quad [HO_2^*]_{ss}^2 = \frac{F_{H_2O_2} \cdot k_H \cdot C_{H_2O_2}}{k_t(k_{OH,i} \cdot C_o + k_H \cdot C_{H_2O_2})} \quad (4)$$

13
 14
 15
 16
 17
 18
 19
 20
 21
 22 where

$$23 \quad F_{H_2O_2} = \frac{I_0}{V} \cdot \phi_{H_2O_2}^{254} \cdot \left(1 - \exp\left(-2.3 \cdot l \cdot (\varepsilon_{H_2O_2}^{254} \cdot C_{H_2O_2} + \varepsilon_i^{254} \cdot C_i)\right)\right) \cdot \frac{\varepsilon_{H_2O_2}^{254} \cdot C_{H_2O_2}}{\varepsilon_{H_2O_2}^{254} \cdot C_{H_2O_2} + \varepsilon_i^{254} \cdot C_i} \quad (5)$$

24
 25
 26
 27
 28 and the concentration of H_2O_2 and ARVs versus time can be determined by solving the
 29 following material balance:
 30

$$31 \quad \frac{dC_{H_2O_2}}{dt} = -F_{H_2O_2} - \frac{k_H \cdot C_{H_2O_2} \cdot F_{H_2O_2}}{k_{OH,i} \cdot C_o + k_H \cdot C_{H_2O_2}} \quad (6)$$

$$32 \quad \frac{dC_i}{dt} = -F_i - \frac{2k_{OH,i} \cdot C_i \cdot F_{H_2O_2}}{k_{OH,i} \cdot C_o + k_H \cdot C_{H_2O_2}} \quad (7)$$

33
 34
 35
 36
 37
 38
 39
 40
 41
 42 where

$$43 \quad F_i = \frac{I_0}{V} \cdot \phi_i^{254} \cdot \left(1 - \exp\left(-2.3 \cdot l \cdot (\varepsilon_{H_2O_2}^{254} \cdot C_{H_2O_2} + \varepsilon_i^{254} \cdot C_i)\right)\right) \cdot \frac{\varepsilon_{ARV}^{254} \cdot C_{H_2O_2}}{\varepsilon_{H_2O_2}^{254} \cdot C_{H_2O_2} + \varepsilon_i^{254} \cdot C_i} \quad (8)$$

44
 45
 46
 47
 48 Similarly to the previous analysis, the equations (6-7) were solved by means of an
 49 optimization routine to minimize the objective function
 50

$$51 \quad \sum_{h=1}^2 \sum_j^m \sum_i^n (y_{j,i} - c_{j,i})^2 \quad (9)$$

slightly modified to account for the number of reacting species (h). Selected experimental runs were not included in the optimization procedure to validate the kinetic model without further adjustment of the estimated kinetic parameter $k_{OH,i}$. All the runs adopted, along with their average standard deviation, are summarized in Table 5. Figure 3 shows the comparisons between the experimental and calculated data, both in optimization and simulation modes. The second order rate constant of ARVs investigated with hydroxyl radicals is shown in Table 6. The kinetic constant of reaction of hydroxyl radical with ZDV, determined by the competition kinetics method, in literature varies over a wide range: $(1.3 \pm 0.026) \cdot 10^{10} \text{ M}^{-1} \cdot \text{s}^{-1}$ determined using para-chlorobenzoic acid as reference compound [57] and $(5.73 \pm 0.76) \cdot 10^9 \text{ M}^{-1} \cdot \text{s}^{-1}$ determined using acetophenone as reference substance [17]. These values are significant higher than the results obtained in this study. One possible explanation for this discrepancy is that the competition kinetics method is only reliable if the contribution of direct photolysis of the investigated compound is absent or negligible. In the absence of that, the $k_{OH,i}$ value tends to be overestimated ($k'_{OH,i}$) since it also accounts for the contribution of direct photolysis to the degradation. To further clarify this aspect, the competition kinetic method using benzoic acid (BA) as reference compound [58] was also used to estimate the second order rate constant of hydroxyl radicals with ARVs. According to this method the $k'_{OH,i}$ value can be calculated according to (10):

$$\ln \frac{C_i}{C_o} = \frac{k'_{OH,i}}{k_{OH,BA}} \ln \frac{C_{BA}}{C_{BA_o}} \quad (10)$$

where C_{BA} and C_{BA_o} are the unconverted and initial concentration of benzoic acid, and $k_{OH,BA}$ the kinetic constant of reaction between benzoic acid and hydroxyl radical ($5.9 \cdot 10^9 \text{ M}^{-1} \cdot \text{s}^{-1}$ [55]). Duplicate experimental runs were carried out under the following conditions: $C_{ZDV_o} = 2.46 \cdot 10^{-5} \text{ M}$, $C_{BA_o} = 2.62 \cdot 10^{-5} \text{ M}$, $C_{H_2O_2_o} = 1.41 \cdot 10^{-3} \text{ M}$, and $C_{STV_o} = 2.21 \cdot 10^{-5} \text{ M}$, $C_{BA_o} = 2.15 \cdot 10^{-5} \text{ M}$, $C_{H_2O_2_o} = 9.8 \cdot 10^{-4} \text{ M}$. Plotting $\ln C_i/C_o$ vs $\ln C_{BA}/C_{BA_o}$, $k'_{OH,i}$ was estimated as

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

$6.39 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ and $5.33 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ for ZDV and STV, respectively. As expected both values overestimate the results reported in Table 6. Moreover, the difference is higher for ZDV, in agreement with the higher significance of direct photolysis in the degradation of this compound compared to STV. It is also important to notice that the $k'_{OH,ZDV}$ value estimated in the present study, adopting the competition kinetics method, was within those reported in the literature [17,57].

3.4. Ecotoxicological assessment

The inhibition of *A. fischeri* luminescence and *D. magna* immobility was not observed on untreated and UV₂₅₄ or UV₂₅₄/H₂O₂ treated solutions (data not shown). The growth inhibition of algae *R. subcapitata* exposed to ZDV and STV containing solutions before and during the photolytic processes (UV₂₅₄ or UV₂₅₄/H₂O₂) is shown in Figures 4a-f. A marked different trend in effects on algae growth according to the dose–response correlation was observed. ZDV inhibited slightly algal growth, while STV had insignificant effect (less than 20%). The toxicity of the treated solutions in both processes slightly increased increasing the UV dose, also for treatment times corresponding to the complete removal of the antiretroviral drugs. In particular, the inhibited algal growth for the ZDV treated solutions increased by 36% and 44% when the conversion of ZDV was 90% and 100% respectively in both UV₂₅₄ and UV₂₅₄/H₂O₂ treatments (Fig. 4a). The same increasing trend of ecotoxicity was observed with the STV treated solutions, although the increase in ecotoxicity to *R. subcapitata* was less significant in comparison to the ZDV treated solutions. The inhibition was 20% and 30% when the STV conversions was 90% and 100% respectively (Fig. 4b).

The antiretroviral drugs showed marked algae growth activity when the dilution factor of untreated and treated solutions was increased from 1:10 (Figs. 4c-d) to 1:100 (Figs. 4e-f), which appeared to stimulate algal growth with a statistically significant extent. This

1 uncommon “apparently beneficial effect” observed at low doses, known as hormesis, has
2 been previously reported for some bioindicators such as crustaceans [59,60] and plants and
3 algae [61] in the presence of nitrogen-containing organic molecules (trinitrotoluene, triazine
4 herbicides, etc.) such as are STV and ZDV.
5
6

7
8
9 Tables 7 and 8 summarize the mutagenicity and genotoxicity results. Both antiretrovirals
10 were not able to determine a significant SOS system induction, while variability among the
11 mutagenic responses was observed in the Salmonella mutagenicity assay (threshold value:
12 2.0). The mutagenicity results indicated that (i) potential mutagenic degradation intermediates
13 could have been formed at significant levels during the photolysis of ZDV than STV, (ii) the
14 mutagenicity of the samples further increased at increasing UV_{254} doses and (iii) the
15 UV_{254}/H_2O_2 treatment produced less mutagenic intermediate products than those formed
16 during UV_{254} photolysis. Residual mutagenic activity was also observed on the ZDV samples
17 treated by UV_{254} photolysis for conversions higher than 90%, after a 1:1000 dilution (Table
18 7). It is useful to point out that the concentration of ZDV in these samples was of the same
19 order of magnitude as the values detected in African surface water (Table 1).
20
21

22 During the UV_{254}/H_2O_2 process it was observed a slight increase of revertants followed by
23 disappearance of revertants at the highest UV_{254} exposure.
24
25

26 Investigations of the genotoxic endpoints demonstrated that the untreated solutions could not
27 be classified as genotoxic since the induction ratio was below the threshold value of 1.5
28 (Tables 7-8). On the contrary, a statistically significant increase in umuC induction was
29 recorded for ZDV undiluted solutions for UV_{254} doses corresponding to ZDV conversions of
30 45% and 90% for both UV_{254} and UV_{254}/H_2O_2 processes (Table 7). The genotoxic activity
31 showed a decreasing trend as the dilution factor of the solutions increased.
32
33

34 STV genotoxic data demonstrated (Table 8) that only the treated solutions were able to
35 induce a significant SOS response with IF higher than 2 also observed for the highest dilution
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 factor (1:1000), which corresponds to an initial concentration of STV of $4.3 \mu\text{g}\cdot\text{L}^{-1}$. The
2 results collectively may indicate that some photoproducts generated during the photolytic
3 processes could have genotoxic activity. However, genotoxic activity was not observed in the
4 solutions treated for prolonged treatment times, thus suggesting that the genotoxic
5 transformation products observed in earlier times might have further evolved to non-
6 genotoxic metabolites.
7
8
9
10
11
12
13
14
15
16

17 **4. Conclusions**

18 The removal of stavudine and zidovudine by UV_{254} radiation without and with hydrogen
19 peroxide was investigated in a microcapillary film photoreactor using minimal quantities of
20 water samples. Higher UV_{254} -photolysis quantum yields were observed for zidovudine,
21 $(2.357 \pm 0.0589) \cdot 10^{-2} \text{ mol}\cdot\text{ein}^{-1}$ in the pH range from 4.0 to 8.0, while stavudine quantum
22 yield was 28-fold lower $(8.34 \pm 0.334) \cdot 10^{-4} \text{ mol}\cdot\text{ein}^{-1}$ in the pH range from 6.0 to 8.0. The
23 second-order rate constant of reaction with hydroxyl radicals was $(9.98 \pm 0.68) \cdot 10^8 \text{ M}^{-1}\cdot\text{s}^{-1}$
24 (pH range 4.0 – 8.0) for zidovudine and $(2.03 \pm 0.18) \cdot 10^9 \text{ M}^{-1}\cdot\text{s}^{-1}$ (pH: 6.0 – 8.0) for
25 stavudine. The well known ecotoxicological tests using *A. fischeri* and *D. magna* as
26 bioindicators did not evidenced acute or chronic effects. A hormetic effect was observed for
27 the first time in *R. subcapitata* for ZDV and STV treated solutions at different UV_{254} doses
28 after a dilution from 1:10 to 1:100.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 On the contrary, specific tests using *Salmonella t.* revealed mutagenic and genotoxic activity
48 of the ZDV and STV samples also at high dilution factors depending on the type of the
49 photolytic treatment and substrate conversion.
50
51
52
53

54 Generally, UV_{254} photolysis in the presence of hydrogen peroxide reduces the
55 ecotoxicological risk associated to direct photolysis of the aqueous solutions containing the
56 antiretrovirals, but for this purpose UV_{254} doses ($\geq 2000 \text{ mJ}\cdot\text{cm}^{-2}$), significantly higher than
57
58
59
60
61
62
63
64
65

1 the levels suggested for the water UV disinfection processes ($50 - 200 \text{ mJ}\cdot\text{cm}^{-2}$) are
2 necessary. This study pointed out the critical importance of selecting suitable bioindicators
3 depending on chemical and biological properties of the selected xenobiotics detected in STP
4 effluents and in surface waters.
5
6
7
8
9

10 **Acknowledgements**

11
12 The Authors are grateful to ERASMUS-Mobility Student Program, and to Ing. Giulio Di
13 Costanzo for his precious support during the experimental campaign.
14
15
16
17
18
19
20
21

22 **References**

- 23
24 [1] B. Petrie, R. Barden, B. Kasprzyk-Hordern, A review on emerging contaminants in
25 wastewater and the environment: current knowledge, understudies areas and
26 recommendations for future monitoring, *Wat. Res.* 72 (2015) 3–27.
27
28
29 [2] T. Deblonde, C. Cossu-Leguille, P. Hartemann, Emerging pollutants in wastewater: a
30 review of the literature, *Int. J. Hyg. Envir. Heal.* 214(6) (2011) 442–448.
31
32 [3] N. Bolong, A.F. Ismail, M.R. Salim, T. Matsuura, A review of the effects of emerging
33 contaminants in wastewater and options for their removal, *Desalination* 239(1-3) (2009) 229–
34 246.
35
36 [4] S. Jain, P. Kumar, R.K. Vyas, P. Pandit, A.K. Dalai, Occurrence and removal of antiviral
37 drugs in environment: a review. *Water Air Soil Poll.* 224(2) (2013) 1410–1428.
38
39 [5] E. Ngumba, A. Gachanja, T. Tuhkanen, Occurrence of selected antibiotics and
40 antiretroviral drugs in Nairobi River Basin, Kenya, *Sci. Total Environ.* 539 (2016a) 206–213.
41
42 [6] E. Ngumba, P. Kosunen, A. Gachanja, T. Tuhkanen, A multiresidue analytical method for
43 trace level determination of antibiotics and antiretroviral drugs in wastewater and surface
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 water using SPE-LC-MS/MS and matrix-matched standards. *Anal. Methods* 8 (2016b) 6720–
2 6729.
3

4 [7] T.P. Wood, C.S.J. Duvenage, E. Rohwer, The occurrence of anti-retroviral compounds
5 used for HIV treatment in South African surface water, *Environ. Pollut.* 199 (2015) 235–243.
6

7 [8] X. Peng, C. Wang, K. Zhang, Z. Wang, Q. Huang, Y. Yu, W. Ou, Profile and behaviour
8 of antiviral drugs in aquatic environments of the Pearl River Delta, China, *Sci. Total Environ.*
9 466–467 (2014) 755–761.
10

11 [9] C. Prasse, M.P. Schlusener, R. Schulz, T.A. Ternes, Antiviral drugs in wastewater and
12 surface water: a new pharmaceutical class of environmental relevance? *Environ. Sci.*
13 *Technol.* 44 (2010) 1728–1735.
14

15 [10] G. Camponeschi, J. Fast, M. Gauval, K. Guerra, M. Moore, S. Ravinutala, D. Ripin, V.
16 Shepel, An overview of the antiretroviral market, *Curr. Opin. HIV AIDS*, 8(6) (2013) 535–
17 543.
18

19 [11] M.J.M. Hitchcock, 2',3'-Didehydro-2',3'-dideoxythymidine (D4T), an anti-HIV agent,
20 *Antivir. Chem. Chemother.* 2(3) (1991) 125–132.
21

22 [12] G. Kumari, R.K. Singh, Highly active antiretroviral therapy for treatment of HIV/AIDS
23 patients: current status and future prospects and the Indian scenario, *HIV AIDS Rev.* 11
24 (2012) 5–14.
25

26 [13] J.S.F. Lee, L.S. Teyssier, B.D. Nguimfack, I.J. Collins, M. Lallemand, J. Perriens, J.
27 Moatti, J., An analysis of volumes, prices and pricing trends of the pediatric antiretroviral
28 market in developing countries from 2004 to 2012, *BMC Pediatrics* 16(41) (2016) 1–8.
29

30 [14] A.A. Khine, K.E. Mokwena, M. Huma, L. Fernandes, Identifying the composition of
31 street drug nyaope using two different mass spectrometer methods, *Afr. J. Drug Alcohol*
32 *Stud.* 14(1) (2015) 49–56.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [15] D.J. Grelotti, E.F. Closson, J.A. Smit, Z. Mabude, L.T. Matthews, S.A. Safren, D.R. Bangsberg, M.J. Mimiaga, Whoonga: potential recreational use of HIV antiretroviral medication in South Africa, *AIDS Behav.* 18(3) (2014) 511–518.
- [16] K.O. K'oreje, L. Vergeynst, D. Ombaka, P. De Wispelaere, M. Okoth, H. Van Langenhove, K. Demeestere, Occurrence patterns of pharmaceutical residues in wastewater, surface water and groundwater of Nairobi and Kisumu city, Kenya, *Chemosphere* 149 (2016) 238–244.
- [17] C. Zhou, J. Chen, Q. Xie, X. Wei, Y. Zhang, Z. Fu, Photolysis of three antiviral drugs acyclovir, zidovudine and lamivudine in surface freshwater and seawater. *Chemosphere* 138 (2015) 792–797.
- [18] K.O. K'oreje, K. Demeestere, P. De Wispelaere, L. Vergeynst, J. Dewulf, H. Van Langenhove, From multi-residue screening to target analysis of pharmaceuticals in water: Development of a new approach based on magnetic sector mass spectrometry and application in the Nairobi River basin, Kenya, *Sci. Total Environ.* 437 (2012) 153–164.
- [19] J. Funke, C. Prasse, T.A. Ternes, Identification of transformation products of antiviral drugs formed during biological wastewater treatment and their occurrence in the urban water cycle. *Wat. Res.* 98 (2016) 75–83.
- [20] Y. Aminot, X. Litrico, M. Chambolle, C. Arnaud, P. Pardon, H. Budzinski, Development and application of a multi-residue method, for the determination of 53 pharmaceuticals in water, sediment, and suspended solids using liquid chromatography-tandem mass spectrometry. *Anal. Bioanal. Chem.* 407 (2015) 8585–8604.
- [21] A.L.L. Silva, R. Cristofolletti, S. Storpirtis, V.D. Sousa, H.E. Junginger, V.P. Shah, S. Stavchansky, J.B. Dressman, D.M. Barends, Biowaiver monographs for immediate-release solid oral dosage forms: stavudine, *J. Pharm. Sci.* 101 (2012) 10–15.

1 [22] M. Vankova, Biodegradability analysis of pharmaceuticals used in developing countries;
2 screening with OxiTopC-110, Tampere University of Technology, Finland, (p.73), 2010.

3
4 [23] Roche, 2007. Pharmaceuticals sustainability database.
5
6 [http://www.roche.com/sustainability/what_we_do/for_communities_and_environment/enviro](http://www.roche.com/sustainability/what_we_do/for_communities_and_environment/environment/safety_data_sheets-row.htm)
7 [nment/safety_data_sheets-row.htm](http://www.roche.com/sustainability/what_we_do/for_communities_and_environment/environment/safety_data_sheets-row.htm) (accessed 12.01.17).
8
9

10 [24] GSK, 2007. Materials safety data sheets from GlaxoSmithKline (GSK).
11
12 <http://www.msds-gsk.com/Default.aspx> (accessed 12.01.17).
13
14

15 [25] H. Sanderson, M. Thomsen, Comparative analysis of pharmaceuticals versus industrial
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

chemicals acute aquatic toxicity classification according to the United Nations classification system for chemicals. Assessment of the (Q)SAR predictability of pharmaceuticals acute toxicity and their predominant acute mode-of-action, *Toxicol. Lett.* 187 (2009) 84–93.

[26] P. Bottoni, S. Caroli, A.B. Caracciolo, Pharmaceuticals as priority water contaminants, *Toxicol. Environ. Chem.* 92(3) (2010) 549–565.

[27] B.A. Wols, C.H.M. Hofman-Caris, Review of photochemical reaction constants of organic micropollutants required for UV advanced oxidation processes in water, *Wat. Res.* 46 (2012) 2815–2827.

[28] M.A. Oturan, J. Aaron, Advanced oxidation processes in water/wastewater treatment: principles and applications. A review, *Crit. Rev. Env. Sci. Tech.* 44(23) (2014) 2577–2641.

[29] I.H. Kim, N. Yamashita, Y. Kato, H. Tanaka, Discussion of the application of UV/H₂O₂, O₃ and O₃/UV processes as technologies for sewage reuse considering the removal of pharmaceuticals and personal care products, *Water Sci. Technol.* 59(5) (2009) 945–955.

[30] A. Y. Hoekstra, Water scarcity challenges to business, *Nat. Clim. Chang.* 4 (2014) 318–320.

[31] F. Yuan, C. Hu, X. Hu, D. Wei, Y. Chen, J. Qu, Photodegradation and toxicity changes of antibiotics in UV and UV/H₂O₂ process, *J. Haz. Mat.* 185 (2011) 1256–1263.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [32] O. Rozas, C. Vidal, C. Baeza, W.F. Jardim, A. Rossner, H.D. Mansilla, Organic micropollutants (OMPs) in natural waters: oxidation by UV/H₂O₂ treatment and toxicity assessment, *Wat. Res.* 98 (2016) 109–118.
- [33] N.M. Reis, G. Li Puma, Novel microfluidics approach for extremely fast and efficient photochemical transformations in fluoropolymer microcapillary films, *Chem. Commun.* 51 (2015) 8414–8417.
- [34] D. Russo, D. Spasiano, M. Vaccaro, R. Andreozzi, G. Li Puma, N.M. Reis, R. Marotta, Direct photolysis of benzoylecgonine under UV irradiation at 254 nm in a continuous flow microcapillary array photoreactor, *Chem. Eng. J.* 283 (2016a) 243–250.
- [35] D. Russo, D. Spasiano, M. Vaccaro, K.H. Cochran, S.D. Richardson, R. Andreozzi, G. Li Puma, N.M. Reis, R. Marotta, Investigation on the removal of the major cocaine metabolite (benzoylecgonine) in water matrices by UV/H₂O₂ process by using a flow microcapillary film array photoreactor as an efficient experimental tool, *Wat. Res.* 89 (2016b) 375–383.
- [36] D. Spasiano, D. Russo, M. Vaccaro, A. Siciliano, R. Marotta, M. Guida, N.M., Reis, G. Li Puma, R. Andreozzi, Removal of benzoylecgonine from water matrices through UV₂₅₄/H₂O₂ process: reaction kinetic modelling, ecotoxicity and genotoxicity assessment, *J. Haz. Mat.* 318 (2016) 515–525.
- [37] I.C. Eom, C. Rast, A.M. Veber, P. Vasseur, Ecotoxicity of a polycyclic aromatic hydrocarbon (PAH)-contaminated soil, *Ecotox. Environ. Safe.* 67 (2007) 190–205.
- [38] O.A. Olivero, G.M. Shearer, C.A. Chougnet, A.A. Kovacs, R. Baker, A.M. Stek, M.M. Khoury, M.C. Poirier, Incorporation of zidovudine into cord blood DNA of infants and peripheral blood DNA of their HIV-1-positive mothers, *Ann. N. Y. Acad. Sci.* 918 (2000) 262–268.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [39] O.A. Olivero, Mechanisms of genotoxicity of nucleoside reverse transcriptase inhibitors Environ. Mol. Mutagen. 48 (2007) 215–223.
- [40] A. Dutra, E. Pak, S. Wincovitch, K. John, M.C. Poirier, O.A. Olivero, Nuclear bud formation: a novel manifestation of zidovudine genotoxicity, Cytogenet. Genome Res. 128 (2010) 105–110.
- [41] I. Nicole, J. De Laat, M. Doré, J.P. Duguet, C. Bonnel, Use of UV radiation in water treatment: measurement of photonic flux by hydrogen peroxide actinometry, Wat. Res. 24 (1990) 157–168.
- [42] S. Goldstein, D. Aschengrau, Y. Diamant, J. Rabani, Photolysis of aqueous H₂O₂: quantum yield and applications for polychromatic UV actinometry in photoreactors, Env. Sci. Tech. 41 (2007) 7486–7490.
- [43] ISO 6341:2012. Water quality: determination of the inhibition of the mobility of *Daphnia magna* Straus (Cladocera, Crustacea)—acute toxicity test.
- [44] ISO 11348-3:2007. Water quality -- Determination of the inhibitory effect of water samples on the light emission of *Aliivibrio fischeri* (Luminescent bacteria test) -- Part 3: Method using freeze-dried bacteria.
- [45] ISO 8692:2012. Water quality—Fresh water algal growth inhibition test with unicellular green algae.
- [46] M. Guida, M. Inglese, S. Meric, A multi-battery toxicity investigation on fungicides, Desalination 226(1–3) (2008) 262–270.
- [47] B.N. Ames, J. McCann, E. Yamasaki, Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test, Mutat. Res. 31(6) (1975) 347–364.
- [48] EBPI, The Muta-chromoPlate Kit S-9 version 3.1. In E. B. P. Inc. (Ed.) 2005.


- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [49] K. Piekarska, J. Karpinska-Smulikowska, Mutagenic activity of environmental air samples from the area of Wroclaw, Poland. *Polish J. Environ. Stud.* 16 (2007) 745–752.
- [50] Y. Oda, S. Nakamura, I. Oki, T. Kato, H. Shinagawa, Evaluation of the new system (umu-test) for the detection of environmental mutagens and carcinogens, *Mutat. Res.* 147(5) (1985) 219–229.
- [51] ISO 13829:2000. Water Quality- Determination of the genotoxicity of water and waste water using the umu-test.
- [52] J.W. Tukey, Comparing individual means in the analysis of variance, *Biometrics* 5 (1949) 99–114.
- [53] A. Dunge, A.K. Chakraborti, S. Singh, Mechanistic explanation to the variable degradation behaviour of stavudine and zidovudine under hydrolytic, oxidative and photolytic conditions, *J. Pharm. Biomed. Anal.* 35(4) (2004) 965–70.
- [54] S. Goldstein, D. Aschengrau, Y. Diamant, J. Rabani, Photolysis of aqueous H₂O₂: quantum yield and applications for polychromatic UV actinometry in photoreactors, *Env. Sci. Tech.* 41 (2007) 7486–7490.
- [55] G.V. Buxton, C.L. Greenstock, W.P. Helman, A.B. Ross, Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals (OH/O) in aqueous solution, *J. Phys. Chem. Ref. Data* 17 (1988) 513–886.
- [56] B.H. Bielski, D.E. Cabelli, R.L. Aruda, A.B. Ross, Reactivity of HO₂/O₂ radicals in aqueous solution, *J. Phys. Chem. Ref. Data* 14 (1985) 1041–1077.
- [57] C. Prasse, J.. Wenk, J.T. Jasper, T.A. Ternes, D.L. Sedlak, Co-occurrence of photochemical and microbiological transformation processes in open-water unit process wetlands, *Environ. Sci. Technol.* 49 (2015) 14136–14145.
- [58] P. Onstein, M.I. Stefan, J.R. Bolton, Competition kinetics method for the determination of rate constants for the reaction of hydroxyl radicals with organic pollutants using the

1 UV/H₂O₂ advanced oxidation technology: the rate constants for the tert-butyl formate ester
2 and 2,4-dinitrophenol, J. Adv. Oxid. Technol. 4(2) (1999) 231–236.
3

4 [59] J. K. Stanley, E.J. Perkins, T. Habib, J. G. Sims, P. Chappell, B. L. Escalon, M.
5 Wilbanks, N. Garcia-Reyero, The good, the bad, and the toxic: approaching hormesis in
6 *Daphnia magna* exposed to an energetic compound, Environ. Sci. Technol., 47 (16) (2013)
7 9424–9433.
8
9

10 [60] S. Li, Y. Tan, Hormetic response of cholinesterase from *Daphnia magna* in chronic
11 exposure to triazophos and chlorpyrifos, J. Environ. Sci. (China) 23(5) (2011) 852–859.
12
13

14 [61] N. Cedergreen, J. C. Streibig, P. Kudsk, S. K. Mathiassen, S. O. Dukec, The Occurrence
15 of hormesis in Plants and Algae, Dose Response 5(2) (2007) 150–162.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



Removal kinetics of Zidovudine and Stavudine under UV_{254} and UV_{254}/H_2O_2 were studied

Zidovudine and Stavudine quantum yields at 254 nm were estimated

Kinetic constants of reaction between OH radicals and antiretrovirals were estimated

No acute/chronic effects of treated and untreated samples on *Vibrio f.* and *Daphnia m.*

Mutagenic/genotoxic activity of treated and untreated diluted samples on *Salmonella t.*



| Zidovudine  <p style="text-align: right;">$pK_a = 9.8$ [17]</p> | | | | |
|--|----------------------|--------------------|--------------|------|
| WWTP effluent (ng/L) | Surface water (ng/L) | Groundwater (ng/L) | Location | Ref |
| -- | 18.3 | -- | Kenya | [18] |
| 110 – 90 | 17410 – 50 | 30 – 20 | Kenya | [16] |
| 513 | 7684 | -- | Kenya | [5] |
| -- | 973 – 51.7 | -- | South Africa | [7] |
| 564 – 98.2 | 170 – 1.2 | -- | Germany | [9] |
| 180 – 57 | 30 – 22 | -- | Germany | [19] |
| 37 – 22 | -- | -- | Finland | [6] |
| 191 – 154 | -- | -- | France | [20] |
| Stavudine  <p style="text-align: right;">$pK_a = 10$ [21]</p> | | | | |
| WWTP effluent (ng/L) | Surface water (ng/L) | Location | Ref | |
| -- | 778 – 102 | Kenya | [18] | |
| -- | 4.3 – 1.3 | Germany | [9] | |

Table 1

| | $\varepsilon_i^{254nm} (M^{-1} \cdot cm^{-1})$ | pH range |
|-----|--|----------|
| ZDV | $1.19 \cdot 10^4$ | 4 - 8 |
| STV | $7.81 \cdot 10^3$ | 6 - 8 |

Table 2

| | $\phi_i^{254} \text{ (mol} \cdot \text{ein}^{-1}\text{)}$ | pH range |
|-----|---|----------|
| ZDV | $(2.357 \pm 0.0589) \cdot 10^{-2}$ | 4 - 8 |
| STV | $(8.34 \pm 0.334) \cdot 10^{-4}$ | 6 - 8 |

Table 3

| ZDV (optimization mode) | | | | |
|--------------------------------|--|-----------|-------------------------------|--------------------------------|
| Run | $C_o \cdot 10^5$ (M) | pH | lamp nominal power (W) | σ (%) |
| <i>1</i> | 3.76 | 6 | 8 | 1.27 |
| <i>2</i> | 3.67 | 4 | 8 | 0.55 |
| <i>3</i> | 1.86 | 6 | 8 | 1.22 |
| <i>4</i> | 1.83 | 4 | 8 | 1.57 |
| <i>5</i> | 1.86 | 6 | 4.5 | 1.98 |
| ZDV (simulation mode) | | | | |
| <i>6</i> | 3.65 | 8 | 8 | 0.48 |
| <i>7</i> | 3.76 | 6 | 4.5 | 1.14 |
| STV (optimization mode) | | | | |
| <i>8</i> | 4.49 | 6 | 8 | 1.65 |
| <i>9</i> | 4.46 | 8 | 8 | 1.39 |
| <i>10</i> | 2.27 | 6 | 8 | 3.87 |
| STV (simulation mode) | | | | |
| <i>11</i> | 4.49 | 6 | 4.5 | 2.41 |

Table 4

| ZDV (optimization mode) | | | | | |
|--------------------------------|--|---|-----------|-------------------------------|--------------------------------|
| Run | $C_0 \cdot 10^5$ (M) | $[H_2O_2]_0 \cdot 10^3$ (M) | pH | lamp nominal power (W) | σ (%) |
| 1bis | 4.89 | 0.7 | 6 | 4.5 | 1.60 |
| 2bis | 4.69 | 1.61 | 6 | 4.5 | 0.90 |
| 3bis | 4.78 | 2.02 | 4 | 4.5 | 0.33 |
| 4bis | 4.86 | 2.69 | 6 | 4.5 | 0.36 |
| 5bis | 4.87 | 3.49 | 6 | 4.5 | 0.63 |
| ZDV (simulation mode) | | | | | |
| 6bis | 4.74 | 1.79 | 8 | 4.5 | 0.83 |
| 7bis | 4.80 | 1.87 | 6 | 8 | 1.59 |
| STV (optimization mode) | | | | | |
| 8bis | 4.47 | 1.83 | 6 | 4.5 | 3.74 |
| 9bis | 4.48 | 2.99 | 6 | 4.5 | 2.58 |
| 10bis | 4.47 | 4.17 | 6 | 4.5 | 3.74 |
| 11bis | 4.43 | 4.16 | 8 | 4.5 | 2.57 |
| 12bis | 4.47 | 1.83 | 6 | 8 | 1.82 |
| STV (simulation mode) | | | | | |
| 13bis | 4.38 | 4.09 | 8 | 8 | 1.17 |

Table 5

| | $k_{OH,i} (M^{-1} \cdot s^{-1})$ | pH range |
|-----|----------------------------------|----------|
| ZDV | $(9.98 \pm 0.68) \cdot 10^8$ | 4 - 8 |
| STV | $(2.03 \pm 0.18) \cdot 10^9$ | 6 - 8 |

Table 6

| | Dilution factor | Mutagenic ratio | | | | Induction ratio | | | |
|--|------------------|-----------------------|-----|-----|-----|-----------------------|-----|-----|-----|
| | | Conversion degree (%) | | | | Conversion degree (%) | | | |
| | | 0 | 45 | 90 | 100 | 0 | 45 | 90 | 100 |
| UV ₂₅₄ | 1 | 8.1 | 2.4 | 9.1 | 9.3 | 1.2 | 2.1 | 2.9 | 1.0 |
| | 10 ⁻¹ | 4.3 | 1.5 | 5.5 | 5.9 | 0.2 | 1.5 | 1.7 | 0.8 |
| | 10 ⁻² | 2.7 | 1 | 3.3 | 3.7 | ND | 1.9 | 1.5 | 0.2 |
| | 10 ⁻³ | 2 | ND | 2.6 | 2.8 | ND | ND | ND | ND |
| UV ₂₅₄ /H ₂ O ₂ | 1 | 7.9 | 6.5 | 2.9 | 0.8 | 1.5 | 0.5 | 3.1 | 0.9 |
| | 10 ⁻¹ | 6 | 4.9 | 0.5 | 0.3 | 0.7 | ND | 1.6 | 0.6 |
| | 10 ⁻² | 2.1 | 0.7 | ND | ND | ND | ND | 1.6 | ND |
| | 10 ⁻³ | ND | ND | ND | ND | ND | ND | ND | ND |

Table 7

| | Dilution factor | Mutagenic ratio | | | | Induction ratio | | | |
|--|------------------|-----------------------|-----|-----|-----|-----------------------|------|------|-----|
| | | Conversion degree (%) | | | | Conversion degree (%) | | | |
| | | 0 | 45 | 90 | 100 | 0 | 45 | 90 | 100 |
| UV ₂₅₄ | 1 | 4.2 | 4.4 | 5.1 | 2.1 | 0.2 | 2.58 | 2.68 | ND |
| | 10 ⁻¹ | 2.2 | 2.7 | 2.5 | 1.6 | ND | 2.02 | 2.22 | ND |
| | 10 ⁻² | 2 | 2.1 | 2.1 | 1.5 | ND | 2.52 | 2.20 | ND |
| | 10 ⁻³ | ND | ND | ND | ND | ND | 2.87 | 2.26 | ND |
| UV ₂₅₄ /H ₂ O ₂ | 1 | 4.1 | 4.6 | ND | ND | 0.5 | 2.68 | 2.50 | ND |
| | 10 ⁻¹ | 1.2 | 1.0 | ND | ND | ND | 2.22 | 2.28 | ND |
| | 10 ⁻² | ND | ND | ND | ND | ND | 2.20 | 2.05 | ND |
| | 10 ⁻³ | ND | ND | ND | ND | ND | 2.20 | 2.22 | ND |

Table 8





Figure 1

OPTIMIZATION MODE



SIMULATION MODE





Figure 2

OPTIMIZATION MODE



SIMULATION MODE




Figure 3




Figure 4

Table 1 - Levels of zidovudine and stavudine in WWTPs effluents, surface water, and ground water.

Table 2 - Estimated molar absorption coefficient of ZDV and STV.

Table 3 - Estimated direct photolysis quantum yields of ZDV and STV at 254 nm.

Table 4 - Experimental runs and experimental conditions used for the estimation of quantum yields photolysis at 254 nm of ZDV and STV along with percentage standard deviation (% σ).

Table 5 - Experimental runs and experimental conditions for $k_{OH,i}$ estimation of ZDV and STV along with percentage standard deviation (% σ).

Table 6 - Estimated kinetic constant of hydroxyl radical attack to ZDV and STV with 97% confidence interval.

Table 7 - Mutagenic results from Ames and Umu tests for ZDV treated solutions. Starting concentration: $4.5 \text{ mg}\cdot\text{L}^{-1}$.

Table 8 - Mutagenic results from Ames and Umu tests for STV treated solutions. Starting concentration: $4.35 \text{ mg}\cdot\text{L}^{-1}$.

Fig. 1 - Absorbance spectra of ZDV (a) and STV (b). $C_{\text{init}} = 5 \cdot 10^{-5} \text{ M}$. (—) pH = 4.0; (- - -) pH = 6.5; (···) pH = 8.0.

Fig. 2 - Comparison between experimental (symbols) and calculated (continuous lines) data for the direct photolysis of ZDV (a, b, e) and STV (c, d, f) under UV_{254} irradiation. Experimental conditions are summarized in Table 4 (a-1; b-4; c-9; d-10; e-6; f-11).

Fig. 3 - Comparison between experimental (symbols) and calculated (continuous lines) data for the $\text{UV}_{254}/\text{H}_2\text{O}_2$ degradation of ZDV (a, b, e) and STV (c, d, f). Experimental conditions are summarized in Table 5 (a-3bis; b-5bis; c-9bis; d-11bis; e-7bis; f-13bis). ZDV and STV concentrations are multiplied by 160 for visual convenience.

Fig. 4: Inhibition algal growth (*R. subcapitata*) for solutions containing the selected antiretrovirals by UV_{254} and $\text{UV}_{254}/\text{H}_2\text{O}_2$. Starting concentration: $4.5 \text{ mg}\cdot\text{L}^{-1}$ for ZDV, $4.35 \text{ mg}\cdot\text{L}^{-1}$ for STV. Data with different letters (a–d) are significantly different ($p < 0.05$).