

Understanding variation in salamander ionomes: A nutrient balance approach

Clay Prater^{1†}, David E. Scott², Stacey L. Lance², Schyler O. Nunziata³, Ryan Sherman⁴, Nathan Tomczyk^{2,5}, Krista A. Capps^{2,5*}, & Punidan D. Jeyasingh^{4*}

¹ Department of Geography, Loughborough University, Loughborough, UK

² Savannah River Ecology Laboratory, University of Georgia, Aiken SC, USA

³ Department of Entomology, University of Maryland, College Park MD, USA

⁴ Department of Integrative Biology, Oklahoma State University, Stillwater OK, USA

⁵ Odum School of Ecology, University of Georgia, Athens GA, USA

†Contact information of corresponding author, mail to: Epinal Way, Loughborough LE11 3TU, UK or prater.clay@gmail.com

*Co-senior authors.

Keywords: Amphibians, *Ambystoma*, ecological stoichiometry, ionomics, and ontogeny.

Running Title: Salamander ionomic variation

Journal: *Freshwater Biology*

Type of manuscript: Original Research

1 **Summary**

2 1. Ecological stoichiometry uses information on a few key biological elements (C, N, and P) to
3 explain complex ecological patterns. Although factors driving variation in these elements are
4 well-established, expanding stoichiometric principles to explore dynamics of the many other
5 essential elements comprising biological tissues (i.e., the ionome) is needed to determine their
6 metabolic relationships and better understand biological control of elemental flows through
7 ecosystems.

8 2. In this paper, we report observations of ionic variation in two species of salamander
9 (*Ambystoma opacum* and *A. talpoideum*) across ontogenic stages using specimens from
10 biological collections of two wetlands sampled over a 30-year period. This unique data set
11 allowed us to explore the extent of ionic variation between species, among ontogenic stages,
12 between sites, and through time.

13 3. We found species- and to a lesser extent site-specific differences in C, N, and P along with 13
14 other elements forming salamander ionomes but saw no evidence of temporal changes.
15 Salamander ionic composition was most strongly related to ontogeny with relatively higher
16 concentrations of many elements in adult males (i.e., Ca, P, S, Mg, Zn, and Cu) compared to
17 metamorphic juveniles, which had greater amounts of C, Fe, and Mn.

18 4. In addition to patterns of individual elements, covariance among elements was used to
19 construct multi-elemental nutrient balances, which revealed differences in salamander elemental
20 composition between species and sites and changes in elemental proportions across ontogenic
21 development. These multi-elemental balances distinguished among species-site-ontogenic stage
22 groups better than using only C, N, and P.

23 5. Overall, this study highlights the responsiveness of consumer ionomes to life-history and
24 environmental variation while reflecting underlying relationships among elements tied to
25 biological function. As such, ionomic studies can provide important insights into factors shaping
26 consumer elemental composition and for predicting how these changes might affect higher-order
27 ecological processes.

28 **Introduction**

29 Elements are the fundamental building blocks of living cells and are involved in all metabolic
30 processes. Organisms must take up elements from the environment and, despite vast differences
31 in environmental supplies, maintain their body elemental composition within a relatively narrow
32 range (Persson *et al.*, 2010). Elemental content differs considerably among species due to
33 diversity in classical traits (e.g., life-history, morphological, and physiological traits) that are
34 constructed using different elements or the same elements in differing proportions (Jeyasingh,
35 Cothran & Tobler, 2014). Thus, it follows that the elemental composition of an individual is
36 determined by the acquisition, assimilation, and allocation of elements within the organism. This
37 abstraction is useful because information on the elemental content of a species enables the
38 application of mass-balance principles to make predictions about sequestration and excretion of
39 elements, both of which have strong impacts on ecological dynamics (Reiners, 1986). This
40 stoichiometric approach has triggered a vibrant area of research integrating data on species
41 elemental composition with cellular processes and organismal life-history to study key
42 ecological phenomena (Sturner & Elser, 2002).

43 In addition to species-level differences, it is apparent that there is considerable
44 intraspecific variation in organismal elemental composition (Jeyasingh *et al.*, 2014; Prater,
45 Wagner & Frost, 2017). For example, Bertram *et al.* (2008) found that variation in the
46 phosphorus (P) content of a single cricket species collected across field populations was as
47 extensive as that observed across several orders of insect taxa (Woods *et al.*, 2004). Further,
48 strong P plasticity among distinct ontogenic stages in metazoans linked to differences in
49 organismal growth and reproduction has commonly been observed (e.g., Villar-argaiz *et al.*
50 2002, Capps *et al.* 2015, Tiegs *et al.* 2016). Besides P, recent work also highlights the potential

51 ecological importance of variation in other elements within organismal tissues, although
52 metabolic causes of this variation are not always well-known (e.g., Baxter 2010; Jeyasingh et al.
53 2017). At a time when it is becoming increasingly clear that growth and productivity can be
54 limited by combinations of elements (Salt, Baxter & Lahner, 2008; Harpole *et al.*, 2011; Parent
55 *et al.*, 2013a), determination of the extent and nature of variation in the content of the other ~20
56 essential elements found in biological tissues is needed to establish their connections with
57 organismal metabolism and ecological functions.

58 The ionome is defined as the mineral nutrient and trace element composition of an
59 organism (Salt, Baxter & Lahner, 2008), which underlies its morphological, anatomical, and
60 physiological state. Although little is known about the ionomes of phagotrophic metazoans (but
61 see Goos et al. 2017), studies on osmotrophs such as *Saccharomyces cerevisiae* (Eide *et al.*,
62 2005) and *Arabidopsis thaliana* (Baxter *et al.*, 2008) in controlled laboratory conditions clearly
63 show that genetics and resource-supply stoichiometry interact to shape organismal growth and
64 ionic profiles. Evidence for ionome-wide shifts from the field are also available. For example,
65 *Synechococcus* cells collected from regions of the Sargasso Sea that vary in nitrogen (N) and P
66 supplies exhibited several-fold cell quota differences in a variety of elements [e.g., manganese
67 (Mn), nickel (Ni), and zinc (Zn); Twining et al. 2010]. These studies depict a strong correlative
68 network among elements likely underpinned by a complex web of metabolic connections (Parent
69 *et al.*, 2013a; Baxter, 2015). However, our knowledge of factors related to elemental variation in
70 vertebrate consumers, particularly in ecologically sensitive taxa such as amphibians, remains
71 limited to a relatively small number of elements.

72 Throughout ontogeny, frog and salamander carbon (C) content generally decreases while
73 %N, P, calcium (Ca), and sulfur (S) tend to increase (Capps *et al.*, 2015; Luhring, DeLong &

74 Semlitsch, 2017). These patterns are correlated with developmental and growth rates (Bumpers
75 et al. 2015; Stephens et al. 2017), arguably due to ontogenic changes in macromolecular
76 demands that differ in stoichiometric composition (e.g., lipids rich in C, proteins rich in N or S,
77 and bones rich in Ca and P; Costello and Michel 2013; Liess et al. 2013; Stephens et al. 2017). In
78 addition to ontogeny, amphibian C:N:P stoichiometry differs considerably across sites at local
79 scales (Milanovich & Hopton, 2014) due to differences in both bottom-up nutrient supplies
80 (Stephens, Berven & Tiegs, 2013; Bumpers *et al.*, 2015) and top-down predation pressure
81 (Costello & Michel, 2013). Furthermore, abiotic factors such as hydroperiod and temperature
82 have been shown to alter population-level stoichiometry at decadal scales (Capps et al. 2015) and
83 might explain regional stoichiometric differences through local adaptation of amphibian
84 populations across latitudinal gradients (Liess *et al.*, 2013). Overall, amphibian composition of a
85 handful of elements appears to be highly responsive to many ecological factors, and these
86 changes could result in shifts of other metabolically connected elements ultimately influencing
87 biogeochemical fluxes through aquatic environments.

88 Here, we report observations on ionic variation in two Ambystomatid salamander
89 species [marbled salamanders (*Ambystoma opacum*) and mole salamanders (*Ambystoma*
90 *talpoideum*)] using specimens that were collected at two wetlands over a 30-yr period. Based on
91 stoichiometric theory, we hypothesized that the ionomes of each species should vary due to
92 differences in life-history and ontogeny, and we sought to identify linkages among elements,
93 including those already known to vary significantly (i.e., C, Ca, N, P, and S; Capps et al. 2015,
94 Tiegs et al. 2016, Luhring et al. 2017). Moreover, because hydroperiod can affect salamander
95 growth and development, we hypothesized that salamander ionomes should vary due to
96 documented differences in hydroperiod between sites. Since hydroperiod differed considerably

97 over the 30-yr time period during which samples were collected (Daszak *et al.*, 2005; Todd *et al.*,
98 2011), we also expected to see temporal changes in salamander ionomes. In addition to
99 examining variation in individual elements, we also constructed nutrient balances (Parent *et al.*,
100 2013a,b) using knowledge of elemental metabolism and multivariate data from 15 elements to
101 see how these basic biological factors influence covariation among suites of elements.

102 **Methods**

103 *Study Sites, Specimen Collection, and Salamander Life-History:* Study animals were collected at
104 the United States Department of Energy's Savannah River Site (SRS) in South Carolina.
105 Salamanders were sampled at two ephemeral wetlands, Rainbow Bay (RB: 33.315696; -
106 81.773643) and Ginger's Bay (GB: 33.260035; -81.631320), as part of a long-term monitoring
107 program that began in 1978 and 1986 at each wetland, respectively (Scott, 1990; Pechmann *et*
108 *al.*, 1991). Rainbow Bay and GB are similar in area at full ponding (1-1.5 ha) but are located 14
109 km apart in different physiographic subregions. Rainbow Bay is found at ~97 m elevation on the
110 Aiken Plateau, whereas GB occurs on an old Pleistocene floodplain terrace of the Savannah
111 River (Sunderland Terrace) at 61 m elevation. Both wetlands are surrounded by well-drained
112 sandy loam soils in the Fuquay and Dothan series, but wetlands themselves are located on
113 poorly-drained sandy fine loam soils (Davis & Janecek, 1997). Wetland soils also differ in
114 elemental composition from surrounding upland soils as depressional wetland soils on the SRS
115 tend to have elevated aluminum (Al) and barium (Ba) concentrations and total N, P, and C levels
116 compared to upland soils (Looney *et al.*, 1990; Dixon *et al.*, 1997). Wetlands on the Sunderland
117 Terrace (GB) usually hold water for more extended periods (i.e., longer hydroperiod) than Aiken
118 Plateau wetlands (RB), have dissimilar zooplankton communities (Mahoney, Mort & Taylor,
119 1990), and may have differences in cation concentrations (Schalles *et al.*, 1989). The mean

120 hydroperiod at RB is 129.6 ± 98.5 days (mean \pm 1 SD), and it has never held water from one year
121 to the next; GB in contrast has a longer hydroperiod (D.E. Scott and K.A. Capps *unpublished*
122 *data*) and remains filled year-round on average once every three years.

123 Each wetland was encircled by a drift fence with paired 19-L pitfall traps positioned at
124 10-m intervals. Individuals entering and leaving the wetlands were censused daily (RB) or
125 periodically (GB) during the breeding season which occurs from late fall through early spring for
126 our study species (Semlitsch *et al.*, 1993; Scott *et al.*, 2013). When mortality occurred during the
127 study (usually due to predation or bucket flooding) deceased salamanders were collected,
128 identified individuals to species, and terrestrial adults were sexed. Partial remains of specimens
129 exposed to predators were used for census counts but excluded from our current study. We
130 classified specimens as one of three ontogenic stages: terrestrial adult male (*A. opacum* only),
131 recently emerged metamorph (both species), or recently emerged aquatic adult paedomorph (*A.*
132 *talpoideum* only; defined below) and stored them in a -70°C freezer until processing (Nunziata,
133 Scott & Lance, 2015). A total of 214 organisms were analyzed in our study representing a mix of
134 species, sites, and ontogenic stages collected between 1986-2015 (Supp. Table 1). All study
135 specimens were collected under annual renewals of South Carolina Department of Natural
136 Resources collection permits and triannual renewals of animal capture and handling protocols
137 approved by the University of Georgia Institutional Animal Care and Use Committee.

138 Ambystomatid salamander life-history differs considerably between species. *Ambystoma*
139 *opacum* breeds terrestrially in late summer/autumn (Sept. – Dec.) and oviposits in nests in dry
140 pond beds prior to wetland filling (Scott, 2005). *Ambystoma talpoideum* breeds aquatically
141 during late fall and winter (Nov. - Mar.), and females lay single eggs or small clusters attached
142 to aquatic vegetation (Pechmann *et al.* 1991, Semlitsch 1987). *Ambystoma talpoideum* eggs

143 typically hatch about two months later than *A. opacum*, which may lead to a predatory or
144 competitive advantage for *A. opacum* larvae (Boone, Scott & Niewiarowski, 2002). In wetlands
145 such as GB that occasionally hold water from one year to the next, *A. talpoideum* are
146 facultatively paedomorphic. Rather than metamorphosing and emigrating to terrestrial uplands,
147 these individuals may remain in the wetland, retain their gills, become sexually mature aquatic
148 adults in the winter (Semlitsch, 1987), and usually exit the wetland in spring after one
149 reproductive bout.

150 Ambystomatid salamander life-history varies with environmental conditions including
151 temperature, predation, food quantity/quality, and population density, but appears to be most
152 strongly tied to pond hydroperiod (Semlitsch 1987, Scott 1990, Pechmann et al. 1991, Daszak et
153 al. 2005, Stephens et al. 2017). Yet, despite this extensive phenotypic variation clear differences
154 exist between our study species. Both species appear to have relatively similar larval growth
155 rates (Semlitsch, 1987; Scott, 1990), although *A. talpoideum* typically metamorphoses later and
156 at a larger body size than *A. opacum* (Pechmann, 1995). Reproductive success (i.e., the
157 production of juveniles that metamorphose and emigrate from wetlands) of both species is tied to
158 hydroperiod, but *A. talpoideum* requires a later date of pond drying and more frequently faces
159 inadequate water at RB, which can result in catastrophic larval mortality (Daszak *et al.*, 2005).
160 Consequently, both species had stable populations at GB over our study period (Nunziata *et al.*,
161 2015), but *A. opacum* populations increased, and *A. talpoideum* decreased apparently due to a
162 shortened hydroperiod at RB (Daszak *et al.*, 2005). Few mortality events occurred for terrestrial
163 *A. talpoideum*; as a result, fewer of these species were collected compared to *A. opacum*. This
164 partially explains the uneven sample sizes and complete lack of adult *A. talpoideum* in our data
165 set (Supp. Table 1).

166 *Sample Processing and Ionomic Profile Generation:* We removed animals from the
167 freezer and placed them individually into trace clean polypropylene grinding vials. Carcasses
168 were then freeze-dried and powdered using a Spex Mill (8000D, Metuchen, NJ) and
169 methacrylate grinding balls, and ground tissues were stored at -20°C. Before elemental analyses,
170 we dried subsamples overnight at 60°C and weighed them to the nearest 0.1 µg. Tissue C and N
171 content was measured using an automated vario MICRO cube analyzer (Elementar Americas
172 Inc., Mt. Laurel, NJ). To measure all other elements, we digested between 3-12 mg of
173 homogenized tissues in a 2:1 v/v solution of trace metal grade nitric acid and hydrogen peroxide
174 (BDH Aristar ® Plus, VWR International, Radnor, PA) for 24h in metal-free polypropylene
175 tubes (VWR International, Radnor, PA) followed by dilution to 10 ml with trace metal grade
176 water. Digested samples were then analyzed through inductively coupled plasma optical
177 emission spectrometry (ICP-OES; Thermo Scientific iCAP 7400, Waltham, MA). Sample
178 nutrient concentrations were determined using standard curves from multi-element standards
179 (CCV stds. 1A&B, CPI International, Santa Rosa, CA) and calibrated using an internal Yttrium
180 standard (Peak Performance Inorganic Y Standard, CPI International, Santa Rosa, CA).

181 We measured a total of 28 elements across all organisms. However, measurements for
182 many trace elements fell below the limits of detection (LOD) for many samples (Supp. Table 2).
183 After omitting all elements with analytical uncertainties, a total of 16 elements: Al, Ba, C, Ca,
184 copper (Cu), iron (Fe), potassium (K), lithium (Li), magnesium (Mg), Mn, N, sodium (Na), P, S,
185 strontium (Sr), and Zn were analyzed. For multivariate statistics, any remaining values falling
186 below the LOD's were replaced with ½ of the LOD value calculated for each individual element.
187 All elemental concentrations were then converted to percentages by dividing them by the total
188 sample dry mass (Supp. Table 3).

190 To visualize relationships between individual elements in Euclidian space, we first
191 conducted partial least squares (PLS) regressions using ontogeny, species, site, and year as
192 independent variables and log-transformed elemental percentages as dependent variables. We
193 also calculated variable importance (VIP) scores to compare the relative importance of each
194 predictor variable (Wold, Sjöström & Eriksson, 2001; Eriksson *et al.*, 2013). Plots of PLS
195 weights revealed correlations between elements and predictor variables, which differentially
196 separated across two factors (Fig. 1). These correlations reflect biologically relevant differences
197 in elemental signatures among organisms. Therefore, we used these partitions (i.e., elemental
198 position in Euclidian space) to construct nutrient balances to further explore differences in
199 elemental profiles among ontogenic stages, species, sites, and years.

200 To more thoroughly investigate sources of variation in salamander ionomes, we
201 constructed a series of isometric log-ratio balances. These balances represent unbiased estimates
202 of multivariate relationships between elements, which avoid violating common statistical
203 assumptions and can be used to describe and interpret elemental interactions in organismal
204 tissues (Parent *et al.*, 2013a). Specifically defined in the context of our study and mathematically
205 defined below, nutrient balances represent orthogonal log contrasts of elements derived from
206 binary partitions of multivariate elemental data projected into Euclidean space. To construct
207 these balances (hereafter referred to as nutrient or elemental balances), we first separated all
208 elements into bulk and trace elements based on the classification of Frausto da Silva and
209 Williams (2001). Then we constructed one bulk and one trace elemental balance to reflect
210 elemental partitions along each PLS axis using absolute PLS weight values of 0.1 as a cutoff for
211 including elements in each balance. Following conventions (Sternner & Elser, 2002), we ordered

212 elements from high to low in each balance as a function of their percentage of dry mass. Nutrient
213 balances were then calculated using the equation:

$$214 \text{ Balance} = \text{SQRT} \left(\frac{rs}{r+s} \right) \ln \left[\frac{g(c^+)}{g(c^-)} \right] \quad (\text{Parent } et al., 2013a)$$

215 where r and s represent the number of elements on the left- and right-hand side of the balance
216 and $g(c^+)$ and $g(c^-)$ minus represent the geometric mean of elemental percentages on the left- and
217 right-hand side of the balance, respectively. To calculate a [C | Ca, P] balance for an adult
218 salamander with an elemental composition of 43% C, 5% Ca, and 3% P for example, we would
219 use the formula $[C | Ca, P] = \text{SQRT} \left[\frac{(1 \times 2)}{(1+2)} \right] \ln \left[\frac{43/g(5,3)}{1} \right] = 1.97$. Balances constructed in
220 this way can then be used to be used to examine proportional differences in elemental
221 composition between groups of organisms such as adults and juveniles that are related to
222 underlying biological factors. For instance, we would expect that [C | Ca, P] balance differences
223 between these two ontogenic stages should reflect proportionally higher investment in C-rich
224 lipids in juveniles and higher Ca and P investment into bone for adults resulting in relatively
225 higher juvenile values (Fig. 2).

226 In total, we constructed 4 novel balances: bulk [C | Ca, P, S, Mg]; [N | Ca, Na, Mg] and
227 trace [Fe, Mn | Zn, Ba, Cu, Sr, Li]; [Fe, Zn, Al, Ba | Cu, Sr, Li] in addition to two traditional
228 stoichiometric balances [C | P] and [N | P]. These balances represent a combination of 1)
229 knowledge of well-known elemental relationships related to organismal macromolecular
230 composition (C, N, P, Ca) and 2) statistically identified biologically relevant relationships related
231 to our study variables (ontogeny, species, site, and time). As such their formation represents a
232 blend of metabolically grounded expert knowledge and exploratory biplot analyses (Parent *et al.*,
233 2013b), which while not exhaustive may serve as a starting point for future functional studies.

234 To compare differences in individual elemental balances, we first separated the data into
235 unique species, site, and ontogenic stage groups (n=7) and conducted one-way analysis of
236 variance (ANOVAs) to examine differences across groups. After identifying significant group
237 differences, we designed contrasts to compare differences across species at the same ontogenic
238 stage (metamorphs averaged across sites), sites (separate comparisons of metamorphs and adult
239 males averaged across species), and stages (*A. opacum* males and metamorphs averaged across
240 sites and *A. talpoideum* metamorphs and paedomorphs from GB) using post-hoc Tukey's tests.
241 Specific comparisons were made rather than full factorial comparisons as complete species, site,
242 stage combinations were not available. All *P*-values were adjusted using Bonferroni corrections
243 to provide conservative estimates of elemental differences for comparison to multivariate
244 statistics.

245 In addition to univariate comparisons, we first used Wilks' Lambda tests from
246 multivariate ANOVA (MANOVA) to determine multivariate differences in nutrient balances
247 among each species, ontogenic stage, and site groups. We used similar methods to examine
248 temporal patterns in salamander balances by conducting MANOVAs on salamanders separated
249 into species(site) data sets for each ontogenic stage, which were grouped by decade to
250 compensate for uneven sample collection and inadequate samples sizes in certain years across
251 the 30-year period. Finally, we conducted two separate discriminant analyses including all
252 balances and one including only [C | P] and [N | P] balances to: a) visualize multivariate trait
253 differences across groups and b) examine the classification accuracy of classical stoichiometric
254 balances vs. all elemental balances. Cross-validation of these models was conducted using leave-
255 one-out methods. Overall, these complimentary univariate and multivariate analyses were
256 designed to quantify ecological sources of variance in individual balances (ANOVAs) and

257 multivariate phenotypes (MANOVAs & discriminant analyses) and reduce the likelihood of
258 statistical artifacts associated with multivariate data analyses.

259 **Results**

260 *Individual Elemental Profiles:*

261 Ontogeny was the strongest predictor of elemental variation according to PLS regressions
262 followed by species and site, which were moderate and weak predictors, respectively (Fig. 1A).
263 Temporal effects were the weakest by far ($VIP < 0.2$) indicating that salamander elemental
264 content was relatively invariant through time. Elemental composition was most divergent along
265 the first PLS factor, which separated strongly between terrestrial adult males (hereby referred to
266 as males) and juvenile metamorphic ontogenic stages with aquatic adult paedomorphs showing
267 intermediate phenotypes (Fig. 1B). Species and sites also separated out to a lesser extent along
268 this axis with *A. opacum* and individuals from GB falling on the left-hand side and *A. talpoideum*
269 and specimens from RB located on the right. Bulk elemental concentrations on the male side of
270 the axis were generally higher for all elements other than C, which was higher in metamorphs
271 and N, Na, and K, which did not differ greatly along this axis. Trace elements were were also
272 higher in males for all elements except for Fe and Mn. Compared to the first factor, factor two
273 explained far less variation but seemed to separate paedo- and metamorphic ontogenic stages
274 along with species and site differences. Relatively fewer bulk elements separated on this axis as
275 N was positively correlated and Ca, Mg, and Na were negatively correlated with factor two,
276 respectively. In contrast, all trace elements differed along the second axis with higher
277 concentrations of Fe and Zn associated with *A. opacum* and RB and *A. talpoideum* and
278 salamanders from GB showing higher amounts of Cu.

279 *Nutrient Balances:*

280 In addition to individual elements, nutrient balances differed across species, sites, and
281 ontogenic stages (Fig. 3). Metamorphs of *A. opacum* and *A. talpoideum* differed significantly in
282 their body [N | P] and [C | P] but did not differ in other balances (Table 1). In contrast,
283 salamander stoichiometric balances did not differ between sites, but we found significant site-
284 specific differences for all other multi-element balances in metamorphs and for [N | Ca, Na, Mg]
285 balances in males. All balances differed significantly among *A. opacum* males and metamorphs,
286 but only [N | P] differed between *A. talpoideum* metamorphs and paedomorphs.

287 Balances for unique species-site-ontogenic stage group combinations also differed in
288 multivariate space ($P < 0.001$), but we found no evidence of significant temporal changes ($P > 0.1$).
289 According to the discriminant analysis, male *A. opacum* phenotypes diverged strongly from other
290 groups along the first discriminant axis (Fig. 4), as males showed lower [C | P] and [N | Ca, Na,
291 Mg] values and other stages had higher [C | Ca, P, S, Mg] and [Fe, Mn | Zn, Ba, Cu, Sr, Li]
292 balances. Species differences separated out on the second discriminant axis as *A. opacum*
293 generally showed lower [C | P] and [Fe, Zn, Al, Ba | Cu, Sr, Li] concentrations and *A. talpoideum*
294 had higher [N | P] and [C | Ca, P, S, Mg]. Elemental balances showed relatively greater overlap
295 between sites in *A. talpoideum* but were more negatively correlated with the second discriminant
296 axis for each ontogenic stage in *A. opacum* from RB. Dual balances explained nearly half of the
297 elemental variation among all groups but were inadequate for distinguishing between
298 metamorphs of each species (Table 2). Including additional multi-elemental balances improved
299 classification accuracy from 49 to 65% among all groups. Despite our conservative analyses, it is
300 important to note that elemental correlations and nutrient balances in our data set could still
301 possibly contain information attributable to artifacts of high-dimensional analyses. Although we

302 cannot quantify the potential for these artifacts, consistency between uni- and multivariate
303 statistics suggests that their likelihood of influencing our findings was minimal.

304 **Discussion**

305 In this study, we found ecologically relevant differences in traditionally studied stoichiometric
306 elements (i.e., C, N, and P) along with a suite of other elements comprising salamander ionomes.
307 In contrast, we saw no significant temporal changes in the elemental composition of either
308 species. Salamander ionic composition was most strongly related to ontogeny with relatively
309 higher concentrations of many elements in males (Ca, P, S, Mg, Zn, and Cu) compared to
310 recently emerged metamorphic juveniles, which had greater amounts of C, Fe, and Mn. In
311 addition to identifying correlations among single elements, we also found differences among
312 species, site, and ontogenic stages in multivariate space. We observed systematic differences in
313 elemental relationships between species and sites and evidence of suites of changes in elemental
314 proportions across ontogenic stages. Together, these results demonstrate the utility of ionic
315 approaches for studying environmental and biological sources of consumer elemental variation
316 and their ecological effects.

317 Similar to previous amphibian studies (Milanovich, Maerz & Rosemond, 2015; Luhring
318 *et al.*, 2017), we found strong differences in the elemental composition of our study species.
319 Consistent with stoichiometric predictions, the species with the smaller body size and faster
320 developmental rates, *A. opacum*, had higher body P content than *A. talpoideum*. The larger-
321 bodied *A. talpoideum* also had higher body %C, which supports previous work showing positive
322 relationships between body size, juvenile lipid content, and body %C in ambystomatid
323 salamanders from this study region (Scott *et al.*, 2007; Luhring *et al.*, 2017). In addition to these
324 traditional stoichiometric elements, we also confirmed species differences in salamander Ca and

325 S content found in previous work (Milanovich *et al.*, 2015; Luhring *et al.*, 2017) and identified
326 differences in several other essential elements. Altogether, these patterns validate the focus on
327 well-studied elements (i.e., C and P) for determining the ecological importance of interspecific
328 variation in body stoichiometry. Our results also suggest that expanding the range of study
329 elements could yield further insights into the ecological roles and nutritional ecology of these
330 species.

331 Compared to species, we found relatively weaker evidence of spatio-temporal effects on
332 salamander ionomes. Site was a poor predictor of stoichiometric variation in males and
333 metamorphs despite strong differences in genetic structure for *A. opacum* between these
334 populations (Nunziata *et al.*, 2015). Male multi-elemental balances were also similar across sites,
335 but metamorphs showed strong contrasts between RB and GB. These patterns seem to indicate
336 that while male nutrient uptake (i.e., diet based) appeared to be similar in the surrounding
337 uplands, metamorph elemental acquisition either from dietary sources or epithelial diffusion
338 directly from the water column (e.g., metals or ions; Motais and Garcia-Romeau 1972, Handy et
339 al. 2002) likely differed across wetlands separated by only a short distance (~14 km). In addition
340 to differences between sites, there was also extensive annual variation in breeding season
341 temperature and hydroperiod within each site (Pechmann *et al.*, 1991; Daszak *et al.*, 2005; Todd
342 *et al.*, 2011). This variation may have limited our ability to identify temporal changes in
343 salamander elemental profiles and likely resulted in a lack of consistent selection pressures on
344 salamander life-history traits tied to their elemental composition. Indeed, previous genetic work
345 examining a subset of animals from our study (Nunziata *et al.*, 2017) along with our phenotypic
346 results seem to preclude the role of evolutionary change in shaping observed ionic patterns in
347 our study populations. However, as longer-term hydroperiod trends have been tied to shifts in

348 species abundances at RB (Daszak *et al.*, 2005), hydroperiod could nevertheless still modify
349 nutrient cycles in these wetlands by causing local species extinctions and thus shaping the
350 elemental profiles of the remaining amphibian communities. However, these ecological changes
351 would almost certainly interact with organismal life-history and ontogeny to ultimately control
352 these dynamics.

353 Salamander ontogenic stage had the strongest effect on elemental variation in our study
354 animals. Consistent with a previous amphibian study, we found that larger-bodied terrestrial
355 adult males had considerably higher body %P than juvenile metamorphs (Tiegs *et al.*, 2016).
356 Higher %P along with %Ca is typically associated with increased bone development and
357 ossification in adults (Kemp & Hoyt, 1969; Milanovich *et al.*, 2015; Luhring *et al.*, 2017) and
358 may explain the lower adult [C | P] and [N | Ca, Na, Mg] balances found in our study. In addition
359 to these elements, we measured higher concentrations of many other essential elements in adult
360 tissues (Zn, S, Li, Cu, and Mg). Our results also confirm previous measurements of higher
361 juvenile body %C and no difference in body N content between terrestrial adults and
362 metamorphic amphibians (Tiegs *et al.*, 2016) while showing differences in Fe and Mn
363 concentrations and in related [Fe, Mn | Zn, Ba, Cu, Sr, Li] nutrient balances between these
364 groups for the first time. This suggests that in addition to traditionally examined stoichiometric
365 elements (e.g., N and P; Stephens *et al.* 2017), ontogenic variation in organismal elemental
366 composition may reflect unique stage-specific nutritional physiologies and elemental
367 requirements. However, it also appears that environmental nutrient supplies may potentially
368 override these differences as we found substantial overlap in elemental profiles between pond-
369 dwelling *A. talpoideum* metamorphs and paedomorphic adults in GB, which only differed in
370 their %N content out of all elements surveyed (Supp. Table 3). As such, it is clear that we should

371 consider moving beyond studies of individual or pairs of elements and examine integrated
372 changes in suites of metabolically connected elements by refining tools such as nutrient balances
373 (Parent *et al.*, 2013a; Baxter, 2015) to better understand the complex relationships between
374 environmental nutrient supplies and organismal life-history traits and ionomes.

375 Traditional stoichiometric ratios relate nutrient interactions to biological functions (e.g.,
376 N:P ratios to ribosomal protein translation; Loladze and Elser 2011), and nutrient balance
377 concepts extend this approach to encompass the entire organismal ionome (Parent *et al.*, 2013a).
378 While it is beyond the scope of this paper to functionally relate changes in salamander balances
379 to specific metabolic pathways and physiological process, our study demonstrates the potential
380 of using nutrient balance principles to further our understanding of relationships between
381 consumer ecology and elemental composition. We found evidence for novel nutrient balances [C
382 | Ca, P, S, Mg] and [Fe, Mn | Zn, Ba, Cu, Sr, Li] that differed across ontogenic stages, and
383 similar to plant-based studies, we documented systematic differences in elemental combinations
384 between species and sites (Watanabe *et al.*, 2007; Parent *et al.*, 2013a). Previously constructed
385 balances inherently differ from ours due to their focus on specific aspects of plant physiology,
386 but the substantial overlap between elemental relationships in our study and previous research is
387 perhaps indicative of fundamental elemental relationships governed by biological processes
388 operating at the cellular level (Sturner & Elser, 2002; Watanabe *et al.*, 2007; Baxter *et al.*, 2008;
389 Parent *et al.*, 2013a).

390 In this study, we examined correlations between salamander ionomic profiles and
391 biological and environmental variables. As previously demonstrated, traditional stoichiometric
392 balances (C | P and N | P) were useful for highlighting species differences in biogeochemically
393 important elements (Sturner & Elser, 2002). However, we also showed that focusing solely on

394 traditional stoichiometric elements alone could neglect other important elements, which might be
395 differentially required for optimal metabolic functioning in terrestrial adult males (Ca, S, Mg,
396 Zn, & Cu) and juveniles (Fe & Mn), respectively. Ionic data analyzed here using balance
397 techniques could thus be used to generate a greater understanding of the nutritional ecology of
398 amphibian development. Such data are vital for providing a starting point for designing
399 manipulative experiments to not only better understand the nutritional ecology of threatened
400 vertebrates such as salamanders but also to illuminate the entire suite of ecological functions that
401 they might influence.

402 **Acknowledgements**

403 Many people have assisted with drift fence checks and specimen collection over the 39-yr
404 Rainbow Bay study, and we especially thank A. Chazal, A. Dancewicz-Helmers, R. Estes, J.
405 Greene-McLeod, R. Semlitsch, J. Caldwell, L. Vitt, and W. Gibbons. We would also like to
406 thank P. Lind and J. Goos for assistance with elemental analyses. This research was partially
407 supported by U. S. Department of Energy under Award Number DE-FC09-07SR22506 to the
408 University of Georgia Research Foundation and was also made possible the long-term protection
409 of Rainbow Bay and Ginger's Bay by the DOE's Set Aside Program, as well as the designation
410 of the SRS as a DOE National Environmental Research Park (NERP). Our study was also
411 partially supported by the DOE and National Nuclear Security Administration under award
412 numbers: DE-EM0004391 to the University of Georgia Research Foundation.

413 **Conflict of Interest Statement**

414 The authors declare no conflicts of interest.

415 **References**

- 416 Baxter I. (2010) Ionomics: The functional genomics of elements. **9**, 149–156.
- 417 Baxter I. (2015) Should we treat the ionome as a combination of individual elements, or should
418 we be deriving novel combined traits? *Journal of Experimental Botany* **66**, 2127–2131.
- 419 Baxter I.R., Vitek O., Lahner B., Muthukumar B., Morrissey J., Guerinot M. Lou, *et al.* (2008)
420 The leaf ionome as a multivariable system to detect a plant's physiological status. **105**,
421 12081–12086.
- 422 Bertram S.M., Bowen M., Kyle M. & Schade J.D. (2008) Extensive natural intraspecific
423 variation in stoichiometric (C:N:P) composition in two terrestrial insect species. *Journal of*
424 *insect science* **8**, 1–7.
- 425 Boone M.D., Scott D.E. & Niewiarowski P.H. (2002) Effects of hatching time for larval
426 ambystomatid salamanders. *Copeia* **2**, 511–517.
- 427 Bumpers P.M., Maerz J.C., Rosemond A.D. & Benstead J.P. (2015) Salamander growth rates
428 increase along an experimental stream phosphorous gradient. *Ecology* **96**, 2994–3004.
- 429 Capps K.A., Berven K.A. & Tiegs S.D. (2015) Modelling nutrient transport and transformation
430 by pool-breeding amphibians in forested landscapes using a 21-year dataset. *Freshwater*
431 *Biology* **60**, 500–511.
- 432 Costello D.M. & Michel M.J. (2013) Predator-induced defenses in tadpoles confound body
433 stoichiometry predictions of the general stress paradigm. *Ecology* **94**, 2229–36.
- 434 Daszak P., Scott D.E., Kilpatrick A.M., Faggioni C., Gibbons J.W. & Porter D. (2005)
435 Amphibian population declines at Savannah River Site are linked to climate, not

436 chytridiomycosis. *Ecology* **86**, 3232–3237.

437 Davis C.E. & Janecek L.L. (1997) *DOE Research Set-Aside Areas of the Savannah River Site*.

438 Dixon K.L., Rogers V.A., Conner S.P., Cummings C.L., Gladden J.B. & Weber J.M. (1997)

439 *Geochemical and Physical Properties of Wetland Soils at the Savannah River Site*. Aiken,

440 S.C.

441 Eide D.J., Clark S., Nair T.M., Gehl M., Gribskov M., Guerinot M. Lou, *et al.* (2005)

442 Characterization of the yeast ionome: a genome-wide analysis of nutrient mineral and trace

443 element homeostasis in *Saccharomyces cerevisiae*. *Genome biology* **6**, R77.

444 Eriksson L., Byrne T., Johansson E., Trygg J. & Wilkstrom C. (2013) *Multi- and Megavariate*

445 *Data Analysis Basic Principles and Applications*, 3rd edn. Umetrics Inc., Umea.

446 Frausto da Silva J.J.R. & Williams R.J.P. (2001) *The Biological Chemistry of the Elements*, 2nd

447 edn. Oxford Univ Press, Oxford.

448 Goos J.M., Cothran R.D. & Jeyasingh P.D. (2017) Within-population variation in the chemistry

449 of life: the stoichiometry of sexual dimorphism in multiple dimensions. *Evolutionary*

450 *Ecology* **31**, 635–651.

451 Handy R.D., Eddy F.B. & Baines H. (2002) Sodium-dependent copper uptake across epithelia: A

452 review of rationale with experimental evidence from gill and intestine. *Biochimica et*

453 *Biophysica Acta - Biomembranes* **1566**, 104–115.

454 Harpole W.S., Ngai J.T., Cleland E.E., Seabloom E.W., Borer E.T., Bracken M.E.S., *et al.*

455 (2011) Nutrient co-limitation of primary producer communities. *Ecology letters* **14**, 852–62.

456 Jeyasingh P.D., Cothran R.D. & Tobler M. (2014) Testing the ecological consequences of

457 evolutionary change using elements. *Ecology and Evolution* **4**, 528–538.

458 Jeyasingh P.D., Goos J.M., Thompson S.K., Godwin C.M. & Cotner J.B. (2017) Ecological
459 stoichiometry beyond redfield: An ionic perspective on elemental homeostasis. *Frontiers*
460 *in Microbiology* **8**, 722.

461 Kemp N.E. & Hoyt J.A. (1969) Sequence of ossification in the skeleton of growing and
462 metamorphosing tadpoles of *Rana pipiens*. *Journal of Morphology* **129**, 415–443.

463 Liess A., Rowe O., Guo J., Thomsson G. & Lind M.I. (2013) Hot tadpoles from cold
464 environments need more nutrients - life history and stoichiometry reflects latitudinal
465 adaptation. *Journal of Animal Ecology* **82**, 1316–1325.

466 Loladze I. & Elser J.J. (2011) The origins of the Redfield nitrogen-to-phosphorus ratio are in a
467 homeostatic protein-to-rRNA ratio. *Ecology Letters* **14**, 244–250.

468 Looney B.B., Eddy C.A., Ramdeen M., Pickett J., Rogers V., Scott M.T., *et al.* (1990)
469 *Geochemical and Physical Properties of Soils and Shallow Sediments at the Savannah*
470 *River Site*.

471 Luhring T.M., DeLong J.P. & Semlitsch R.D. (2017) Stoichiometry and life-history interact to
472 determine the magnitude of cross-ecosystem element and biomass fluxes. *Frontiers in*
473 *Microbiology* **8**, 1–11.

474 Mahoney D.L., Mort M.A. & Taylor B.E. (1990) Species richness of calanoid copepods,
475 cladocerans and other branchiopods in Carolina bay temporary ponds. *The American*
476 *Midland Naturalist* **123**, 244–258.

477 Milanovich J.R. & Hopton M.E. (2014) Stoichiometry of a semi-aquatic plethodontid

478 salamander: Intraspecific variation due to location, size and diet. *Integrative Zoology* **9**,
479 613–622.

480 Milanovich J.R., Maerz J.C. & Rosemond A.D. (2015) Stoichiometry and estimates of nutrient
481 standing stocks of larval salamanders in Appalachian headwater streams. *Freshwater*
482 *Biology* **60**, 1340–1353.

483 Motais R. & Garcia-Romeau F. (1972) Transport mechanisms in the teleostean gill and amphibian
484 skin. *Am Rev Physiol* **34**, 141–176.

485 Nunziata S.O., Lance S.L., Scott D.E., Lemmon E.M. & Weisrock D.W. (2017) Genomic data
486 detect corresponding signatures of population size change on an ecological time scale in
487 two salamander species. *Molecular Ecology* **26**, 1060–1074.

488 Nunziata S.O., Scott D.E. & Lance S.L. (2015) Temporal genetic and demographic monitoring
489 of pond-breeding amphibians in three contrasting population systems. *Conservation*
490 *Genetics* **16**, 1335–1344.

491 Parent S., Parent L.E., Egozcue J.J., Rozane D., Hernandez A., Lapointe L., *et al.* (2013a) The
492 plant ionome revisited by the nutrient balance concept. *Frontiers in Plant Science* **4**, 1–10.

493 Parent S., Parent L.E., Rozane D.E. & Natale W. (2013b) Plant ionome diagnosis using sound
494 balances : case study with mango (*Mangifera Indica*). *Frontiers in Plant Science* **4**, 1–12.

495 Pechmann J.H.K. (1995) Use of large field enclosures to study the terrestrial ecology of pond
496 breeding amphibians. *Herpetologica* **51**, 434–450.

497 Pechmann J.H.K., Scott D.E., Semlitsch R.D., Caldwell J.P., Vitt L.J. & Gibbons J.W. (1991)
498 Declining amphibian populations: The problem of separating human impacts from natural

499 fluctuations. *Science* **253**, 892–895.

500 Persson J., Fink P., Goto A., Hood J.M., Jonas J. & Kato S. (2010) To be or not to be what you
501 eat: Regulation of stoichiometric homeostasis among autotrophs and heterotrophs. *Oikos*
502 **119**, 741–751.

503 Prater C., Wagner N.D. & Frost P.C. (2017) Interactive effects of genotype and food quality on
504 consumer growth rate and elemental content. *Ecology* **98**, 1399–1408.

505 Reiners W.A. (1986) Complementary models for ecosystems. *The American Naturalist* **127**, 59–
506 73.

507 Salt D.E., Baxter I. & Lahner B. (2008) Ionomics and the study of the plant ionome. *Annual*
508 *review of plant biology* **59**, 709–733.

509 Schalles J.F., Sharitz R.R., Gibbons J.W., Laversee G.J. & Knox J.N. (1989) *Carolina Bays of*
510 *the Savannah River Plant*.

511 Scott D.E. (2005) *Ambystoma opacum* (Marbled Salamander). In: *Amphibian Declines: The*
512 *Conservation Status of United States Species*. (Ed. M.J. Lannoo), pp. 627–632. University
513 of California Press, Berkeley.

514 Scott D.E. (1990) Effects of larval density in *Ambystoma opacum*: An experiment large-scale
515 field enclosures. *Ecology* **71**, 296–306.

516 Scott D.E., Casey E.D., Donovan M.F. & Lynch T.K. (2007) Amphibian lipid levels at
517 metamorphosis correlate to post-metamorphic terrestrial survival. *Oecologia* **153**, 521–532.

518 Scott D.E., Komoroski M.J., Croshaw D.A. & Dixon P.M. (2013) Terrestrial distribution of
519 pond-breeding salamanders around an isolated wetland. *Ecology* **94**, 2537–2546.

520 Semlitsch R.D. (1987) Density-dependent growth and fecundity in the paedomorphic salamander
521 *Ambystoma talpoideum*. *Ecology* **68**, 1003–1008.

522 Semlitsch R.D., Scott D.E., Pechmann J.H.K. & Gibbons J.W. (1993) Phenotypic variation in the
523 arrival time of breeding salamanders: individual repeatability and environmental influences.
524 *Journal of Animal Ecology* **62**, 334–340.

525 Stephens J.P., Berven K.A. & Tiegs S.D. (2013) Anthropogenic changes to leaf litter input affect
526 the fitness of a larval amphibian. *Freshwater Biology* **58**, 1631–1646.

527 Stephens J.P., Stoler A.B., Sckrabulis J.P., Fetzer A.J., Berven K.A., Tiegs S.D., *et al.* (2017)
528 Ontogenetic changes in sensitivity to nutrient limitation of tadpole growth. *Oecologia* **183**,
529 263–273.

530 Sterner R.W. & Elser J.J. (2002) *Ecological Stoichiometry*. Princeton University Press,
531 Princeton.

532 Tiegs S.D., Berven K.A., Carmack D.J. & Capps K.A. (2016) Stoichiometric implications of a
533 bi-phasic life cycle. *Oecologia* **179**, 853–863.

534 Todd B.D., Scott D.E., Pechmann J.H.K. & Gibbons J.W. (2011) Climate change correlates with
535 rapid delays and advancements in reproductive timing in an amphibian community.
536 *Proceedings of the Royal Society B: Biological Sciences* **278**, 2191–2197.

537 Twining B.S., Nun D., Vogt S., Johnson R.S. & Sedwick P.N. (2010) Variations in
538 *Synechococcus* cell quotas of phosphorus, sulfur, manganese, iron, nickel, and zinc within
539 mesoscale eddies in the Sargasso Sea. *Limnol. Oceanogr* **55**, 492–506.

540 Villar-argaiz M., Medina-sánchez J.M. & Carrillo P. (2002) Linking life history strategies and

541 ontogeny in crustacean zooplankton: Implications for homeostasis. *Ecology* **83**, 1899–1914.

542 Watanabe T., Broadley M.R., Jansen S., White P.J., Takada J., Satake K., *et al.* (2007)

543 Evolutionary control of leaf element composition in plants. *The New phytologist* **174**, 516–

544 523.

545 Wold S., Sjöström M. & Eriksson L. (2001) PLS-regression: A basic tool of chemometrics.

546 *Chemometrics and Intelligent Laboratory Systems* **58**, 109–130.

547 Woods H.A., Fagan W.F., Elser J.J. & Harrison J.F. (2004) Allometric and phylogenetic

548 variation in insect phosphorus content. *Functional Ecology* **18**, 103–109.

549

Table 1. Trait differences between species, sites, and ontogenic stages. Degrees of freedom (*df*), f-ratio of mean squares (F), and *P*-values (*P*) are reported for 1-way main effects ANOVAs and multiple comparisons. Significant differences are shown in bold for *P*-values adjusted using Bonferroni corrections (Meta: 0.05/3= 0.017; Adult: 0.05/2=0.025). Specific contrast details include: averaged across sites¹, averaged across species², *Ambystoma opacum* only³, average across sites *A. opacum* only⁴, and *A. talpoideum* from Ginger’s Bay only⁵. All elements are reported in standard scientific notation. Other abbreviations include: metamorph (Meta) and paedomorph (Paedo).

Model	<i>df</i>	[C P]		[N P]		[C Ca,P,S,Mg]		[N Ca,Na,Mg]		[Fe,Mn Zn,Ba,Cu,Sr,Li]		[Fe,Zn,Al,Ba Cu,Sr,Li]	
		<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
Group	6	44.88	<0.001	34.86	<0.001	53.56	<0.001	19.52	<0.001	64.52	<0.001	27.07	<0.001
R ²		0.565		0.503		0.608		0.361		0.652		0.440	
Contrast	<i>df</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
¹ Species (Meta)	1	7.12	0.008	8.26	0.005	0.85	0.358	0.02	0.877	1.53	0.217	0.00	0.998
² Site (Meta)	1	5.02	0.026	3.36	0.068	6.02	0.015	9.93	0.002	10.40	0.002	20.66	<0.001
³ Site (Male)	1	0.54	0.463	1.21	0.272	2.11	0.148	14.54	<0.001	0.95	0.332	0.49	0.484
⁴ Stage (Male vs. Meta)	1	165.07	<0.001	122.92	<0.001	221.84	<0.001	60.10	<0.001	259.22	<0.001	89.82	<0.001
⁵ Stage (Meta vs. Paedo)	1	0.04	0.838	9.33	0.003	0.01	0.906	3.26	0.072	0.26	0.611	4.60	0.033

Table 2. Comparison of discriminant analysis classification accuracy based on A) stoichiometric balances only and B) all elemental balances. Cross-validation results are reported as percentages, and percentages correctly assigned to each group are shown in bold. Abbreviations include: *Ambystoma opacum* (Amop), *A. talpoideum* (Amta), metamorph (Meta), paedomorph (Paedo), Ginger’s Bay (GB), and Rainbow Bay (RB).

A. [C|P] & [N|P]

Species	Ontogenic Stage	Site	Group	1	2	3	4	5	6	7	Total
Amop	Male	GB	1	61.5	12.8	0	25.6	0	0	0	49.1
Amop	Male	RB	2	43.2	35.1	0	18.9	0	0	2.7	
Amop	Meta	GB	3	0	0	0	100	0	0	0	
Amop	Meta	RB	4	0	0	0	98.4	0	1.6	0	
Amta	Meta	GB	5	0	5.6	0	83.3	0	11.1	0	
Amta	Meta	RB	6	0	0	0	71.4	0	9.5	19	
Amta	Paedo	GB	7	0	0	0	50	0	0	50	

B. All Balances

Species	Ontogenic Stage	Site	Group	1	2	3	4	5	6	7	Total
Amop	Male	GB	1	76.9	17.9	0	0	2.6	2.6	0	64.5
Amop	Male	RB	2	35.1	62.2	0	2.7	0	0	0	
Amop	Meta	GB	3	0	0	33.3	48.1	11.1	7.4	0	
Amop	Meta	RB	4	0	3.2	8.1	87.1	1.6	0	0	
Amta	Meta	GB	5	0	0	16.7	16.7	38.9	22.2	5.6	
Amta	Meta	RB	6	0	0	4.8	19	23.8	33.3	19	
Amta	Paedo	GB	7	0	0	10	0	10	0	80	

Figure Captions

Figure 1. Relationships between ontogeny, species, site, date, and salamander elemental composition. A) Variable importance predictor scores show the relative influence of predictor variables on organismal ionic profiles. Scores <0.8 are considered weakly important, those between $0.8-1$ are moderately important, and scores >1 are strong predictors. B) Partial least squares regression (PLS) loadings plot of variables weights demonstrates correlations between independent (x) and dependent (y) variables where relationships between variables are directly proportional to the sign and distance of all other variables in Euclidian space. Abbreviations of x-variables include: metamorph (meta), paedomorph (Paedo), *Ambystoma opacum* (Amop), *A. talpoideum* (Amta), Ginger's Bay (GB), and Rainbow Bay (RB).

Figure 2. Hypothetical differences in salamander elemental composition between adult and juvenile salamanders. Panel A) depicts univariate differences in body elemental composition between an adult and a juvenile salamander. Panel B) represents a mobile-and-fulcrum plot (*sensu* Parent et al. 2013) showing multivariate differences in [C | Ca, P] balances between these two ontogenic stages. The black circle represents an equilibrium point where salamander C and [Ca, P] content is equal for a given specimen, and the other circles correspond to balance points for each stage.

Figure 3. Variation in nutrient balances among species, sites, and ontogenic stages. Boxplots depict medians, 25th and 75th percentiles (boxes), and 10th and 90th percentiles (error bars) for each balance. All balances differed significantly among species (left- and right-hand panels), sites, and ontogenic stages according to Wilks's lambda scores ($P < 0.001$). Abbreviations include: metamorph (Meta), paedomorph (Paedo), Ginger's Bay (GB; Panels A&B), and Rainbow Bay (RB; Panels C&D).

Figure 4. Multivariate relationships among species, sites, and ontogenic stages. Discriminant analysis (DA) loadings plots of centroids and 95% confidence intervals are shown for each group, and the two highest positive and negative standardized canonical discriminant function coefficient scores out of six nutrient balances are reported for each axis. All balances differed significantly among groups according to Wilks's lambda scores ($P < 0.001$). Abbreviations include: metamorph (Meta), paedomorph (Paedo), *Ambystoma opacum* (Amop), *A. talpoideum* (Amta), Ginger's Bay (GB), and Rainbow Bay (RB).

Figure 1




Figure 2




Figure 3




Figure 4

