This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Compulsive exercise and eating disorders

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1002/erv.1122>

PUBLISHER

© John Wiley & Sons, Ltd and Eating Disorders Association

VERSION

SMUR (Submitted Manuscript Under Review)

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Meyer, Caroline, Lorin Taranis, Huw Goodwin, and Emma Haycraft. 2019. "Compulsive Exercise and Eating Disorders". figshare. <https://hdl.handle.net/2134/11292>.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Compulsive exercise and eating disorders

Caroline Meyer, Lorin Taranis, Huw Goodwin and Emma Haycraft

Loughborough University Centre for Research into Eating Disorders

From the very earliest descriptions, clinicians have observed a tenacious drive for physical activity as a prominent characteristic of many eating disordered patients, even in the face of severe emaciation (Bruch, 1965; Crisp, 1967; Dally, 1969; Eisler & LeGrange, 1990; Gull, 1874; Inches, 1895; Kron, Katz, Gorzynski, & Viner, 1978; Lasegue, 1873; Long, Smith, Midgley, & Cassidy, 1993; Penas-Lledo, Vaz Leal, & Waller, 2002; Shroff, Reba, Thornton, Tozzi, Klump et al., 2006; Touyz, Beumont & Hoek, 1987). Over the last few decades there has been considerable research into compulsive exercise among both eating disordered patients and non-clinical groups (Davis, 2000). Some researchers have argued that problematic exercise occupies a central role in the pathogenesis of the eating disorders (e.g., Davis, Kennedy, Ralevski, & Dionne, 1994; Epling & Pierce, 1992; Kron et al., 1978). Indeed, it is now recognised as a significant factor in the aetiology, development and maintenance of eating disorders across diagnoses (Brewerton, Stelfox, Hibbs, Hodges & Cochrane, 1995; Davis, Katzman, Kaptein, Kirsh, Brewer et al., 1997; Hechler, Beumont, Marks, & Touyz, 2005), although particularly in anorexia nervosa (Beumont, Arthur, Russell, & Touyz, 1994; Davis, 1997; Davis et al., 1997; Katz, 1996; Touyz et al., 1987; Thien, Thomas, Markin, & Birmingham, 2000).

The importance of compulsive exercise in the eating disorders is highlighted by the fact that it is often one of the last symptoms to subside (Crisp, Hsu, Harding & Hartshorn, 1980; Davis et al., 1994; Kron et al., 1978). For example, a 3-year in-patient hospital records analysis found that eating disorder patients across diagnoses who were compulsive exercisers required a longer length of hospitalisation than non-compulsive exercisers (Solenberger, 2001). In addition, compulsive exercise has been identified as a significant risk factor for relapse in AN (Carter, Blackmore, Sutandar-Pinnock & Woodside, 2004; Casper & Jabine, 1996). A 10-year follow up study of 95 patients with anorexia nervosa found that

increased exercise not only significantly predicted a shorter time to relapse, but was also one of only two variables (out of more than 50) that was associated with a chronic outcome (Strober, Freeman & Morrell, 1997).

Despite the documented importance of exercise within the context of the eating disorders, there is currently no consensus about the cognitive attributes that define compulsive exercise. This is problematic, since any interventions tailored at reducing exercise behaviour or altering exercise cognitions requires a clear underpinning theoretical model. Therefore, the broad aim of this review paper is to develop an empirically supported, testable, cognitive behavioural model of compulsive exercise within the context of the eating disorders. In order to achieve this aim, two extensive literature reviews were undertaken. The first, a systematic review aimed at identifying the correlates and predictors of compulsive exercise among eating disordered patients, identified four key correlates of compulsive exercise: eating psychopathology; obsessive-compulsiveness; affect regulation; and perfectionism. Second, a broader review incorporating both clinical and non-clinical samples, as well as literature from the exercise science field, identified similar key indices thereby validating the proposed model. The findings from these searches were then amalgamated. First, an outline of the review methodologies used will be provided. This will be followed by presentation of the proposed model and a subsequent review of the evidence for the inclusion of the key constructs.

Review methods:

Inclusion / exclusion criteria:

For both reviews, empirical work, published in English, which investigated factors associated with compulsive exercise behaviour in eating disordered individuals, was included. Given the variety of terms used to describe compulsive exercise, wider terminology was included (see viewpoint in this edition). For the initial systematic review, any study that recruited all, or part of their sample from a clinical eating disordered population and which investigated exercise (and/or physical activity) behaviour or any qualitative dimension of

exercise (e.g. commitment to exercise, obligatory exercise, exercise addiction, exercise dependence) was included. Eating disorders were restricted to anorexia nervosa, bulimia nervosa, and eating disorder not otherwise specified (EDNOS). Consequently, studies on obesity and binge eating disorder were excluded, as were studies that simply investigated body image or body dissatisfaction, without any further assessment of eating pathology. Additionally, studies that investigated eating disorders among athletic or exercising populations were also excluded from this review, unless the study's specific focus was on compulsive exercise (and/or one of the above terms). Qualitative studies were included, but individual case studies were excluded. Finally, outcome studies were also excluded as these were beyond the focus of this review. With the second review, the sample selection was broadened out to include non-clinical community samples as well as studies that specifically recruited exercisers. In all cases, the studies included were those that had researched links between exercise and eating disorder-related attitudes and behaviours.

Importantly, several identified studies investigated predictors of extreme weight control behaviours and/or eating disordered behaviours, of which excessive exercising was sometimes mentioned. However, these studies were often simply an investigation of exercising for weight management reasons and are not measuring compulsive exercise. Therefore, these studies were also excluded.

Search method:

Evidence for both reviews was gathered via a comprehensive three-stage literature search. First, key electronic databases were used. These were Web of Science, PubMed, PsycInfo, Science Direct, EBSCO, OVID and SCOPUS. Search terms included, but were not limited to: *exercise, activity, excessive, compulsive, obsessive, obligatory, addiction, dependence, habitual, commitment, abuse, disordered, eating, anorexia, bulimia*. Wildcards were used wherever required to enable all possible permutations of a root word to be found. For example **exercis** would find the word "exercise" together with all possible prefixes and suffixes, such as *over-exercise, over-exercising, over-exerciser, exercising, exerciser,*

exercised, and so on. In addition, in order to narrow the search to relevant findings only, Boolean Operators were used to specify relationships between specific search terms/word. For example, **exercis** AND *eat** OR *excessive*. Second, eating disorder journals were subjected to individual searches using the same search terms (including European Eating Disorders Review, International Journal of Eating Disorders, Eating Behaviors, and Eating & Weight Disorders). Finally, reference lists of relevant review papers as well as reference lists of identified articles were systematically searched to ensure that any further articles missed by the database and journal searches were also identified and included for the review.

Results of the systematic review:

The following section outlines the results specifically of the initial systematic review. Due to the large number of search terms employed, over 500 possible articles were produced from the databases. However, after accounting for duplications, and using the exclusion and inclusion criteria stipulated, 39 articles were retained for closer inspection. Following on from the procedures outlined in Berkman, Lohr & Bulik (2007) and Atlantis & Baker (2008), these 39 articles were graded according to relevance, from '0' representing irrelevance, through to '4', which represented high relevance. A total of 22 articles were graded as either 3 or 4 and were subsequently retained for this review. The remaining articles, which had rankings of below 3, were excluded.

Description of the studies

The characteristics of the different samples were largely similar across the studies. Participants were predominantly recruited at eating disorder clinics and the majority of the studies used young adult female samples. This is reflective of the demographics typical of eating disorder clinics. The only exception to the use of a female-only sample was that of Blaydon, Linder, and Kerr (2004), who also recruited highly active male exercisers.

The vast majority of the studies adopted cross-sectional designs ($n = 15$). One paper used a prospective design (Davis & Kaptein, 2006) and two were quasi-experimental (Davis,

Kaptein, Kaplan, Olmsted, & Woodside 1998; Klein, Mayer, Schebendach, & Walsh, 2007). Several studies involved a retrospective case series design, by investigating connections between various features of the eating disorder patients (Kron et al., 1978; Penas-Lledo et al., 2002; Solenberger, 2001). Finally, one study adopted a daily process design (Vansteelandt, Rijmen, Pieters, Probst, & Vanderlinden, 2007).

The majority of studies utilised one of two methods for gathering data. Self report measures were used either on their own, or as part of an overall assessment. Structured interviews were the other key method used to identify key features and qualitative experiences. Objective measures of height and weight were taken in several studies, whilst physical activity was assessed using objective measures in only one study (Klein et al., 2007).

From this point on, the outcome of the systematic review and the subsequent broader review of the non-clinical and exercise science fields will be combined to provide a rationale for inclusion of the key constructs within the ensuing model.

Development of an empirical model of compulsive exercise in the eating disorders

The reviews outlined above revealed several key indices of the compulsive dimension of exercise that are consistently described. Each of these indices can be viewed from a cognitive behavioural perspective as constituting potential maintenance factors for compulsive exercise. These key factors are schematically presented in Fig 1. This review will now critically consider the evidence supporting these factors in the maintenance of compulsive exercise.

Figure 1. A schematic representation of the maintenance model of compulsive exercise

1 Links between compulsive exercise and eating pathology

Among eating disordered samples, increased compulsive exercise has been shown to be significantly associated with higher levels of eating disordered symptoms (Shroff et al., 2006), dietary restraint, weight and shape concerns (Dalle Grave, Calugi, & Marchesini, 2008; Holtkamp, Hebebrand, & Herpertz-Dahlman, 2004), drive for thinness (Solenburger, 2001), and body dissatisfaction (Brewerton et al., 1995; Solenburger, 2001). Likewise, among non-eating disordered samples, the relationship between compulsive exercise and elevated scores on measures of eating psychopathology is similarly well established (e.g., Elbourne & Chen, 2007; Lipsey, Barton, Hulley, & Hill, 2006). For example, compulsive exercise has been shown to be associated with higher scores on measures of eating and weight restriction (Thome & Espelage, 2007), drive for thinness (Ackard, Brehm, & Steffen, 2002; Adkins & Keel, 2005; Gulker, Laskis, & Kuba, 2001; Matheson & Crawford-Wright,

2000; Steffen & Brehm, 1999), bulimic tendencies (Ackard et al., 2002; Adkins & Keel, 2005; Steffen & Brehm, 1999), and body dissatisfaction (Ackard, Brehm, & Steffen, 2002; Adkins & Keel, 2005; Gulker, Laskis, & Kuba, 2001; Steffen & Brehm, 1999). In addition, Mond and colleagues (Mond, Hay, Rodgers, Owen, & Beumont, 2004;2006a; Mond, Hay Rodgers, & Owen, 2006b; Mond, Myers, Crosby, Hay, & Mitchell, 2008; Mond & Calogero, 2009) have consistently demonstrated that the extent to which exercise is undertaken solely or primarily to change weight or shape is one of the dimensions of compulsive exercise most strongly associated with eating disorder psychopathology and reduced quality of life. Indeed, such compulsive exercise was shown to clearly differentiate between eating disordered and non-eating disordered samples (Mond & Calogero, 2009).

Furthermore, it has been noted that behaviours such as dieting and exercise tend to reinforce each other because of the many personal and social benefits that accrue (Davis, 1997). Indeed, several studies have shown that exercising for weight loss is one of the principal reasons reported by men and women for beginning an exercise program (Davis et al., 1995; Markland & Hardy, 1993; McDonald & Thompson, 1992). However, worryingly, exercise can foster weight and diet concerns in certain susceptible individuals because the initial weight loss that often accompanies regular exercise elicits certain social reinforcers which, in turn, may lead to a heightened interest in physical appearance (Katz, 1986). This is in contrast to the prevalent social-cognitive/cognitive-behavioural perspective that views compulsive exercise as symptomatic of, rather than antecedent to, an eating disorder (Davis, Fox, Cowles, Hastings, & Schwass, 1990; Fairburn, Cooper, & Shafran, 2003) and suggests a reciprocally reinforcing relationship between compulsive exercise and eating pathology.

The importance of the links between exercise and eating pathology are underpinned by compelling empirical support for weight and shape concerns being a central feature of compulsive exercise, such that in the absence of eating pathology, compulsive exercise is not considered to constitute a clinically significant syndrome (Keski-Rahkonen, 2001; Mond et al., 2006a). Indeed, Mond et al. (2006a) demonstrated that there was no association

between exercise and reduced quality of life when eating pathology was statistically controlled for.

The evidence presented above clearly supports eating pathology as a central component of compulsive exercise both in terms of its development and maintenance. Critically, the evidence also suggests that the relationship between compulsive exercise and eating pathology is not one sided. Rather, each is suggested to reciprocally reinforce the other. However, despite the central importance of eating pathology in the development and maintenance of compulsive exercise, it is now recognised that compulsive exercise is more than simply a strategy for wasting energy or burning calories in weight preoccupied individuals (Davis, Katzman, & Kirsh, 1999). Whilst weight and shape concerns remain an essential component of compulsive exercise (e.g., Hubbard, Gray, & Parker, 1998; Keski-Rahkonen, 2001; Mond et al., 2006a), there is evidence to suggest that this model is too simplistic. Indeed, a review of the literature clearly suggests other key factors in addition to weight and shape concerns that may also be contributing to the maintenance of compulsive exercise. One such factor is the regulation of negative affect (e.g., Thome & Espelage, 2004), which will now be considered.

2 Links between affect regulation and compulsive exercise

Researchers and clinicians have long recognised eating disorders as, in part, ineffective coping mechanisms used to manage uncomfortable emotions (Heilbrun & Harris, 1986; Slade, 1982; Soukup, Beiler, & Terrell, 1998). Indeed, links between affect regulation and eating psychopathology are well established (e.g., Killen, Taylor, Hayward, Wilson, Haydel et al., 1996; Meyer, Waller & Waters, 1998). Given the affect regulatory function of restriction and bulimic behaviours (Fairburn, 1997; Fairburn, Cooper, & Shafran, 2003; Hawkins & Clement, 1984; Heatherton & Baumeister, 1991; Meyer et al., 1998), a plausible model would suggest that compulsive behaviours (such as exercise) serve a similar regulatory function in the eating disorders; indeed this was the conclusion reached by

Lawson, Waller and Lockwood (2007). This notion is supported by the fact that, in general, compulsivity is associated with a wide range of negative emotions (Spinella, 2005).

Within the exercise science literature, the affect regulatory role of exercise has been well established (Callaghan, 2004), with mood regulation being one of the most frequently endorsed motivations for exercising (e.g., Markland & Ingledew, 1997). As has been pointed out by several researchers, any behaviour that is under voluntary control and that can influence mood can be used to regulate emotions (e.g., Jackson, Cooper, Mintz, & Albino, 2003; Parkinson & Totterdell, 1999; Sher & Trull, 1994). The suggestion that physical activity may be used as a compensatory behaviour not just to burn calories but to alleviate or counteract dysphoric mood states, is further supported by frequent patient reports that exercise is often being maintained primarily for the control of negative affect as much as it is to expend calories (Blumenthal, O'Toole, & Chang, 1984; Davis & Woodside, 2002; Long et al., 1993).

Certainly among eating disordered patients, compulsive exercise has been shown to be associated with elevated levels of negative affect (e.g., Brewerton et al., 1995; Penas-Lledo et al., 2002; Vansteelandt et al., 2007; Vansteelandt, Vandereycken, Claes, Probst, & Mechelen, 2004). Furthermore, the association between drive for thinness and physical activity is most pronounced in those patients with the highest levels of chronic negative affect (Vansteelandt et al., 2007).

Similar associations between compulsive exercise and increased levels of negative affect have also been found in non-eating disordered samples (e.g., Siegel & Hetta, 2001; Spano, 2001). For example, in those with high levels of eating pathology, exercise is associated with negative affect (Thome & Espelage, 2004). Conversely, among females with relatively low eating pathology, exercise is associated with positive affect (Thome & Espelage, 2004). This not only suggests that exercise might be differentially associated with affect based on the presence or absence of eating pathology (Thome & Espelage, 2004), but also supports the previously proposed model that exercise does not constitute a clinically

significant syndrome in the absence of eating pathology (Keski-Rahkonen, 2002; Mond et al., 2006a).

There is also evidence that increased exercise is associated with specific negative emotional states. For example, in both anorexia nervosa (AN) and bulimia nervosa (BN) patients, compulsive exercise has consistently been found to be associated with elevated ratings of anxiety (Brewerton et al., 1995; Klein, Bennett, Schebendach, Foltin, Devlin & Walsh, 2004; Penas-Lledo et al., 2002). The anxiolytic properties of strenuous physical activity are well established in both animal and human studies (e.g., Binder, Droste, Ohl, & Reul, 2004; Cox, Thomas, Hinton, & Donahue, 2004; Greenwood, Foley, Day, Burhans, Brooks et al., 2005). To date, there have been over 35 published reviews dealing with the anxiolytic effects of exercise providing consistent empirical support for the anxiolytic effects of exercise in clinical and non-clinical populations (Callaghan, 2004; Salmon, 2001; Scully, Kremer, Meade, Graham & Dudgeon, 1998). Indeed, it has been repeatedly shown that exercise can reduce anxiety symptoms in healthy adults (e.g., Hale & Raglin, 2002), in anxious women (e.g., Breus & O'Connor, 1998), and in psychiatric patients (Broocks, Bandelow, Pekrun, George, Meyer et al., 1998). In one study, among patients in the acute phase of AN, physical activity levels were predicted only by the level of food restriction and anxiety symptoms, supporting the notion that physical activity may be a way of coping with anxiety in some patients (Holtkamp et al., 2004).

In addition to anxiety, compulsive exercise has been shown to be associated with higher levels of depression in eating disorder patients (Penas-Lledo et al., 2002) and in non-clinical samples (Seigel & Hetta, 2001; Thome & Espelage, 2004). In addition, exercise has been shown to significantly decrease depressive symptoms following a single bout of exercise (Maraki, Tsofliou, Pitsiladis, Malkova, Mutrie & Higgins, 2005) and in regular exercisers (Steinberg, Nicholls, Sykes, LeBoutillier, Ramlakham et al., 1998). Meta-analyses of studies examining the efficacy of exercise in the prevention and/or alleviation of depression have consistently found that there is consistent empirical support for the efficacy of exercise in preventing and alleviating the symptoms of depression in clinically and non-

clinically depressed populations. This anti-depressive effect is largely independent of the frequency, mode, or intensity of exercise – and that exercise is as effective as alternative treatments for depression (e.g., Craft & Landers, 1998; North, McCullagh & Tran, 1990). Similarly, several more recent reviews have also concluded that exercise is associated with decreased depression and is effective at reducing depression in clinically and non-clinically depressed populations (Callaghan, 2004; Phillips, Kiernan & King, 2003).

The suggestion that compulsive exercise may function as a means of compensating for or suppressing adverse emotional states in ED patients is not new (e.g., Geller, Cockell & Goldner, 2000; Markland & Ingledew, 1997; Penas-Lledo et al., 2002). Fairburn et al. (2003) suggest that the inability to cope appropriately with adverse mood states, such as anxiety and depression is a central component in the maintenance of the eating disorders. Termed “mood intolerance”, it is characterised by engagement in various dysfunctional mood modulatory behaviours, such as compulsive exercising, that serve the purpose of neutralising the mood state as well as reducing awareness of the associated cognitions (Fairburn et al., 1993). They further suggest that these mood modulatory behaviours may then become habitual forms of mood modulation. This also supports the suggestion that compulsive exercise may be functioning as a mood modulatory behaviour in eating disordered patients. The use of exercise as a mood modulatory behaviour may not be so pronounced among bulimic patients as they exhibit a range of behaviours (e.g., bingeing, purging, self-harm, or drug abuse) that serve the same function more effectively (McManus & Waller, 1995). The idea that compulsive exercise may become a habitual method of affect regulation is supported by evidence that among eating disordered patients who exercise compulsively, exercise may become the primary and even the only means by which they regulate their emotional state (Geller et al., 2000; Markland & Ingledew, 1997).

In summary, the functional utility of exercise in regulating emotion further suggests a functional equivalence with eating pathology. It is probable that an inability to cope appropriately with adverse mood states results in exercise being maintained as an affect regulation strategy. It follows, therefore, that affect regulation may constitute a further

maintenance factor for compulsive exercise. Moreover, given the assumption that the use of exercise to regulate affect can become habitual, it is little wonder that the concepts of exercise dependence and/or exercise addiction have been proposed to explain the observed *reliance* on exercise among eating disordered patients to regulate their emotional state (Adams & Kirkby, 2002; Klein et al., 2004).

Within the literature there are two plausible mechanisms for the maintenance of the observed *reliance on exercise* that characterise some eating disordered patients. These mechanisms are positive and negative reinforcement. This notion is compatible with the 'addiction' model of compulsive exercise. For example, Davis (2000) postulates that in order for exercise to have the potential for addiction, it has to be either a positively or negatively reinforcing activity. In other words, it must either increase pleasure or ameliorate distress and suffering. Indeed, on the nature of addiction, Goodman (1990) notes that the behaviour indulged in (albeit no longer able to be controlled) is of a kind that can function not only to produce pleasure, but also to provide escape (avoidance/alleviate) from internal discomfort.

Positive reinforcement

The positive reinforcement function of exercise is reflected in the positive addiction model of exercise dependence which proposed that exercise dependence was attributable to exercise-induced euphoric states that were highly enjoyable (Glasser, 1976). However, strenuous exercising of the type typical of compulsive exercise is not a primary reinforcing behaviour for most people since relatively few individuals engage in exercise for the euphoria it produces (Davis, 2000).

Whilst there is evidence that exercise is associated with positive affect (Thome & Espelage, 2004), which no doubt is a positively reinforcing factor, other studies have demonstrated that doing exercise for enjoyment is by far the least endorsed reason among men and women (e.g., Davis et al., 1995). In particular, there is strong evidence that compulsive exercise is primarily maintained by secondary reinforcing factors of which weight and shape concerns are the strongest predictors (Brehm & Steffen, 1998; Davis, Brewer, &

Ratusny, 1993) followed by affect regulation (Geller et al., 2000; Markland & Ingledrew, 1997).

Supporting the affect regulatory model of compulsive exercise is the model of exercise dependence whereby exercise is maintained primarily to mitigate the affective withdrawal symptoms that manifest when unable to exercise (e.g., Bamber, Cockerill, Rodgers, & Carroll, 2003; Hausenblas & Symons Downs, 2002a; Morgan, 1979; Veale, 1987;1995), essentially constituting a form of negative reinforcement.

Negative reinforcement (withdrawal symptoms)

Of all the various definitional and diagnostic criteria proposed for exercise dependence (e.g., Hausenblas & Symons Downs, 2002b; Veale, 1995), the largest body of evidence is for the experience of affective withdrawal symptoms as a consequence of being unable to exercise (e.g., Bamber, Cockerill, Rodgers, & Carroll, 2003). Indeed, Aidman and Woollard (2003) make the point that exercise dependence is characterised by a pattern of withdrawal symptoms. Based on the premise that in some exercisers the cessation of regular exercise may result in negative psychological states (Hausenblas & Symons Downs, 2002a;b), withdrawal symptoms have been posited as the cardinal characteristic of exercise dependence (Szabo, 1995).

Withdrawal symptoms have been described as the psychological effects that occur as a result of not exercising, and are manifested either by emotional symptoms, or further exercise to avoid such symptoms (American Psychiatric Association, 1994). The most frequently reported symptoms of withdrawal were guilt, anxiety, depression, and irritability (Bamber, Cockerill, & Carroll, 2000; Szabo, Frenkl, & Caputo, 1997). It has been suggested that as little as 24 hours without exercise may be sufficient to produce significant withdrawal symptoms in some individuals (e.g., Sachs, 1981; Thaxton, 1982).

Whilst the experience of withdrawal symptoms is common to many regular exercisers, it has been suggested that only some can be classed as dependent (Szabo, 1995). However, in a recent study attempting to delineate diagnostic criteria, withdrawal

symptoms emerged as one of only two criteria implying dependence (Bamber et al., 2003). It has certainly been suggested that habitual exercisers may learn to rely on exercise to counteract anxiety or depression, control mood states, or lose weight, and that a subsequent inability to exercise results in negative psychological consequences in the form of withdrawal symptoms (Baekeland, 1970; Morris, Steinberg, Sykes, & Salmon, 1990; Thaxton, 1982). Indeed, such exercise-related dysphoria has even been shown to discriminate between eating disordered and non-clinical groups (Boyd, Abraham, & Luscombe, 2007).

Withdrawal symptoms are known to be universally present at, if not responsible for, the onset of most addictions, including opiates, nicotine and alcohol (e.g., Galanter, 1993; Niaura, Rohsenow, Binkoff, & Monti, 1988; Powel, Bradley, & Gray, 1992) and the withdrawal phenomenon is considered a common consequence of dependence (Farrell, 1994). From the earliest descriptions, emotional withdrawal symptoms have been consistently reported as a consequence of exercise deprivation among habitual exercisers (e.g., Aidman & Woollard, 2003; Baekeland, 1970; Chan & Grossman, 1988; Conboy, 1994; Mondin et al., 1996; Morgan, 1979; Morris et al., 1990; Ogles, Masters, & Richardson, 1995; Veale, 1987; Yates, Leehey, & Shisslak, 1983).

Whilst it has been suggested that withdrawal can take the forms of physiological and/or psychological disturbance (Sachs & Pargman, 1979; West & Gossop, 1994), the evidence for a physiological basis to exercise dependence remains equivocal (Adams & Kirkby, 2002; Davis, 2000; Hamer & Karageorghis, 2007). Indeed, the model currently most supported by the empirical literature suggests that the presence of affective withdrawal symptoms is indicative of a *psychological* dependence on exercise whereby exercise is maintained via negative reinforcement specifically to avoid the experience of such withdrawal.

In summary, evidence has been presented suggesting that among compulsive exercisers, the experience of affective withdrawal symptoms when unable to exercise is common in both clinical and non-clinical samples (Szabo, 1995). In some cases this may result in a psychological dependency/addiction upon exercise whereby exercise behaviour is

maintained specifically to avoid the withdrawal symptoms (i.e., negative reinforcement). In addition, for some individuals there is potentially a further powerful maintenance factor of positive reinforcement as a result of elevated positive affect (i.e., positive addiction) (Glasser, 1976). Taken together, the evidence discussed so far supports the notion that in some susceptible individuals (i.e., those who are mood intolerant), engaging in exercise may inadvertently result in a psychological dependence, which then reciprocally maintains the exercise behaviour.

3 Compulsivity and the maintenance of exercise

It is plausible to suggest that *addictive* behaviours and *compulsive* behaviours are essentially similar, since they are both characterised by a need to perform a behaviour that is no longer intrinsically rewarding. The following section will focus on the compulsivity element of exercise in more detail.

Compulsivity describes an insistent urge to perform some action (i.e., behaviour) to relieve the *anxiety* stemming from a fear of perceived negative consequences if the behaviour is not performed (American Psychiatric Association, 2000; Frost, Steketee, Cohn, & Griess, 1994). Therefore, the primary function of compulsive behaviours is to mitigate *anxiety* via the *avoidance* of perceived negative consequences. Furthermore, such compulsive behaviours are typically characterised by being performed persistently, repetitively and excessively, without leading to any reward or pleasure (Evans, Lewis, & Iobst, 2004).

Compulsivity has been shown to be significantly related to levels of activity in both eating disordered patients and non-eating disordered exercisers (Gulker et al., 2001; Davis et al., 1995; Shroff et al., 2006). Furthermore, eating disorder symptomatology has been shown to be associated with compulsive attitudes to exercise, rather than just exercise quantity (Adkins & Keel, 2005; Seigel & Hetta, 2000). Indeed, excessive exercising has been described as one of the most obvious compulsions in the eating disorders with a lifetime occurrence in up to 84% of patients (Davis, 1997).

Supporting the importance of the avoidance of perceived negative consequences component of compulsivity, it has been demonstrated that compulsive exercise is associated with heightened concerns about the consequences of not exercising (Bamber et al., 2000; Penas-Lledo, Sancho, & Waller, 2002). For example, avoidance of failure has been identified as a maintaining factor for compulsive exercise (Hall, Kerr, Kozub, & Finnie, 2007). In addition, a tendency towards avoidance (i.e., an avoidant coping style) has been found to be associated with disordered eating attitudes and behaviours in both eating disordered and non-eating disordered women (Sherwood, Crowther, Wills, & Ben-Porath, 2000; VanBoven & Espelage, 2006). More specifically, avoidance of emotional arousal has been shown to be associated with increased compulsivity in both eating disordered (Lawson, Waller, & Lockwood, 2007) and non-eating disordered samples (Abramowitz, Lackey, & Wheaton, 2009). It is plausible that the experience of affective withdrawal symptoms when unable to exercise leads to a fear of such negative consequences, resulting in compulsive exercise behaviour in order to avoid experiencing such withdrawal. Exercise behaviour therefore becomes maintained by the need to avoid the experience of the withdrawal component of compulsivity. This suggestion that a fear of the negative consequences of not exercising is a maintaining factor for compulsive exercise is not new (Penas-Lledo, Sancho, & Waller, 2002; Thaxton, 1982) and supports the centrality of compulsivity in the maintenance of compulsive exercise.

In addition to the evidence described above, there are also established links between low body weight and compulsivity. For example, Crisp (1967) concluded that starvation facilitates generalised compulsive behaviour so that exercising, like various other activities, can involuntarily become stereotyped, ritualised, and excessive, simply as a consequence of the disorder. Supporting this, serotonin dysregulation, a frequent consequence of malnutrition, has been implicated in the development of compulsive behaviours (Barr, Goodman, Price, McDougle, & Charney, 1992; Holden, 1990; Kaye, Weltzin, & Hsu, 1993). The links between starvation and compulsivity can readily be observed within the context of anorexia nervosa. Specifically, it is frequently observed in AN that as the condition worsens,

patients feel an increasingly strong compulsion to be physically active even when they no longer enjoy it, and continuing even when the process is painful and exhausting (Davis, 1997). Indeed, AN patients showed both a higher level of physical activity and more compulsive exercise than controls (Davis, Blackmore, Katzman, & Fox, 2005).

Obsessive-compulsive disorder and obsessive-compulsive personality disorder

Both ED and compulsive exercise share a strong relationship with obsessionality. For example, it has been suggested that an inherent obsessionality may underlie both compulsive exercise and eating pathology (Davis et al., 1998; Gulker, Laskis, & Kuba, 2001) and several studies report that compulsive exercisers typically display an obsessional and rigid personality profile as well as a compulsive behaviour pattern (Davis, Brewer, & Ratusny, 1993; Goldfarb & Plante, 1984; Kagan, 1987; Yates, 1991). As a result, compulsive exercise is often viewed as being related to obsessive-compulsiveness (Davis & Claridge, 1998; Davis et al., 1995;1998;1999). Consequently, much of the research into compulsive exercise has tended to focus on its links with obsessive-compulsive personality traits and obsessive-compulsive symptomatology (Penas-Lledo et al., 2002). Indeed, it has even been suggested that it may be useful to consider compulsive exercise as stemming from an anxiety disorder (Gulker et al., 2001), which fits with the idea that the function of compulsive exercise in the ED is to reduce anxiety, much like the function of compulsions in obsessive-compulsive disorder (Davis & Kaptein, 2006).

The existing literature clearly suggests a significant relationship between AN and obsessive-compulsive personality traits (Serpell, Livingstone, Neiderman, & Lask, 2002), with evidence to support a more obsessive-compulsive personality profile in exercising AN patients than their non-exercising counterparts (Davis & Claridge, 1998; Davis et al., 1998; Davis & Kaptein, 2006). Furthermore, both anorexic and bulimic patients classified as compulsive exercisers had higher scores on measures of obsessive-compulsive personality than their non-exercising counterparts (Davis & Claridge, 1998). As a result, obsessive-compulsive personality factors are presumed to be probable risk factors for both disordered

eating (Wonderlich, Lilenfeld, Riso, Engel, & Mitchell, 2005) and compulsive exercising (Gulker et al., 2001).

Guilt

Guilt has been shown to be associated with obsessive-compulsive symptomatology in both clinical and non-clinical samples (e.g., Frost et al., 1994; Mancini, Gangemi, Perdighe, & Marini, 2008; Shafran, Watkins, & Charman, 1996). Indeed, guilt has been shown to predict compulsivity independent of anxiety and depression (Shafran, Watkins, & Charman, 1996) and fear of guilt has been linked to increased rates of compulsive behaviours (Mancini & Gangemi, 2006). In addition, guilt has been implicated as an aetiological factor in the development of obsessive compulsive disorder (OCD) (Gibbs, 1996).

Mond and colleagues (2004a;2006;2008;2009) have consistently demonstrated that the degree of guilt (feeling bad if one has been prevented from exercising a certain amount) experienced whenever exercise is unable to be undertaken is one of the components of compulsive exercise most strongly associated with eating disorder psychopathology and reduced quality of life. The importance of guilt is further supported by the finding that guilt when unable to exercise discriminated between ED and non-ED groups (Boyd, Abraham, & Luscombe, 2007; Mond & Calogero, 2009). Indeed, the experience of guilt has consistently been reported as a consequence of being unable to exercise among compulsive exercisers (Bamber et al., 2000; Szabo et al., 1997).

In summary, compulsivity is linked with both increased exercise and eating disorders. Specifically, the avoidance of perceived negative consequences and the experience of guilt when exercise is unable to be undertaken have been shown to be the components of compulsivity most strongly implicated in the maintenance of compulsive exercise.

Taken together, the evidence presented so far clearly supports compulsivity as an important factor in the maintenance of compulsive exercise and further supports the conception of problematic exercise as a compulsive behaviour.

4 Perfectionism and rigidity in the maintenance of compulsive exercise

Of all the obsessive-compulsive personality traits associated with disordered eating such as perfectionism, inflexibility, rigidity, and preoccupation with interpersonal and mental control (Halmi, 2005), it is perfectionism that is posited as the most strongly associated with both compulsive exercising (Flett & Hewitt, 2005) and disordered eating (Cassidy, Allsopp & Williams, 1999; Halmi et al., 2000). In addition, rigidity has long been described as a feature of compulsive exercise (e.g., Beumont et al., 1994; Wyatt, 1997; Yates, 1991) and a cluster analysis of personality pathology in ED patients found that nearly half of patients were included in a cluster characterised by rigid compulsivity (Goldner, Srikameswaran, Schroeder, Livesley, & Birmingham, 1999).

Perfectionism

Perfectionism typically describes a personality trait characterised by having and striving for high personal standards (Frost, Marten, Lahart, & Rosenblate, 1990), and is well established as an important factor in the development and maintenance of the eating disorders (Fairburn, Cooper, Doll, & Welch, 1999; Fairburn et al., 2003; Fairburn, Shafran, & Cooper, 1999; Lilenfeld, Wonderlich, Riso, Crosby, & Mitchell, 2006; Stice, 2002). In terms of compulsive exercise, recent evidence suggests that perfectionism is a robust correlate of compulsive exercise (e.g., Shroff et al., 2006). Indeed, compulsive exercise has been shown to be consistently associated with higher levels of perfectionism in both eating disordered and non-eating disordered samples (Ackard et al., 2002; Brehm & Steffen, 1998; Coen & Ogles, 1993; Cook 1996; Symons Downs, Hausenblas, & Nigg, 2004; Gulker et al., 2001; Hagan & Hausenblas, 2003; Hall et al., 2007; Hausenblas & Symons Downs, 2002b; Seigel & Hetta, 2001; Shroff et al., 2006).

The importance of perfectionism is further highlighted by the finding that AN patients classified as compulsive exercisers report significantly higher levels of perfectionism than their non-exercising counterparts (Davis et al., 1998). Perfectionism has been shown to be

independent of the state of nutrition in AN (Kaye, 1997), and in a sample of compulsively exercising AN patients, elevated perfectionism was associated with poor outcome from the treatment of their compulsive exercise (Long & Hollin, 1995). This supports the notion that perfectionism is a significant factor in compulsive exercise and furthermore, may be an important maintenance factor in its own right (i.e., independent of compulsivity).

Despite the preceding evidence, understanding of the relationship between perfectionism and compulsive exercise is impaired by the use of disparate conceptualizations of perfectionism (e.g., Frost et al., 1990; Hewitt & Flett, 1991). For example, empirical findings have suggested that whilst high personal standards remain an undisputed component of perfectionism they are not necessarily, by themselves, pathological (Dunkley, Blankstein, Halsall, Williams & Winkworth, 2000; Frost et al., 1990; Serpell, Waller, Fearon, & Meyer, 2009). Indeed, recent findings have consistently demonstrated the existence of two distinct, albeit related, dimensions of perfectionism (e.g., Bieling, Israeli, & Antony, 2004; Blankstein & Dunkley, 2002). Whereas the first dimension reflects the setting and striving for high personal standards, the second reflects a tendency towards self-criticism (Dunkley, Blankstein, Masheb, & Grilo, 2006) and it is the self-critical dimension that is posited as the primary indicator of the dysfunctional component of perfectionism (Dunkley et al., 2006).

Unfortunately, no studies examining the relationship between perfectionism and compulsive exercise have specifically considered the self-critical dimension of perfectionism. However, despite the lack of studies explicitly considering the two proposed dimensions of perfectionism, there is evidence that both are associated with compulsive exercise. For example, whilst not explicitly considered, many current perfectionism measures have been shown to tap both dimensions (e.g., Bieling et al., 2004; Dunkley, Zuroff, & Blankstein, 2003). An examination of studies using these measures with measures of compulsive exercise clearly supports an association between compulsive exercise and both the high personal standards and self-critical dimensions of perfectionism (e.g., Cook 1996; Symons Downs et al., 2004; Shroff et al., 2006). In addition, compulsive exercisers are reported to

often strive to achieve perfection with any failure to reach this standard resulting in self-criticism (Beumont et al., 1994) and Davis (1997) has posited a co-occurrence of self-criticism and high personal standards as a potential risk factor for compulsive exercise.

Rigidity

Rigidity has been defined as a pattern of resisting "...the acquisition of new behaviour patterns by holding onto previous and non-adaptive styles of performance" (Schaie & Parman, 1975: p. 1) and is regarded as an essential component of dysfunctional perfectionism (Shafran & Mansell, 2001). Indeed, Ferrari and Mautz (1997) previously demonstrated an association between rigidity and perfectionism, and a recent qualitative analysis of perfectionism identified rigidity as a consistent theme implicated in its maintenance (Riley & Shafran, 2005), thereby supporting previous descriptions of dysfunctional perfectionism (e.g., Burns, 1980). Endorsing rigidity as a significant component of compulsive exercise is the fact that it remains central to the definition of obsessive-compulsive personality disorder, which has been demonstrated to have a significant association with both ED (Halmi, 2005) and compulsive exercise (Davis & Claridge, 1998; Davis et al., 1999; Serpell et al., 2002).

In terms of compulsive exercise, rigidity has been linked to the emotional consequences of not meeting personal standards. Specifically, it is suggested that repetitive behaviour helps filter out both extraneous stimuli and disruptive affect-laden cognitions (Rauch & Savage, 2000). Rauch and Savage (2000) proposed that repetitive behaviours in the anxious state represent an attempt to homeostatically reduce anxiety-caused increased levels of arousal by soothing limbic tone. Because compulsive exercisers often express their exercise in a stereotyped and repetitive manner (e.g., Beumont et al., 1994; Wyatt, 1997; Yates, 1991), it has been suggested that this characteristic of their exercise further supports the anxiolytic effect of the exercise (Holtkamp et al., 2004). Indeed, Bamber, Cockerill, Rodgers and Carroll (2003) found that inflexible attitudes towards exercising was an important signifier of compulsive exercise resulting in inflexible and stereotyped exercise

behaviour. Furthermore, Boyd et al. (2007) found that the setting of rigid exercise rules was a significant predictor of an ED diagnosis.

In summary, the evidence above clearly supports perfectionism as a significant component of compulsive exercise. Evidence was presented supporting perfectionism as a maintenance factor for compulsive exercise in the ED. Similarly, evidence was presented that suggests rigidity may also be a central component in the maintenance of compulsive exercise, likely stemming from its anxiolytic function and association with perfectionism. Whilst both perfectionism and rigidity overlap with the construct of compulsivity the evidence supports their conceptualisation as separate constructs in terms of maintaining compulsive exercise

Conclusions

This review has presented a novel conceptualisation of compulsive exercise within the context of the eating disorders. Several key constructs have been identified that are empirically supported as being potentially maintaining of exercise within this domain. Future research into compulsive exercise should take into account these factors when providing working definitions of exercise with the eating disorders field.

References:

- Abramowitz, J.S., Lackey, G.R., & Wheaton, M.G. (2009). Obsessive-compulsive symptoms: The contribution of obsessional beliefs and experiential avoidance. *Journal of Anxiety Disorders, 23*, 160-166.
- Ackard, D.M., Brehm, B.J., & Steffen, J.J. (2002). Exercise and eating disorders in college-aged women: Profiling excessive exercisers. *Eating Disorders, 10*(1), 31-47.
- Adams, J., & Kirkby, R.J. (2002). Excessive exercise as an addiction: A review. *Addiction Research and Theory, 10*(5), 415- 437.
- Adkins, E.C. & Keel, P.K. (2005). Does “excessive” or “compulsive” best describe exercise as a symptom of bulimia nervosa? *International Journal of Eating Disorders, 38*, 24-29.
- Aidman, E.V. & Woollard, S. (2003). The influence of self-reported exercise addiction on acute emotional and physiological responses to brief exercise deprivation. *Psychology of Sport and Exercise, 4*(3), 225-236.
- American Psychiatric Association. (1994). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.) (DSM-IV). Washington, DC: American Psychiatric Association.
- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed., text revision) (DSM-IV-TR). Washington, DC: American Psychiatric Association.
- Atlantis, E., & Baker, M. (2008). Obesity effects on depression: systematic review of epidemiological studies. *International Journal of Obesity, 32*, 881-891.
- Baekeland, F. (1970). Exercise deprivation: Sleep and psychological reactions. *Archives of General Psychiatry, 22*, 365–369.
- Bamber, D., Cockerill, I.M., & Carroll, D. (2000). The pathological status of exercise dependence. *British Journal of Sports Medicine, 34*, 125-132.
- Bamber, D., Cockerill, I.M., Rodgers, S., & Carroll, D. (2003). Diagnostic criteria for exercise dependence in women. *British Journal of Sports Medicine, 37*, 393–400.

- Barr, L.C., Goodman, W.K., Price, L.H., McDougle, C.J., & Charney, D.S. (1992). The serotonin hypothesis of obsessive-compulsive disorder: Implications of pharmacologic challenge studies. *Journal of Clinical Psychiatry*, 53, 17-28.
- Berkman, N. D., Lohr, K. N., & Bulik, C. M. (2007). Outcomes of eating disorders: a systematic review of the literature. *IJED*, 40, 293-309.
- Beumont, P.J.V., Arthur, B., Russell, J.D., & Touyz, S.W. (1994). Excessive physical-activity in dieting disorder patients - proposals for a supervised exercise program. *International Journal of Eating Disorders*, 15, 21-36.
- Bieling, P.J., Israeli, A.L., & Antony, M.M. (2004). Is perfectionism good, bad, or both? Examining models of the perfectionism construct. *Personality and Individual Differences*, 36, 1373-1385.
- Binder, E., Droste, S.K., Ohl, F., & Reul, J.M.H.M. (2004). Regular voluntary exercise reduces anxiety-related behaviour and impulsiveness in mice. *Behavioural Brain Research*, 155, 197-206.
- Blankstein, K.R., & Dunkley, D.M. (2002). Evaluative concerns self-critical and personal standards perfectionism: A structural equation modelling strategy. In G.L. Flett, & P.L. Hewitt (Eds.), *Perfectionism: Theory, research and treatment* (pp. 285-315). Washington, DC: American Psychological Association.
- Blaydon, M.J., Linder, K.J., & Kerr, J.H. (2004). Metamotivational characteristics of exercise dependence and eating disorders in highly active amateur sport participants. *Personality and Individual Differences*, 36(6), 1419-1432.
- Blumenthal, J.A., O'Toole, L.C., & Chang, J.L. (1984). Is running an analogue of anorexia nervosa? An empirical study of obligatory running and anorexia nervosa. *Journal of the American Medical Association*, 252(4), 520-523.
- Boyd, C., Abraham, S., & Luscombe, G. (2007). Exercise behaviours and feelings in eating disorder and non-eating disorder groups. *European Eating Disorders Review*, 15, 112-118.

- Brehm, B.J. & Steffen, J.J. (1998). Relation between obligatory exercise and eating disorders. *American Journal of Health Behavior*, 22, 1008-1019.
- Breus, M.J., & O'Connor, P.J. (1998). Exercise-induced anxiolysis: A test of the "time out" hypothesis in high anxious females. *Medical Science in Sports and Exercise*, 30, 1107–1112.
- Brewerton, T.D., Stelfox, E.J., Hibbs, N., Hodges, E.L., & Cochrane, C.E. (1995). Comparison of eating disorder patients with and without compulsive exercising. *International Journal of Eating Disorders*, 17(4), 413-416.
- Broocks, A., Bandelow, B., Pekrun, G., George, A., Meyer, T., Bartmann, U., et al. (1998). Comparison of aerobic exercise, clomipramine, and placebo in the treatment of panic disorder. *American Journal of Psychiatry*, 155, 603–609.
- Bruch, H. (1965). Anorexia nervosa – its differential diagnosis. *Journal of Mental and Nervous Disorders*, 141, 555-566.
- Burns, D.D. (1980). The perfectionist's script for self defeat. *Psychology Today*, November, 34-51.
- Callaghan, P. (2004). Exercise: a neglected intervention in mental health care? *Journal of Psychiatric and Mental Health Nursing*, 11, 476-483.
- Carter, J.C., Blackmore, E., Sutandar-Pinnock, K., & Woodside, D.B. (2004). Relapse in anorexia nervosa: a survival analysis. *Psychological Medicine*, 34, 671-679.
- Casper, R.C., & Jabine, L.N. (1996). An Eight-Year Follow-Up: Outcome from Adolescent Compared to Adult Onset Anorexia Nervosa. *Journal of Youth and Adolescence*, 25, 499-517.
- Cassidy, E., Allsopp, M., & Williams, T. (1999). Obsessive compulsive symptoms at initial presentation of adolescent eating disorders. *European Child & Adolescent Psychiatry*, 8, 193–199.
- Chan, C. S., & Grossman, H. Y. (1988). Physiological effects of running loss on consistent runners. *Perceptual and Motor Skills*, 66, 875–883.

- Coen, S.P., & Ogles, B.M. (1993). Psychological characteristics of the obligatory runner - a critical-examination of the anorexia analogue hypothesis. *Journal of Sport & Exercise Psychology, 15*, 338-354.
- Conboy, J.K. (1994). The effects of exercise withdrawal on mood states in runners. *Journal of Sport Behavior, 17*, 188-2-3.
- Cook, C. A. (1996). The psychological correlates of exercise dependence in aerobics instructors. Unpublished master's thesis, University of Alberta, Alberta, Canada.
- Cox, R.H., Thomas, T.R., Hinton, P.S., & Donahue, O.M. (2004). Effects of acute 60 and 80% $\text{VO}_{2\text{max}}$ bouts of aerobic exercise on state anxiety of women of different age groups across time. *Research Quarterly for Exercise and Sport, 75*, 165-175.
- Craft, L.L. & Landers, D.M. (1998). The effect of exercise on clinical depression and depression resulting from mental illness: a meta-analysis. *Journal of Sport and Exercise Psychology, 20*, 339-357.
- Crisp, A.H. (1967). The possible significance of some behavioural correlates of weight and carbohydrate intake. *Journal of Psychosomatic Research, 11*, 117-131.
- Crisp, A.H., Hsu, L.K.G., Harding, B., & Hartshorn, J. (1980). Clinical features of anorexia nervosa: A study of a consecutive series of 102 female patients. *Journal of Psychosomatic Research, 24*, 179-191.
- Dalle Grave, R., Calugi, S., & Marchesini, G. (2008b). Compulsive exercise to control shape or weight in eating disorders: prevalence, associated features, and treatment outcome. *Comprehensive Psychiatry, 49*, 346-352.
- Dally, P.J. (1969). *Anorexia nervosa*. New York: Grune & Stratton.
- Davis, C. (1997). Eating disorders and hyperactivity: a psychobiological perspective. *Canadian Journal of Psychiatry, 42*, 168-175.
- Davis, C. (2000). Exercise abuse. *International Journal of Sport Psychology, 31*, 278-289.
- Davis, C., Blackmore, E., Katzman, D., & Fox, J. (2005). Female adolescents with anorexia nervosa and their parents: A case-control study of exercise attitudes and behaviours. *Psychological Medicine, 35*, 377-386.

- Davis, C., Brewer, H., & Ratusny, D. (1993). Behavioral frequency and psychological commitment - necessary concepts in the study of excessive exercising. *Journal of Behavioral Medicine*, 16, 611-628.
- Davis, C., & Claridge, G. (1998). The eating disorders as addiction: A psychobiological perspective. *Addictive Behaviors*, 23, 463-475.
- Davis, C., Fox, J., Cowles, M.P., Hastings, P., & Schwass, K. (1990). The functional role of exercise in the development of weight and diet concerns in women. *Journal of Psychosomatic Research*, 34, 563-574.
- Davis, C., & Kaptein, S. (2006). Anorexia nervosa with excessive exercise: A phenotype with close links to obsessive-compulsive disorder. *Psychiatry Research*, 142, 209-217.
- Davis, C., Kaptein, S., Kaplan, A.S., Olmsted, M.P., & Woodside, D.B. (1998). Obsessionality in anorexia nervosa: The moderating influence of exercise. *Psychosomatic Medicine*, 60, 192-197.
- Davis, C., Katzman, D.K., Kaptein, S., Kirsh, C., Brewer, H., & Kalmbach, K., Olmstead, M.P., Woodside, D.B., & Kaplan, A.S. (1997). The prevalence of high-level exercise in the eating disorders: Etiological implications. *Comprehensive Psychiatry*, 38, 321-326.
- Davis, C., Katzman, D.K., & Kirsh, C. (1999). Compulsive physical activity in adolescents with anorexia nervosa - A psychobehavioral spiral of pathology. *Journal of Nervous and Mental Disease*, 187, 336-342.
- Davis, C., Kennedy, S.H., Ravelski, E. & Dionne, M. (1994). The role of physical activity in the development and maintenance of eating disorders. *Psychological Medicine*, 24, 957-967.
- Davis, C., Kennedy, S.H., Ravelski, E., Dionne, M., Brewer, H., & Neitzert, C., & Ratusny, D. (1995). Obsessive compulsiveness and physical activity in anorexia nervosa and high-level exercising. *Journal of Psychosomatic Research*, 39, 967-976.
- Davis, C. & Woodside, D.B. (2002). Sensitivity to the Rewarding Effects of Food and Exercise in the Eating Disorders. *Comprehensive Psychiatry*, 43, 189-194.

- Dunkley, D.M., Blankstein, K.R., Halsall, J., Williams, M., & Winkworth, G. (2000). The relation between perfectionism and distress: Hassles, coping, and perceived social support as mediators and moderators. *Journal of Counselling Psychology*, 47, 437-453.
- Dunkley, D.M., Blankstein, K.R., Masheb, R.M., & Grilo, C.M. (2006). Personal standards and evaluative concerns dimensions of “clinical” perfectionism: A reply to Shaffran et al. (2002, 2003) and Hewit et al. (2003). *Behaviour Research and Therapy*, 44, 63-84.
- Dunkley, D.M., Zuroff, D.C., & Blankstein, K.R. (2003). Self-critical perfectionism and daily affect: dispositional and situational influences on stress and coping. *Journal of Personality and Social Psychology*, 84, 234–252.
- Eisler, I., & le Grange, D. (1990). Excessive exercise and anorexia nervosa. *International Journal of Eating Disorders*, 9, 377-386.
- Elbourne, K.E., & Chen, J. (2007). The continuum model of obligatory exercise: A preliminary investigation. *Journal of Psychosomatic Research*, 62, 73-80.
- Epling, W.F., & Pierce, W.D. (1992). *Solving the anorexia puzzle*. Toronto: Hogrefe & Huber Publishers.
- Evans, D.W., Lewis, M.D., & Lobst, E. (2004). The role of the orbitofrontal cortex in normally developing compulsive-like behaviours and obsessive-compulsive disorder. *Brain and Cognition*, 55, 220-234.
- Fairburn, C.G. (1997). Eating disorders. In D.M. Clark & C.G. Fairburn (Eds.), *Science and practice of cognitive behaviour therapy* (pp. 209-241). Oxford: Oxford University Press.
- Fairburn, C.G., Cooper, Z., Doll, H.A., & Welch, S.L. (1999). Risk factors for anorexia nervosa: Three integrated case-control comparisons. *Archives of General Psychiatry*, 56, 468-476.
- Fairburn, C.G., Cooper, Z. & Shafran, R. (2003). Cognitive behaviour therapy for eating disorders: a “transdiagnostic” theory and treatment. *Behaviour Research and Therapy*, 41, 509-528.

- Fairburn, C.G., Shafran, R., Cooper, Z. (1999). A cognitive-behavioural theory of anorexia nervosa. *Behaviour Research and Therapy*, 37, 1-13.
- Farrell, M. (1994). Opiate withdrawal. Special issue: Comparing drugs of dependence. *Addiction*, 89, 1471-1475.
- Ferrari, J.R., & Mautz, W.T. (1997). Predicting perfectionism: Applying tests of rigidity. *Journal of Clinical Psychology*, 53, 1–6.
- Flett, G.L., & Hewitt, P.L. (2005). The perils of perfectionism in sports and exercise. *Current Directions in Psychological Science*, 14, 14-18.
- Frost, R.O., Marten, P.A., Lahart, C., & Rosenblate, R. (1990). The dimensions of perfectionism. *Cognitive Therapy and Research*, 14, 449-468.
- Frost, R.O., Steketee, G., Cohn, L., & Griess, K.E. (1994). Personality traits in subclinical and non-obsessive compulsive volunteers and their parents. *Behaviour, Research and Therapy*, 32, 47-56.
- Galanter, M. (1993). Network therapy for addiction: a model for office practice. *American Journal of Psychiatry*, 150, 28–36.
- Geller, J., Cockell, S.J., & Goldner, E.M. (2000). Inhibited expression of negative emotions and interpersonal orientation in anorexia nervosa. *International Journal of Eating Disorders*, 28, 8-19.
- Gibbs, N.A. (1996). Nonclinical populations in research on obsessive-compulsive disorder: A critical review. *Clinical Psychology Review*, 16, 729-773.
- Glasser, W. (1976). *Positive addiction*. New York: Harper and Row.
- Goldfarb, L.A., & Plante, T.G. (1984). Fear of fat in runners: an examination of the connection between anorexia nervosa and distance running. *Psychological Reports*, 55, 296.
- Goldner, E.M., Sriameswaran, S., Schroeder, M.L., Livesley, W.J., Birmingham, C.L. (1999). Dimensional assessment of personality pathology in patients with eating disorders. *Psychiatry Research*, 85, 151-159.

- Goodman, A. (1990). Addiction: Definition and implications. *British Journal of Addiction*, 85(11), 1402-1408.
- Greenwood, B.N., Foley, T.E., Day, H.E.W., Burhans, D., Brooks, L., Campeau, S., & Fleshner, M. (2005). Wheel running alters serotonin (5-HT) transporter, 5-HT_{1A}, 5-HT_{1B} and alpha_{1b}-adrenergic receptor mRNA in the rat raphe nuclei. *Biological Psychiatry*, 57, 559-568.
- Gulker, M.G., Laskis, T.A., & Kuba, S.A. (2001). Do excessive exercisers have a higher rate of obsessive-compulsive symptomatology? *Psychology, Health and Medicine*, 6, 387-398.
- Gull, W.W. (1874). Anorexia nervosa (apepsia hysterica, anorexia hysterica). *Transactions of the Clinical Society of London*, 7, 22-28.
- Hagan, A.L. & Hausenblas, H.A. (2003). The relation between exercise dependence symptoms and perfectionism. *American Journal of Health Studies*, 18, 133-137.
- Hale, B.S., & Raglin, J.S. (2002). State anxiety responses to acute resistance training and step aerobic exercise across eight weeks of training. *Journal of Sports Medicine and Physical Fitness*, 42, 108–112.
- Hall, H.K., Kerr, A.W., Kozub, S.A., & Finnie, S.B. (2007). Motivational antecedents of obligatory exercise: The influence of achievement goals and multidimensional perfectionism. *Psychology of Sport and Exercise*, 8, 297-316.
- Halmi, K.A. (2005). Obsessive-compulsive personality disorder and eating disorders. *Eating Disorders: The Journal of Treatment & Prevention*, 13, 85-92.
- Halmi, K. A., Sunday, S. R., Strober, M., Kaplan, A., Woodside, D. B., Fichter, M., Treasure, J., Berretini, W.H., & Kaye, W.H. (2000). Perfectionism in anorexia nervosa: Variation by clinical subtype, obsessionality, and pathological eating behavior. *American Journal of Psychiatry*, 157(11), 1799-1805.
- Hamer, M., & Karageorghis, C. (2007). Psychobiological mechanisms of exercise dependence. *Sports medicine*, 37, 477-84.

- Hausenblas, H. A., & Downs, D. S. (2002a). Exercise dependence: A systematic review. *Psychology of Sport and Exercise*, 3, 89-123.
- Hausenblas, H.A. & Downs, D.S. (2002b). How much is too much? The development and validation of the exercise dependence scale. *Psychology and Health*, 17, 387-404.
- Hawkins R.C., & Clement P.F. (1984). Binge eating: measurement problems and a conceptual model. In R.C. Hawkins, W.J. Fremoun, & P.F.Clement (Eds.), *The Binge-Purge Syndrome*, (pp. 229-253). Springer, New York. Herman C.
- Heatherton, T.F., & Baumeister, R.F. (1991). Binge eating as escape from self-awareness. *Psychological Bulletin*, 110, 86-108.
- Hechler, T., Beumont, P., Marks, P., & Touyz, S. (2005). How do clinical specialists understand the roles of physical activity in eating disorders. *European Eating Disorders Review*, 13, 125-132.
- Heilbrun, A.B., & Harris, A. (1986). Psychological defences in females at-risk for anorexia nervosa: An explanation for excessive stress found in anorexic patients. *International Journal of Eating Disorders*, 5, 503-516.
- Hewitt, P.L., & Flett, G.L. (1991). Perfectionism in the self and social contexts: conceptualization, assessment and association with psychopathology. *Journal of Personality and Social Psychology*, 60, 456-470.
- Holden, N.L. (1990). Is anorexia nervosa an obsessive-compulsive disorder? *British Journal of Psychiatry*, 157, 1-5.
- Holtkamp, K., Hebebrand, J., & Herpertz-Dahlmann, B. (2004). The contribution of anxiety and food restriction on physical activity levels in acute anorexia nervosa. *International Journal of Eating Disorders*, 36, 163-171.
- Hubbard, S.T., Gray, J.J., & Parker, S. (1998). Differences among women who exercise for 'food related' and 'non-food related' reasons. *European Eating Disorders Review*, 6, 255-265.
- Inches, P. (1895). Anorexia nervosa. *Maritime Medical News*, 7, 73-75.

- Jackson, B., Cooper, M.L., Mintz, L., & Albino, A. (2003). Motivations to eat: Scale development and validation. *Journal of Research in Personality*, 37, 297-318.
- Kagan, D. (1987). Addictive personality factors. *Journal of Psychology*, 121, 533-538.
- Katz, J.L. (1986). Long-distance running, anorexia nervosa and bulimia: A report of two cases. *Comprehensive Psychiatry*, 27, 74-78.
- Kaye, W.H. (1997). Anorexia nervosa: Obsessional behavior, and serotonin. *Psychopharmacology Bulletin*, 33, 335-344.
- Kaye, W.H., Weltzin, T.E., & Hsu, L.K.G. (1993). Relationship between anorexia nervosa and obsessive and compulsive behaviours. *Psychiatric Annals*, 23, 365-373.
- Keski-Rahkonen, A. (2001). Exercise dependence - a myth or a real issue? *European Eating Disorders Review*, 9, 279-283.
- Killen, J.D., Taylor, C.B., Hayward, C., Wilson, D., Haydel, K., Hammer, L., et al. (1994). Pursuit of thinness and onset of eating disorder symptoms in a community sample of adolescent girls: A three year prospective analysis. *International Journal of Eating Disorders*, 16, 227-238.
- Klein, D.A., Mayer, L.E.S., Schebendach, J.E., & Walsh, B.T. (2007). Physical activity and cortisol in Anorexia Nervosa. *Psychoneuroendocrinology*, 32(5), 539-547.
- Klein, D.A., Bennett, A.S., Schebendach, J., Foltin, R.W., Devlin, M.J., & Walsh, B.T. (2004). Exercise "addiction" in anorexia nervosa: Model development and pilot data. *CNS Spectrums*, 9, 531-537.
- Kron, L., Katz, J.L., Gorzynski, G., & Weiner, H. (1978). Hyperactivity in anorexia nervosa: A fundamental clinical feature. *Comprehensive Psychiatry*, 19, 433-439.
- Lasegue, E.C. (1873). De l'anorexie hysterique. *Archives Generales de Medecine*, 1, 384-403. In: *Evolution of Psychosomatic Concepts. Anorexia Nervosa: A Paradigm*, (eds. R.M. Kaufmann and M. Heiman). New York: International Universities Press, 1964.
- Lawson, R., Waller, G., & Lockwood, R. (2007). Cognitive content and process in eating disordered patients with obsessive-compulsive features. *Eating Behaviors*, 8, 305-310.

- Lilenfeld, L.R., Wonderlich, S., Riso, L.P., Crosby, R., & Mitchell, J. (2006). Eating disorders and personality: A methodological and empirical review. *Clinical Psychology Review*, 26, 299-320.
- Lipsey, Z., Barton, S.B., Hulley, A., & Hill, A.J. (2006). After a workout..." Beliefs about exercise, eating and appearance in female exercisers with and without eating disorder features. *Psychology of Sport and Exercise*, 7, 425-436
- Long, C.G., & Hollin, C.R. (1995). Assessment and management of eating disordered patients who overexercise: A four-year follow-up of six single case studies. *Journal of Mental Health*, 4, 309-317.
- Long, C.G., Smith, J., Midgley, M., & Cassidy, T. (1993). Over-exercising in anorexic and normal samples: Behaviour and attitudes. *Journal of Mental Health*, 2, 321-327.
- Mancini, F., & Gangemi, A. (2006). The role of responsibility and fear of guilt in hypothesis testing. *Journal of Behavior Therapy and Experimental Psychiatry*, 37, 333-346.
- Mancini, F., Gangemi, A., Perdighe, C., & Marini, C. (2008). Not just right experience: Is it influenced by feelings of guilt. *Journal of Behavior Therapy and Experimental Psychiatry*, 39, 162-176.
- Maraki, M., Tsofliou, F., Pitsiladis, Y.P., Malkova, D., Mutrie, N., & Higgins, S. (2005). Acute effects of a single exercise class on appetite, energy intake, and mood. Is there a time of day effect? *Appetite*, 45, 272-278.
- Markland, D. & Hardy, L. (1993). The exercise motivations inventory: preliminary development and validity of a measure of individuals' reasons for participation in regular physical exercise. *Personality and Individual Differences*, 15, 289-296.
- Markland, D. & Ingledrew, D.K. (1997). The measurement of exercise motives: Factorial validity and invariance across gender of a revised Exercise Motivations Inventory. *British Journal of Health Psychology*, 2, 361-376.
- Matheson, H. & Crawford-Wright, A. (2000). An examination of eating disorder profiles in student obligatory and non-obligatory exercisers. *Journal of Sport Behavior*, 23, 42-50.

- McDonald, K. & Thompson, J.K. (1992). Eating disturbance, body image dissatisfaction, and reasons for exercising: gender differences and correlational findings. *International Journal of Eating Disorders*, 11, 289-292.
- McManus, F., & Waller, G. (1995). A functional analysis of binge-eating. *Clinical Psychology Review*, 8, 845-863.
- Meyer, C., Waller, G., & Waters, A. (1998). Emotional states and bulimic psychopathology. In H.W. Hoek, J.L. Treasure, & M.A. Katzman (Eds.), *Neurobiology in the treatment of eating disorders* (pp. 271-287). Chichester: Wiley.
- Mond, J. M. & Calogero, R.M. (2009). Excessive exercise in eating disorder patients and healthy women. *The Australian and New Zealand Journal of Psychiatry*, 49, 227-234.
- Mond, J.M., Hay, P.J., Rodgers, B., & Owen, C. (2006). An update on the definition of “excessive exercise” in eating disorders research. *International Journal of Eating Disorders*, 39, 147-153.
- Mond, J.M., Hay, P.J., Rodgers, B., Owen, C., & Beumont, P.J. (2004a). Relationships between exercise behaviour, eating disordered behaviour and quality of life in a community sample of women: When is exercise excessive? *European Eating Disorders Review*, 12, 265-272.
- Mond, J. M., Hay, P. J., Rodgers, B., Owen, C., & Beumont, P. J. V. (2004b). Validity of the eating disorder examination questionnaire (EDE-Q) in screening for eating disorders in community samples. *Behaviour Research and Therapy*, 42(5), 551-567.
- Mond, J.M., Myers, T.C., Crosby, R., Hay, P., & Mitchell, J. (2008). Excessive exercise and eating-disordered behaviour in young adult women: further evidence from a primary care sample. *European Eating Disorders Review*, 16, 215-221.
- Mondin, G.W., Morgan, W.P., Piering, P.N., Stegner, A.J., Stotesbery, C.L., Trine, M.R., & Wu, M.Y. (1996). Psychological consequences of exercise deprivation in habitual exercisers. *Medicine and Science in Sports and Exercise*, 28, 1199–1203.
- Morgan, W.P. (1979). Negative addiction in runners. *Physician and Sports Medicine*, 7, 57-70.

- Morris, M., Steinberg, H., Sykes, E. A., & Salmon, P. (1990). Effects of temporary withdrawal from regular running. *Journal of Psychosomatic Research*, 34, 493–500.
- Niaura, R., Rohsenow, D., Binkoff, J., & Monti, P. (1988). Relevance of cue reactivity to understanding alcohol and smoking relapse. *Journal of Abnormal Psychology*, 97, 133–152.
- North, T.C., McCullagh, P., & Tran, Z.V. (1990). Effects of exercise on depression. *Exercise and Sport Science Reviews*, 18, 379-415.
- Ogles, B.M., Masters, K.S., & Richardson, S.A. (1995). Obligatory running and gender: an analysis of participative motives and training habits. *International Journal of Sport Psychology*, 26, 233–248.
- Parkinson, B., & Totterdell, P. (1999). Classifying affect-regulation strategies. *Cognition and Emotion*, 13, 277–303.
- Penas-Lledo, E., Vaz Leal, F.J., & Waller, G. (2002). Excessive Exercise in Anorexia Nervosa and Bulimia Nervosa: Relation to Eating Characteristics and General Psychopathology. *International Journal of Eating Disorders*, 31, 370-375.
- Phillips, W.T., Kiernan, M., & King, A.C. (2003). Physical Activity as a Non-pharmacological Treatment for Depression: A Review. *Complementary Health Practice Review*, 8, 139-152.
- Powell, J., Bradley, B., & Gray, J. (1992). Classical conditioning and cognitive determinants of subjective craving for opiates: An investigation of their relative contributions. *British Journal of Addiction*, 87, 1133–1144.
- Rauch, S.L. & Savage, C.R. (2000). Investigating cortico-striatal pathophysiology in obsessive-compulsive disorders: Procedural learning and imaging probes. In W.K. Goodman, M.V. Rudorfer, & J.D. Maser (Eds.), *Obsessive-compulsive disorder* (pp. 133–154). Mahwah, NJ: Lawrence Erlbaum Associates, Inc.
- Riley, C. & Shafran, R. (2005). Clinical perfectionism: A preliminary qualitative analysis. *Behavioural and Cognitive Psychotherapy*, 33, 369-374.

- Sachs, M.L. (1981). Running addiction. In: Sacks, M. and Sachs, M. (Eds.), *Psychology of Running*, pp. 116–126. Human Kinetics, Champaign, Ill.
- Sachs, M.L., & Pargman, D. (1979). Running addiction: A depth interview examination. *Journal of Sport Behavior*, 2, 143-155.
- Salmon, P. (2001). Effects of physical exercise on anxiety, depression, and sensitivity to stress: A unifying theory. *Clinical Psychology Review*, 21, 33-61.
- Schaie, K.W., & Parham, I.A. (1975). *The test of behavioral rigidity*. Palo Alto, CA: Consulting Psychologists Press.
- Scully, D., Kremer, J., Meade, M.M., Graham, R., & Dedgeon, K. (1998). Physical exercise and psychological well-being: A critical review. *British Journal of Sports Medicine*, 32, 111-120.
- Seigel, K. & Hetta, J. (2001). Exercise and eating disorder symptoms among young females. *Eating and Weight Disorders*, 6, 32-39.
- Serpell, L., Livingstone, A., Neiderman, M., & Lask, B. (2002). Anorexia nervosa: Obsessive-compulsive disorder, obsessive-compulsive personality disorder, or neither? *Clinical Psychology Review*, 22, 647-669.
- Serpell, L., Waller, G., Fearon, P., & Meyer, C. (2009). The roles of persistence and perseveration in psychopathology. *Behaviour Therapy*, 40, 260-271.
- Shafran, R., & Mansell, W. (2001). Perfectionism and psychopathology: a review of research and treatment. *Clinical Psychology Review*, 21, 879–906.
- Shafran, R., Watkins, E., & Charman, T. (1996). Guilt in obsessive-compulsive disorder. *Journal of Anxiety Disorders*, 10, 509-516.
- Sher, K.J., & Trull, T.J. (1994). Personality and disinhibitory psychopathology: Alcoholism and antisocial personality disorder. *Journal of Abnormal Psychology*, 103, 92–102.
- Sherwood, N.E., Crowther, J.H., Wills, L., & Ben-Porath, Y.S. (2000). The perceived function of eating for bulimic, subclinical bulimic and non-eating disordered women. *Behaviour Therapy*, 31, 777-793.

- Shroff, H., Reba, L., Thornton, L.M., Tozzi, F., Klump, K.L., Berrettini, W.H., et al. (2006). Features Associated With Excessive Exercise in Women with Eating Disorders. *International Journal of Eating Disorders*, 39, 454-461.
- Slade, P. (1982). Toward a functional analysis of anorexia nervosa and bulimia nervosa. *British Journal of Clinical Psychology*, 21, 167-179.
- Solenberger, S.E. (2001). Exercise and eating disorders: A 3-year inpatient hospital records analysis. *Eating Behaviors*, 2, 151-168.
- Soukup, V.M., Beiler, E.M., & Terrell, F. (1998). Stress, coping style, and problem-solving ability among eating disordered inpatients. *Journal of Clinical Psychology*, 46, 592-599.
- Spano, L. (2001). The relationship between exercise and anxiety, obsessive-compulsiveness, and narcissism. *Personality and Individual Differences*, 30, 87-93.
- Spinella, M. (2005). Mood in relation to subclinical obsessive-compulsive symptoms. *International Journal of Neuroscience*, 115, 433-443.
- Steffen, J.J., & Brehm, B.J. (1999). The dimensions of obligatory exercise. *Eating Disorders*, 7, 219-226.
- Steinberg, H., Nicholls, B.R., Sykes, E.A., LeBoutillier, N., Ramlakham, N., Moss, T.P., & Dewey, A. (1998). Weekly Exercise Consistently Reinstates Positive Mood. *European Psychologist*, 3, 271-280.
- Stice, E. (2002). Risk and Maintenance Factors for Eating Pathology: A Meta-Analytic Review. *Psychological Bulletin*, 128(5), 825-848.
- Strober, M., Freeman, R., & Morrell, W. (1997). The long-term course of severe anorexia nervosa in adolescents: Survival analysis of recovery, relapse, and outcome predictors over 10-15 years in a prospective study. *International Journal of Eating Disorders*, 22, 339-360.
- Symons Downs, D., Hausenblas, H.A., & Nigg, C.R. (2004). Factorial validity and psychometric examination of the Exercise Dependence Scale-Revised. *Measurement in Physical Education and Exercise Science*, 8, 183-201.

- Szabo, A. (1995). The impact of exercise deprivation on well-being of habitual exercisers. *The Australian Journal of Science and Medicine in Sport*, 27, 68-75.
- Szabo, A., Frenkl, R., & Caputo, A. (1997). Relationships between addiction to running, commitment to running, and deprivation from running: A study on the internet. *European Yearbook of Sport Psychology*, 1, 130-147.
- Thaxton, L. (1982). Physiological and psychological effects of short term exercise addiction on habitual runners. *Journal of Sport Psychology*, 4, 73–80.
- Thien, V., Thomas, A., Markin, D., & Birmingham, C.L. (2000). Pilot study of a graded exercise program for the treatment of anorexia nervosa. *International Journal of Eating Disorders*, 28, 101-106.
- Thome, J.L., & Espelage, D.L. (2004). Relations among exercise, coping, disordered eating, and psychological health among college students. *Eating Behaviors*, 5, 337-351.
- Thome, J.L., & Espelage, D.L. (2007). Obligatory exercise and eating pathology in college females: Replication and development of a structural model. *Eating Behaviors*, 8, 334-349.
- Touyz, S.W., Beumont, P.J.V., & Hoek, S. (1987). Exercise anorexia: A new dimension in anorexia nervosa? In P. J. V. Beumont, G. D. Burrows, & R. C. Caspar (Eds.), *Handbook of eating disorders: Part 1. Anorexia and bulimia nervosa* (pp. 143-157). Amsterdam: Elsevier.
- VanBoven, A., & Espelage, D. L. (2006). Influence of depression on coping assessment and disordered eating among college females. *Journal of Counselling and Development*, 84, 341-348.
- Vansteelandt, K., Pieters, G., Vandereycken, W., Claes, L., Probst, M., & Van Mechelen, I. (2004). Hyperactivity in anorexia nervosa: A case study using experience sampling methodology. *Eating Behaviors*, 5, 67–74.
- Vansteelandt, K., Rijmen, F., Pieters, G., Probst, M., & Vanderlinden, J. (2007). Drive for thinness, affect regulation and physical activity in eating disorders: A daily life study. *Behaviour, Research and Therapy*, 45, 1717-1734.

- Veale, D. (1987). Exercise dependence. *British Journal of Addiction*, 82, 735-740.
- Veale, D. (1995). Does primary exercise dependence really exist? In J. Annett, B. Cripps & H. Steinberg (Eds.), *Exercise addiction: Motivation for participation in sport and exercise*. (pp. 1-5). Leicester, UK: British Psychological Society.
- Waller, G. (2002). Eating attitudes and the use of alcohol, tobacco, and exercise among male and female adolescents. *Eating Behaviors*, 3, 101-111.
- West, R. & Gossop, M. (1994). Overview: A comparison of withdrawal symptoms from different drug classes. *Addiction*, 89, 1483-1489.
- Wonderlich, S. A., Lilienfeld, L. R., Riso, L. P., Engel, S., & Mitchell, J. E. (2005). Personality and anorexia nervosa. *International Journal of Eating Disorders*, 37, S68-S71.
- Wyatt, L.M. (1997). *Obsessive-Compulsiveness and Disordered Eating in Obligatory and Non-Obligatory Exercisers*. (PhD ed.). Los Angeles: California School of Professional Psychology.
- Yates, A. (1991). *Compulsive exercise and the eating disorders*. New York: Brunner/Mazel.
- Yates, A., Leehey, K., & Shisslak, C.M. (1983). Running: an analogue of anorexia? *New England Journal of Medicine*, 308, 251–255.