This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Erratum: effects of lattice geometry and interaction range on polaron dynamics

PLEASE CITE THE PUBLISHED VERSION

PUBLISHER

© American Physical Society

VERSION

VoR (Version of Record)

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Hague, J.P., P.E. Kornilovitch, A.S. Alexandrov, and J.H. Samson. 2019. "Erratum: Effects of Lattice Geometry and Interaction Range on Polaron Dynamics". figshare. <https://hdl.handle.net/2134/4274>.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Erratum: Effects of lattice geometry and interaction range on polaron dynamics [Phys. Rev. B 73, 054303 (2006)]

J. P. Hague, P. E. Kornilovitch, A. S. Alexandrov, and J. H. Samson
(Received 5 December 2006; published 2 February 2007)

DOI: [10.1103/PhysRevB.75.059901](https://doi.org/10.1103/PhysRevB.75.059901)

PACS number(s): 71.38.-k, 02.70.Ss, 74.20.-z, 99.10.Cd

We have recently discovered a mistake in the strong coupling, long-range interaction results in this paper, changing Figs. 9–12. The corrected figures are shown below. We also note that γ computed by Eq. (23) should read 0.32(0) for the triangular lattice and 0.33(4) for the square lattice with Fröhlich interaction. We note that the Lang-Firsov limit and spectra now agree even better with our numerical results at large λ (>5) and ω , Fig. 10. Numerical masses of lattice Fröhlich polarons (Fig. 9) in the intermediate coupling and adiabaticity regimes are instrumental for better understanding of transport properties and angle-resolved photo-emission spectra of cuprate superconductors, where the finite-range Fröhlich interaction with c -axis polarized optical phonons is important.^{1,2} The conclusions of the paper are unchanged.

FIG. 9. Inverse mass of the Fröhlich polaron on the square and triangular lattices when $\bar{\omega}=1$. Inset: approaching the strong coupling limit at large λ .

FIG. 10. Dispersion for a screened Fröhlich polaron with $R_{sc}=1$, $\lambda=8$, and $\bar{\omega}=8$, compared with the Lang-Firsov (LF) result to first order in $1/\lambda$.

FIG. 11. DOS for a screened Fröhlich polaron with $R_{sc}=1$, $\lambda=8$, and $\bar{\omega}=8$, compared with the Lang-Firsov result to first order in $1/\lambda$.

FIG. 12. Ratio of the effective bandwidths of the screened Fröhlich polaron on square and triangular lattices. $\lambda=1.4$ and $\bar{\omega}=1$.

¹A. S. Alexandrov, Phys. Rev. B **53**, 2863 (1996); Phys. Rev. Lett. **82**, 2620 (1999).

²W. Meevasana, N. J. C. Ingle, D. H. Lu, J. R. Shi, F. Baumberger, K. M. Shen, W. S. Lee, T. Cuk, H. Eisaki, T. P. Devereaux, N. Nagaosa, J. Zaanen, and Z.-X. Shen, Phys. Rev. Lett. **96**, 157003 (2006).