This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Response of benthic cave invertebrates to organic pollution events

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1002/aqc.933>

PUBLISHER

© John Wiley & Sons

VERSION

SMUR (Submitted Manuscript Under Review)

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Wood, Paul J., J. Gunn, and S.D. Rundle. 2019. "Response of Benthic Cave Invertebrates to Organic Pollution Events". figshare. <https://hdl.handle.net/2134/13019>.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Response of benthic cave invertebrates to organic pollution events

WOOD, P.J.^{1*}, GUNN, J.² and RUNDLE, S.D.³

1. *Department of Geography, Loughborough University, Loughborough, Leicestershire, LE11 3TU, United Kingdom.*

2. *Limestone Research Group, School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom.*

3. *School of Biological Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, United Kingdom.*

Corresponding Author *

Dr P.J. Wood
Department of Geography
Loughborough University
Loughborough
Leicestershire
LE11 3TU
United Kingdom

Email:- p.j.wood@lboro.ac.uk

Tel:- 00 (44) +1509 223012

Fax:- 00 (44) +1509 223930

Abstract

1. Even though the fragility and vulnerability of subterranean ecosystems (caves, groundwater and hyporheic habitats) is widely acknowledged, the impacts of anthropogenic disturbances have been poorly quantified when compared to surface waters. In particular, limited data exist regarding the impact of organic pollution upon aquatic cave invertebrate communities.
2. The Peak-Speedwell Cavern system (Derbyshire, UK) was affected by two organic pollution events, during a 7-year study (1997-2003), originating from the same source in the surface catchment but resulting in markedly different ecological responses. The first event led to the elimination of most taxa from affected sites while the second resulted in an increase in abundance of organisms within the cave associated with the increased availability of trophic resources. The second event also coincided with the invasion of the stygophilic amphipod, *Gammarus pulex*, at a site where it had not previously been recorded.
3. Recovery of the invertebrate community following both organic pollution events occurred within 12-months. Recolonisation of the affected sites was facilitated by annual flooding of the cave and by the presence of refugia on unaffected subterranean tributaries.
4. The data highlight the problems associated with the conservation and management of subterranean ecosystems where impacts in distant surface catchments may have unseen repercussions for the subterranean environment. Aquatic subterranean habitats are not widely monitored and the impacts of pollution/disturbance may not be detected in surface waters for some time, if at all, due to dilution effects. Caves supporting obligate subterranean organisms (stygobites) are particularly vulnerable to these pressures and require clear management strategies to protect both the subterranean and surface catchments which support them.

Key Words:- subterranean ecosystem; disturbance; recovery; point source pollution; invertebrates.

Introduction

The dark zones of caves are naturally typified by low organismal abundance and diversity (Holsinger, 1988; Jasinska et al., 1996; Culver and Sket, 2000), and due to the relatively constant abiotic conditions, their biological communities are widely considered to be more stable compared with those in epigean systems (Culver, 1985; Simon et al., 2003). In the absence of light and primary producers, cave habitats are largely oligotrophic, relying almost exclusively on dissolved or particulate organic matter originating in surface (epigean) habitats (Poulson and Lavoie, 2000; Simon et al., 2003). Hence, it is highly likely that changes in landuse and/or management practices within surface epigean catchments may result in significant changes to the trophic dynamics of subterranean (hypogean) food webs (Poulson and Lavoie, 2000; Hancock et al., 2005). Consequently, aquatic subterranean habitats (hyporheic zone, groundwater and wet caves) are considered to be vulnerable to anthropogenic activities (Sket, 1999; Gunn et al., 2000; van Beynen and Townsend, 2005; Boulton, 2005), yet our understanding of the impacts of such activities upon subterranean ecosystems is much more limited compared with epigean waterbodies (Elliott, 2000; Hancock et al., 2005).

Although there has been some recent increase in interest regarding the impact of disturbances upon biological communities within groundwater aquifers (e.g., Danielopol et al., 2003; Hancock, 2002), research exploring the influence of disturbances upon cave ecosystems has been limited despite wide recognition of their high conservation/biodiversity value (Culver and Sket, 2000). Anthropogenic disturbances and modifications of cave ecosystems associated with heavy metals (Graening and Brown, 2003), faecal bacteria (Green et al., 1990; Simon and Buikema, 1997; Graening and Brown, 2003) and waste disposal (Halliday, 2003)

81 have been reported. However, the response of cave communities and individual species to
82 organic pollution remains poorly quantified. This paucity of information reflects the absence
83 of pre-disturbance baseline data and/or absence of adjacent control sites which could be used
84 to determine the nature and magnitude of impacts. Those studies that have documented the
85 response of cave invertebrate communities and individual species to organic enrichment and
86 pollution show that responses are variable (Table 1). Significant changes to the structure of
87 cave benthic invertebrate communities, particularly reductions in abundance or exclusion of
88 obligate subterranean aquatic fauna (stygobites) as a result of organic pollution have been
89 reported (Culver et al., 1992, Simon and Builkema, 1997, Graening and Brown, 2003).
90 However, in some instances there have been increases in the abundance of obligate
91 subterranean (hypogean/stygobitic) fauna and/or an increase in species richness of
92 epigean/stygophilic faunal populations within caves, particularly when trophic resource
93 availability is enhanced (Holsinger, 1966, Sket, 1977, Simon and Builkema, 1997, Graening
94 and Brown, 2003). It has even been suggested that mild organic enrichment may be beneficial
95 to stygobitic populations under some circumstances, provided that highly competitive
96 stygophiles, epigean taxa able to complete their life-cycles within the cave but usually
97 occurring in surface waters, do not invade (Sket, 1999, Graening and Brown, 2003).
98
99 Pollution of groundwater dominated habitats has been implicated as one of the greatest threats
100 to the long term provision of groundwater resources and subterranean biodiversity (e.g.,
101 Boulton, 2005; Danielopol et al., 2003; Hancock et al., 2005) and in particular, cave ecology
102 (e.g., Gunn et al., 2000; Finlay et al., 2006; Panno et al., 2006). However, data clearly
103 demonstrating the ecological impact of pollution within caves are limited due to the
104 difficulties associated with conducting research within subterranean habitats, the absence of
105 pre-disturbance (pollution) data and/or information regarding the source and nature of
106 pollutants (Gunn et al., 2000). Here we examine the response of freshwater cave invertebrates

to two point-source organic pollution events that occurred during a seven-year study (1997-2003). Our main aims were to gauge the impact of pollution by: (1) quantifying the invertebrate community response to pollution episodes and comparing the impacts of separate events; and (2) investigating changes to the local populations (i.e. extinctions or invasions) resulting from pollution.

Methods

Study Site

The study was undertaken from 1997-2003 within the Peak-Speedwell Cave system, Derbyshire (UK). Peak Cavern and Speedwell Cavern are interconnected and contain more than 16 km of active (wet) and relict (dry) cave passages that have formed within Carboniferous limestone (karst geology). There is limited hydrological connectivity between the caves, except under high flow (flood) conditions, when water from Speedwell Cavern may rise into the higher passages within Peak Cavern. Water within Peak Cavern is largely derived from autogenic sources (water that has only been in contact with limestone bedrock and overlying soil, and percolates into the cave) which are concentrated into two main subterranean streams that enter the cave from flooded conduits, Ink Sump and Far Sump. These streams flow along the Peak Cavern streamway, enter another flooded conduit and emerge as Peak Cavern Rising, a large spring at the head of Peakshole Water (Figure 1). Water within Speedwell Cavern is largely derived from allogenic sources - twelve streams that flow on the surface over non-limestone geologies before sinking underground. The streams combine underground, enter Speedwell Cavern via two flooded conduits, Main Rising and Whirlpool Rising, flow through the cave, enter another flooded conduit and finally emerge from two springs, Russet Well and Slop Moll, which both flow into Peakshole Water (Gunn et al., 2000). Landuse in both catchments is dominated by livestock grazing,

which has historically resulted in inputs of faecal bacteria to the subterranean ecosystem (Gunn et al., 1998, Hunter et al., 1999).

Detection of pollution and tracing the source

Two major point source pollution episodes were experienced during the study period: (1) during early 1999; and (2) between December 2001-January 2002. Both events occurred when parts of the cave were inaccessible due to flooding and as a result the passage of the pollutant could not be directly monitored *in situ*. Following the detection of pollution within Peak Cavern due to the first event a survey of the surface catchment identified an orange liquor draining from a large mound into a small stream-sink. The pollutant was organic rich material which was being stock-piled prior to spreading on land as an ameliorant and was principally composed of paper pulp and organic rich peat from a water treatment works. This material formed a mound that covered >500 m² to a depth of at least 1m. When the pollution became evident within Peak Cavern the landowner was asked to take action to prevent runoff/pollution entering the cave. However, the second pollution event occurred after this same material had been partially dispersed on the surface catchment. Once the soils were in a saturated state, following heavy rain in late 2001, water re-entered the same sink holes leading to further degradation.

The cave passage downstream of Ink Sump (Figure 1) was heavily stained following both events and the substratum was covered by an orange residue. The staining was observed and reported by recreational cavers and divers but no visible evidence of pollution was detected outside the cave within the springs or river draining the caves (Peakshole Water) during the first event. The second pollution event occurred over a longer period but discolouration of the water was only observed for 24 hours. Hydrological connectivity between the pollutant and the cave was demonstrated by a tracing experiment using two fluorescent dyes, sodium

fluorescein (CI 45359 Acid Yellow 73) and rhodamine WT (CI Acid Red 388). The dyes were detected at the head of Ink Sump, the most upstream visible point where pollution was recorded within Peak Cavern and also entered Far Sump (the second major percolation input to Peak Cavern - Figure 1). The experiment also indicated that a large proportion of the tracer (and therefore the pollutant) travelled ~4 km in an easterly direction and was discharged by a natural spring and two anthropogenic sources (soughs) draining water from disused lead mines, and that a small volume of tracer also entered Speedwell Cavern (Wood et al., 2002). Microfloral analysis of water samples from Peak Cavern indicated the presence of a number of cellulose degrading bacteria associated with the biodegradation of the paper pulp following the second event (Hibberd, 2003).

Monitoring and laboratory processing

The invertebrate community was routinely sampled monthly over the 7-year period (84 months; January 1997- December 2003) from 5 sites within Peak Cavern and from the Peak Cavern Rising (n = 480) and from 6 sites (n = 472) within Speedwell Cavern (Figure 1). Benthic invertebrates were sampled using a 0.05 m² cylinder sampler (fitted with a 90 µm mesh net) over a 30-second period. Additional examination of larger clasts within the cylinder was also undertaken, where they occurred. Due to the potential disturbance and degradation associated with extensive sampling of subterranean habitats single cylinder samples were collected and sampling occasions were used as replicates (Gunn et al., 2000). Sampling could not be undertaken at all sites each month due to flooding of some subterranean passages during the winter and early spring months (5 months within the 84-month study period). At Peak Cavern, three sites were all downstream of the pollution source (Figure 1 – Peak Polluted: PP1, PP2 and PP3) and three sites (control sites) were located on unaffected tributaries (Peak Control: PC1, PC2 and PC3).

All specimens were preserved in the field with 70% industrial methylated spirits (IMS) and returned to the laboratory for processing and identification. Samples were washed and screened on 250µm and 90µm mesh sieves. Material >250µm was manually inspected by removing all invertebrates from an illuminated sorting tray. All sediment retained on the 90µm mesh sieve was examined in a grooved (5 mm) Bogorov sorting tray at 10-50 magnifications to ensure all material from the samples was examined. All macroinvertebrate taxa were identified to species level where possible. Chironomidae, Oligochaeta and Copepoda specimens were examined individually and mounted on microscope slides for examination (up to 400 magnifications) as required for species level identification.

Water temperature (°C), conductivity (µS cm⁻¹), pH and dissolved oxygen (mg l⁻¹) were measured in the field using a portable YSI 600R water quality probe. Replicate water samples were collected from the caves and associated springs and analysed for nitrate (mg l⁻¹) and phosphate (mg l⁻¹) concentrations. Preliminary analysis indicated that there were no significant differences between samples pre- and post-pollution, or between those sites affected by the pollution and those on unaffected tributaries. This reflects the fact that the pollution entered the cave on the flood hydrograph when most of the cave was inaccessible.

Data analysis

Differences in the invertebrate community between the two caves, and sites affected and unaffected by pollution were examined on an annual basis (calendar year January-December). This corresponded to the timing of flood events and the detection of pollution (disturbance events) within Peak Cavern, and provided 2-years of pre-disturbance data (1997 and 1998), 2-years when pollution events occurred (1999 and 2002), and 3 other years (2000, 2001 and 2003). The invertebrate community was characterised by the following metrics: total abundance (individuals m⁻²), number of taxa, Shannon-Wiener diversity index and the Berger

Parker dominance index. The latter two indices were calculated using the α Species Diversity and Richness software (Pisces Conservation, 1998). Preliminary examination of the data for the different sites and years using Levene's test for homogeneity of variances were significant for some groups ($P < 0.05$). Hence, the non-parametric Kruskal-Wallis test was applied to examine differences between the caves, polluted and control sites, and for the different time periods.

Results

Invertebrate community

A total of 34 aquatic invertebrate taxa were recorded during the study period (Table 2). The pre-disturbance Peak Cavern invertebrate community was dominated by Oligochaeta (5 taxa: *Limnodrilus hoffmeisteri*, *Lumbriculus variegatus*, *Spirosperma ferox*, *Stylodrilus* sp. and *Tubifex tubifex*) and Copepoda (4 taxa: *Acanthocyclops venustus*, *A. vernalis*, *Diacyclops bicuspidatus* and *Megacyclops viridis*) in terms of abundance. Other invertebrate taxa typically comprised less than 15% of the total abundance for individual sampling occasions. The community within Speedwell Cavern was more variable and was dominated by Oligochaeta (*Limnodrilus hoffmeisteri* and *Tubifex tubifex*), Chironomidae (particularly two Orthocladiinae: *Rheocricotopus fuscipes* and *Brillia modesta*), Copepoda (*Acanthocyclops venustus*, *A. vernalis*, *Diacyclops bicuspidatus*) and the amphipod *Gammarus pulex*. Faunal abundance displayed seasonal variability, demonstrating the influence of epigeal inputs of water and organic matter. Examination of the invertebrate communities for the pre-disturbance period (1997 and 1998) indicated that samples from Speedwell Cavern supported a greater abundance of invertebrates, although the community was dominated by a smaller number of taxa compared to Peak Cavern (Kruskal-Wallis test : abundance - $P < 0.001$, Berger Parker dominance – $P < 0.005$). Samples from Peak Cavern supported a greater number of taxa and had a higher Shannon-Wiener diversity than Speedwell Cavern (Kruskal-Wallis test:

number of taxa – $P < 0.001$; Shannon-Wiener - $P < 0.001$ – see Wood et al., 2002 for further details).

Pollution episode 1

At sites affected by pollution in Peak Cavern no benthic invertebrates were recorded in the month after the pollution event, although a large number of dead and decaying earthworms (*Lumbricus terrestris*) were recorded at the channel margins. The abundance of freshwater taxa remained low at polluted sites for the rest of the year compared with control sites (Figure 2), with the invertebrate community being almost exclusively composed of two oligochaetes (*Limnodrilus hoffmeisteri* and *Tubifex tubifex*). The first pollution episode resulted in a significant reduction in the abundance at affected sites (1999 in Figure 3a) compared to pre-disturbance data (1997 and 1998 in Figure 3a) and control sites (Figure 3b). A similar pattern was observed for the number of taxa and the Shannon-Wiener diversity index, and an inverse pattern for the Berger-Parker dominance index (see Table 3 for pair-wise comparisons). No significant differences in benthic abundance, number of taxa, Shannon-Wiener diversity or Berger Parker dominance were recorded between the polluted and control sites within Peak Cavern in the following 2 years (2000 and 2001) (Table 3), and there were no differences in any invertebrate community parameters for Speedwell Cavern between the pre-disturbance (1997 and 1998), disturbance (1999) or post-disturbance periods (2000 and 2001).

Pollution episode 2

As a result of the second input of pollutant in 2002, a significant increase in benthic abundance occurred at the affected sites within Peak Cavern (Figure 3a) compared with two of the control sites (Figure 3b). A similar pattern was observed for the Berger-Parker dominance index and an inverse pattern for the number of taxa and the Shannon-Wiener diversity index (see Table 3 for pair-wise comparisons). The abundances of two oligochaetes

(*Limnodrilus hoffmeisteri* and *Tubifex tubifex*) increased significantly (to $>500 \text{ m}^{-2}$) within one month of the input of the pollutant (Kruskal-Wallis test – $P<0.001$). In the following months, numbers of the epigeal amphipod, *Gammarus pulex*, also increased significantly at polluted sites compared with two control sites (Kruskal-Wallis test – $P<0.001$). In February 2002, *G. pulex* was recorded for the first time during the study period at one control site (PC1 in Figure 1). Following the discovery of *G. pulex* at the site the total abundance of invertebrates, particularly Oligochaeta and Copepoda, was reduced compared to the other control sites (Figure 4a-e). No significant differences between any invertebrate community parameters were recorded for Speedwell Cavern following the second pollution episode.

Discussion

The nature of cave pollution and disturbance

Differences in the physical nature of perturbations, in the form of pulse, press and ramp disturbances can result in multiple and markedly different biotic responses within aquatic ecosystems (*sensu* Lake, 2000). Pollution disturbances of groundwater dominated ecosystems can be associated with both press and pulse disturbances. Press disturbances are typically associated with the diffuse entry of material from a relatively large geographical area which percolates into the subterranean groundwater environment (Hancock et al., 2005; Rinaudo et al., 2005). Pulse events are usually associated with the rapid transfer of material into the subterranean environment from a specific location within the surface catchment and may be associated with high water input (Culver et al., 1992; Graening and Brown, 2003). Both of the events recorded in this investigation were clearly point-source disturbances associated with flood events. However, flood events occurring between the two pollution events, during 2000 and 2001, did not appear to result in any significant input of pollutant and acted as ‘flushing flows’ which facilitated the recovery of the benthic invertebrate community (abundance, number of taxa, diversity and dominance) to pre-disturbance levels (Figure 3).

288

289 Both pollution events resulted in significant changes to the benthic invertebrate community of
290 affected sites within Peak Cavern. However, no impact was recorded within the adjacent
291 system (Speedwell Cavern) despite water tracing experiments indicating limited hydrological
292 connectivity with the stream-sink through which the pollutant entered the groundwater system
293 (Wood et al., 2002). This reflects the different hydrological characteristics of the two caves.
294 Water in Speedwell Cavern is primarily derived from sinking streams and as a result the
295 residence time of water within the cave is short, dissolved and particulate organic matter input
296 is relatively high, and pollutants are likely to be diluted and transported through the system
297 relatively quickly (Gunn et al., 2000; Simon et al., 2003). In contrast, water within Peak
298 Cavern is principally derived from percolation water that has passed through the overlying
299 soil and rock and, as a result, the residence time of water is longer. In addition, the volume
300 and delivery of dissolved and particulate organic matter and abundance of invertebrates is
301 usually lower within percolation water dominated systems such as Peak Cavern (Poulson and
302 Lavoie, 2000; Simon et al., 2003). These natural hydrological characteristics reflect a well
303 known gradient of differences that strongly influences the volume, timing and processing rate
304 of trophic resources within subterranean ecosystems (e.g., Poulson and Lavoie, 2000; Simon
305 and Benfield, 2001, Simon et al., 2003).

306

307 *Faunal response to pollution*

308 Faunal response to the pollution events was marked and indicative of significant disturbance
309 events. Direct faunal community response to the pollution of caves has only been recorded in
310 a limited number of previous studies (e.g. Culver et al., 1992), with several studies comparing
311 degraded systems with reference sites in the absence of non-affected control sites (e.g.
312 Holsinger, 1966, Simon and Buikema, 1997). Few studies have included detailed pre- and
313 post-disturbance data or have been undertaken over a comparable length of time. The greatest

changes to the invertebrate community of Peak Cavern were associated with a limited number of taxa (2 oligochaetes: *Limnodrilus hoffmeisteri* and *Tubifex tubifex*, the amphipod *Gammarus pulex* and 4 Copepoda - *Acanthocyclops venustus*, *A. vernalis*, *Diacyclops bicuspidatus* and *Megacyclops viridis*).

During the period immediately following both pollution episodes the invertebrate community at affected sites was dominated by the oligochaetes *Limnodrilus hoffmeisteri* and *Tubifex tubifex*. Both of these taxa are widespread, occur in most surface waters and have been recorded from caves across the globe where they have been associated with organic enrichment (Swayne et al., 2004; Wetzel and Taylor, 2001). During the first pollution event densities were lower than baseline conditions ($<50 \text{ m}^{-2}$) and during the second event they were significantly higher ($>200 \text{ m}^{-2}$) at degraded sites (Figure 3). Their dominance of the invertebrate community within Peak Cavern during these events suggests they are relatively resilient and good indicators of organic pollution within caves and other groundwater dominated ecosystems (Lafont et al., 1996; Lafont and Vivier, 2006).

The increased abundance of cyclopoid copepods following the input of organic material during the second event probably reflects an increased food supply for these taxa. Several cyclopoid copepods (including those in the genus *Acanthocyclops*) are known to be predatory (Fryer, 1957; Galassi et al., 2002), feeding on taxa such as ciliates, rotifers, small oligochaetes and other small crustaceans, all of which may have increased abundances in conditions of high organic matter. At the same time, other cyclopoid taxa are more reliant on fine detrital material (Galassi et al., 2002) that, again, is likely to be more plentiful during an organic pollution event. The abundance of *Gammarus pulex* also increased ($>20 \text{ individuals m}^{-2}$) at affected sites as a result of the second pollution episode, as well as invading one of the adjacent control sites. *G. pulex* have been recorded in many cave systems in the UK, where

they frequently occur in relatively high abundances (Proudlove et al., 2003). Stygophilic gammarids have been recorded within a number of caves around the world where some populations display adaptations to the subterranean environment (e.g. Culver et al., 1995). Epigean *Gammarus* species have been widely reported to be highly competitive and invasive in some instances (MacNeil et al., 2003). It is now widely acknowledged that some gammarids are omnivorous and may be active and effective predators (Kelly et al., 2002) and the invasion of epigean (stygophilic) taxa into subterranean habitats may result in the displacement and/or elimination of hypogean (stygobitic) taxa (Sket, 1977).

The aquatic invertebrate communities of both caves were almost exclusively composed of stygophiles, and none can be regarded as obligate subterranean taxa (stygobites); although the larvae of the dytiscid beetle *Hydroporus ferrugineus* has only been recorded from the Peak-Speedwell system and may be an obligatory subterranean life stage (Alarie et al., 2001). No stygobitic taxa have been recorded from 48 karstic springs within the wider limestone region of the English Peak District (Wood et al., 2005), suggesting that the absence of stygobitic fauna from the Peak-Speedwell Cavern system is not due to pollution alone. Absence of hypogean taxa may reflect glacial activity during the Pleistocene, the maximum extent of which was thought to mark the limits of subterranean faunal distributions. However, there is increasing evidence that stygobitic fauna persisted in sub-glacial refugia beneath the ice in many areas (e.g. Holsinger et al., 1997) including the UK where stygobitic taxa have been recorded some distance north of the maximum extent of glaciation (Proudlove et al., 2003; Bratton, 2006).

Parallels and contrasts between pollution events

A number of parallels and contrasts between the events and their impact on the cave ecosystem can be identified. Both of the pollution events recorded during the study period

coincided with floods and originated from the same location within the surface catchment. The impact on the benthic community at polluted sites was rapid (one month following their detection) and persisted until the next major flood event. However, the response of the community and individual taxa to the two events was markedly different. The first pollution episode resulted in a significant reduction in community abundance, number of taxa and Shannon- diversity index but an increase in the Berger-Parker dominance index at affected sites. The second episodes led to a marked increase in the community abundance and Berger-Parker dominance index, and a reduction in the number of taxa, and Shannon- diversity index. The differences in the community response to the events probably reflects differences in the magnitude of the flood events and associated pollutant loading. The first event resulted in the input of pollutants which were largely contained within the cave and led to the exclusion of almost all fauna from affected sites. The second event was associated with a period of sustained high flow and it is likely that a large proportion of the pollutant was transported through the cave, and was observed as discolouration of the water emerging from Peak Cavern Rising. The pollution load retained within the cave associated with the second event was probably lower, did not lead to sub-lethal concentrations and may have actually enhanced the trophic resources available within the cave leading to the marked increase in the abundance of some members of the invertebrate community (Graening and Brown, 2003; Simon and Buikema, 1997; Sket, 1999).

Recovery of the benthic community was relatively rapid following both pollution events, possibly due to the presence of a large number of refugia within non-polluted sites. Subsequent flooding of the cave in the proceeding years (2000 and 2003 respectively) appeared to “cleanse” the system of the pollutant and facilitated the recovery of fauna at all sites following the first event and all but one site (which was invaded by *Gammarus pulex*) following the second (Figure 3 and Figure 4). In other studies, recovery of aquatic cave

invertebrate communities following pollution disturbances has not been as rapid as reported in the current investigation and the impacts have persisted for some time (in excess of 3-years - see Culver et al., 1992). However, data on recovery times for cave communities are usually absent (Graening and Brown, 2003, Simon and Buikema, 1997), reflecting the long-term and diffuse nature of the impact of pollution on some systems but also the fragility of cave ecosystems, the difficulty of undertaking research in subterranean environments and the paucity of pre-disturbance baseline data available for most systems. In the case of the Peak-Speedwell Cavern system, the relatively rapid recovery may have reflected the legacy of impacts upon the subterranean ecosystem (Gunn et al., 2000). This may also explain the absence of stygobitic taxa which are less competitive and more vulnerable to pollution disturbances than most stygophilic taxa (Graening and Brown, 2003; Panno et al., 2006; Sket, 1999).

Implications for conservation and management

Managing groundwater/subterranean ecosystems is particularly difficult since the most damaging activities usually occur in the surface catchment (van Beynen and Townsend, 2005; Danielopol et al., 2003; Gunn et al., 2000). There may be an extended time-period between a disturbance event occurring in the surface catchment and its detection within the subterranean system, by which time irreversible damage may have already occurred (Hancock et al., 2005). Even after the detection of any pollutant, tracing the source may be problematic because the pollution may have ceased and/or the input may be episodic, as recorded in the current investigation.

In Great Britain (England, Wales, Scotland), the major mechanism for legally protecting, and thereby conserving wildlife and earth science features is through notification as a 'Site of Special Scientific Interest' (SSSI). A list of "operations likely to damage the special interest"

is issued to each owner of land in the boundaries of a SSSI at the time the site is designated and the relevant country authority (Natural England, Countryside Council for Wales, Scottish Natural Heritage) must be consulted before any of the listed operations are undertaken. If it is considered that the proposed action will damage the scientific interest of the site then permission may be denied and the authority may enter into a management agreement with the land owner. Following a Geological Conservation Review (GCR) which began in 1977 (Ellis et al., 1996) 48 ‘cave’ sites were identified and subsequently have been designated as SSSI. Descriptions and evaluations of the geomorphological evolution of each Cave and Karst GCR site have been published (Waltham et al., 1997). At the time of the GCR the boundaries of the 48 sites encompassed 879 named caves, ~30% of the total caves in Britain (Hardwick and Gunn, 1996). These 879 caves included all of the longer cave systems so that ~75% of known cave passage (and hence of the total cave resource) was within areas proposed for conservation.

Some of the caves designated as SSSIs in Great Britain were base on their biological interest, although almost exclusively on the basis of bats (Chiroptera) and/or bat roosts. Aquatic invertebrates are only listed as an additional reason for notification at one site (Pridhamsleigh Cave SSSI, Devon) where *Niphargus glenniei* (Crustacea: Amphipoda), an endemic amphipod which is abundant within the cave, occurs. The Peak-Speedwell Cavern system forms part of a Site of Special Scientific Interest (SSSI) but its designation only covers the earth science interests and does not include any subterranean ecological/biological interests (Gunn et al., 2000). However, designation of a cave SSSI does provide limited, even if unintentional, protection for aquatic cave ecosystems and the communities they support because each SSSI has a list of ‘operations requiring consent’. These have been drawn up to protect the earth science features of interest but by providing controls on water quality and water quantity they may also benefit the whole subterranean ecosystem. Initially the protection of sites was confined to operations on the overlying land surface and the land owner was held responsible for any

infringement. However, in England and Wales, part of the Countryside and Rights of Way Act 2000 (CROW) makes it possible for action to be taken against any person damaging the scientific interest of a SSSI even if the action took place outside the SSSI boundaries. As the current research demonstrates, this is particularly important in the case of active cave systems that often receive inputs of water from surface streams whose catchment is outside of the SSSI. All notified water pollution incidents, whether outside or inside of a SSSI, are subject to investigation by the Environment Agency (England and Wales) or the Scottish Environment Protection Agency. However, if the investigating agency is unaware of the composition, sensitivity or even existence of potentially vulnerable aquatic communities in caves then their conservation is not likely to be considered. Knowledge regarding subterranean biodiversity and its conservation value in the UK is severely limited due to an absence of historic and contemporary scientific research compared to other geographical localities (e.g., Ferreira et al., 2007; Culver et al., 2000) and therefore requires an urgent reassessment.

Many obligate aquatic subterranean organisms (stygobites) are confined to relatively small geographical locations (Christman et al., 2005; Ferreira et al., 2007), and display morphological and physiological adaptations to their environment (Coineau, 2000, Culver et al., 1995). As a result, many aquatic cave communities are scientifically important and of high conservation value (Sket, 1999). Managing and mitigating the effects of organic pollution within groundwater dominated habitats may be particularly difficult due to the highly diffuse nature in which many pollutants enter aquifers and cave ecosystems (Boulton, 2005; Sket, 1999) and the long residence time of water compared to epigean riverine systems. Across North America and in some Europe countries, a greater awareness of subterranean biodiversity exists (Culver et al., 2000; Ferreira et al., 2007; Sket, 1999), and some faunal species have been recognised as threatened by the International Union for Conservation of Nature and Natural Resources (IUCN 2006). However, conservation of subterranean fauna is

problematic since while individual species and caves may be protected, the wider community and the surface catchment usually have limited or no protection.

There is a growing need to consider the importance of groundwater quality within subterranean systems since it has major implications for obligate subterranean taxa, and may ultimately have a significant impact on surface waters and their ecology (Boulton, 2005, Hancock, 2002). However, the identification of indicator organisms and the development of biotic indices for groundwater dominated ecosystems, including caves, are currently limited (e.g. Lafont et al., 1996; van Beynen and Townsend, 2005; Hahn, 2006). Greater awareness regarding the impact and implication of disturbances, particularly pollution, upon groundwater dominated ecosystems is required. Given the limited biological monitoring of subterranean groundwater dependant ecosystems, and the largely unseen consequences of pollution within them, a significant knowledge gap exists regarding their impacts. Future research should address these issues to ensure the continued conservation and protection of subterranean faunal communities and the subterranean and surface water ecosystems within the wider drainage basin.

Acknowledgements

PJW acknowledges the support of the British Ecological Society (Small Ecological Project Grant – 1371) and the Natural Environment Research Council (NER/M/S/1999/00152 and GR8/04287) for funding parts of this research. Thanks to Dr Paul Hardwick, Laura Chapman, Garry Rushworth, Robin Kenyon and Richard Battye for field and laboratory assistance and to John Harrison and Tony Marsden for providing access to the caves. Thanks to Prof. P. Armitage, J. Blackburn, G. Fryer, M. Greenwood and Dr D. Horne for confirmation of faunal identifications. Thanks to Ben Le Bas (Natural England) and David Ottewell (Environment

Agency for comments on cave conservation and management; and to Phil Boon and an anonymous reviewer for comments on a draft of this manuscript.

References

Alarie Y, Wood PJ, DeBruyn AMH, Cuppen JGM. 2001. Description of the larvae of *Hydroporus ferrugineus* Stephens and *H. polaris* Fall (Coleoptera: Adephaga: Dytiscidae). *Aquatic Insects* **23**: 123-133.

van Beynen PV, Townsend K. 2005. A disturbance index for karst environments. *Environmental Management* **36**: 101-116.

Boulton AJ. 2005. Chances and challenges in the conservation of groundwaters and their dependent ecosystems. *Aquatic Conservation: Marine and Freshwater Ecosystems* **15**: 319-323.

Bratton JH. 2006. Occurrence of the well shrimp *Niphargus aquilex* (Crustacea: Niphargidae) in Anglesey, North Wales, UK. *Cave and Karst Science* **33**: 29-30.

Christman MC, Culver DC, Madden MK, White D. 2005. Patterns of endemism of the eastern North American cave fauna. *Journal of Biogeography* **32**: 1441-1452.

Coineau N. 2000. Adaptations to interstitial groundwater life. In *Subterranean Ecosystem: Ecosystems of the World*. 30, Wilkins H, Culver DC, Humphries WF (eds). Elsevier Science: Amsterdam, The Netherlands; 189-210.

Culver DC. 1985. Trophic relationships in aquatic cave environments. *Stygologia* **1**: 43-53.

Culver DC, Jones WK, Holsinger JR. 1992. Biological and hydrological investigation of the Cedars, Lee County, Virginia, an ecologically significant and threatened karst area. In *Proceedings of the 1st International Conference on Ground Water Ecology*, Stanford JA, Simons JJ (eds). American Water Resources Association: Bethesda, MD: 281-290.

Culver DC, Kane TC, Fong DW. 1995. *Adaptation and Natural Selection in Caves: The Evolution of Gammarus minus*. Harvard University Press: Cambridge, Massachusetts.

521 Culver DC, Master LL, Christman, MC and Hobbs HH III 2000. Obligate cave fauna of the
522 48 contiguous United States. *Conservation Biology* **14**: 386-401.

523 Culver DC, Sket B. 2000. Hotspots of subterranean biodiversity in caves and wells. *Journal of*
524 *Cave and Karst Studies* **62**: 11-17.

525 Danielopol DL, Griebler C, Gunatilaka A, Notenboom J. 2003. Present state and future
526 prospects for groundwater ecosystems. *Environmental Conservation* **30**: 104-130.

527 Elliott WR. 2000. Conservation of the North American cave and karst biota. In *Subterranean*
528 *Ecosystem: Ecosystems of the World. 30*, Wilkins H, Culver DC, Humphries WF (ed).
529 Elsevier Science: Amsterdam, The Netherlands; 665-689.

530 Ellis NV, Bowen, DQ, Campbell S, Knill JL, McKirdy AP, Prosser CD, Vincent MA, Wilson
531 RCL. 1996. *An introduction to the Geological Conservation Review*. London, Chapman
532 Hall.

533 Ferreira D, Malard F, Dole-Olivier MJ, Gibert J. 2007. Obligate groundwater fauna of France:
534 diversity patterns and conservation implications. *Biodiversity and Conservation* **16**: 567-
535 597.

536 Finlay JB, Buhay, JE Crandall KA. 2006. Surface to subsurface freshwater connections :
537 phylogeographic and habitat analyses of *Cambarus tenebrosus*, a facultative cave-
538 dwelling crayfish. *Animal Conservation* **9**: 375-387.

539 Fryer G. 1957. The food of some freshwater cyclopoid copepods and its ecological
540 significance. *Journal of Animal Biology* **26**: 263-286.

541 Galassi D, Marmonier P, Dole-Olivier M-J, Rundle SD. 2002. Microcrustacea. In *Freshwater*
542 *Meiofauna: Biology and Ecology*, Rundle SD, Robertson AL, Schmid-Araya JM (eds).
543 Backhuys Publishers: Leiden, The Netherlands; 135-175.

544 Graening GO, Brown AV. 2003. Ecosystem dynamics and pollution effects in an Ozark cave
545 stream. *Journal of the American Water Resources Association* **36**:1497-1507.

546 Green WD, Elliott LP, Crawford NC. 1990. Investigation on nonpoint source pollution
 547 associated with karst aquifer systems. *Transactions of the Kentucky Academy of Science*
 548 **51**:177-181.

549 Gunn J, Tranter J, Perkins J, Hunter C. 1998. Sanitary Bacterial Dynamics in a Mixed Karst
 550 Aquifer. In *Karst Hydrology*, Leibungut, C, Gunn, J. Dassargues A. (eds.). Publication
 551 247, IAHS; 61-70.

552 Gunn J, Hardwick P, Wood PJ. 2000. The invertebrate community of the Peak-Speedwell
 553 cave system, Derbyshire, England – pressures and considerations for conservation
 554 management. *Aquatic Conservation: Marine and Freshwater Ecosystems* **10**: 353-369.

555 Hahn HJ. 2006. The GW-Faunal-Index: A first approach to a quantitative ecological
 556 assessment of groundwater habitats. *Limnologica* **36**: 119-137.

557 Halliday WR. 2003. Raw sewage and solid waste dumps in lava tube caves of Hawaii Island.
 558 *Journal of Cave and Karst Studies* **65**:68-75.

559 Hancock PJ. 2002. Human impacts on the stream-groundwater exchange zone. *Environmental*
 560 *Management* **29**: 763-781.

561 Hancock PJ, Boulton AJ, Humphreys WF. 2005. Aquifers and hyporheic zones: Towards an
 562 ecological understanding of groundwater. *Hydrogeology Journal* **13**: 98-111.

563 Hardwick P, Gunn J. 1996. The conservation of Britain's limestone cave resource.
 564 *Environmental Geology* **28**: 121-127.

565 Hibberd L. 2003. The impact of agricultural paper pulp on the microflora of a Derbyshire
 566 Karst. Unpublished Thesis, University of Huddersfield, UK.

567 Holsinger JR. 1966. A preliminary study of the effects of organic pollution of Banners Corner
 568 Cave, Virginia. *International Journal of Speleology* **2**: 75-89.

569 Holsinger JR. 1988. Troglobites – The evolution of cave dwelling organisms. *American*
 570 *Scientist* **76**: 146-153.

571 Holsinger JR, Carlson KR, Shaw DP. 1997. Biogeographic significance of recently
 572 discovered amphipod crustaceans (*Stygobromus*) in caves of southeastern Alaska and
 573 Vancouver Island. In *Proceedings of the 12th International Congress of Speleology, 10-
 574 17th August 1999, La Chaux-de-Fonds: Switzerland*; 347-349.

575 Hunter C, Perkins J, Tranter J, Gunn J. 1999. Agricultural land-use effects on the indicator
 576 bacterial quality of an upland stream in the Derbyshire Peak District in the UK. *Water
 577 Research* **33**: 3577-3586.

578 IUCN 2006. *2006 IUCN Red List of Threatened Species*. <www.iucnredlist.org>.
 579 Downloaded on 21 August 2007.

580 Jasinska EJ, Knott B, McComb AJ. 1996. Root mats in ground water: a fauna-rich cave
 581 habitat. *Journal of the North American Benthological Society* **15**: 508-519.

582 Kelly DW, Dick JTA, Montgomery WI. 2002. The functional role of *Gammarus* (Crustacea,
 583 Amphipoda): shredders, predators or both. *Hydrobiologia* **485**: 199-203.

584 Lafont M, Camus JC, Rosso A. 1996. Superficial and hypohelic oligochaeta communities as
 585 indicators of pollution and water exchange in the River Moselle, France. *Hydrobiologia*
 586 **334**: 147-155.

587 Lafont M, Vivier A. 2006. Oligochaete assemblages in the hypohelic zone and coarse surface
 588 sediments: Their importance for understanding of ecological functioning of watercourses.
 589 *Hydrobiologia* **546**: 171-181.

590 Lake PS. 2000. Disturbance, patchiness, and diversity in streams. *Journal of the North
 591 American Benthological Society* **19**: 573-592.

592 MacNeil C, Bigsby E, Dick JTA, Hatcher MJ, Dunn AM. 2003. Differential physico-chemical
 593 tolerances and intraguild predation among native and invasive amphipods (Crustacea); a
 594 field study. *Archiv für Hydrobiologie* **156**: 165-179.

595 Panno SV, Hackley KC, Kelly WR, Hwang HH, Wilhelm FM, Taylor SJ, Stiff BJ. 2006.
 596 Potential effects of recurrent low oxygen conditions on the Illinois cave amphipod.
 597 *Journal of Cave and Karst Studies* **68**: 55-63.

598 Pisces Conservation. 1998. α Species Diversity and Richness, Pisces Conservation Ltd.
 599 Lymington, Hampshire, UK.

600 Poulson TL, Lavoie KH. 2000. The trophic basis of subsurface ecosystems. *Subterranean*
 601 *Ecosystem: Ecosystems of the World*, 30,. In Wilkins H, Culver DC, Humphries WF (eds).
 602 Elsevier Science: Amsterdam, The Netherlands; 231-249.

603 Proudlove GS, Wood PJ, Harding PT, Horne DJ, Gledhill T, Knight LRFD. 2003. A review
 604 of the status and distribution of the subterranean aquatic Crustacea of Britain and Ireland.
 605 *Cave and Karst Science* **30**: 51-74.

606 Rinaudo JD, Arnal C, Blanchin R, Elsass R, Meilhac A, Loubier S. 2005. Assessing the cost
 607 of groundwater pollution: the case of diffuse agricultural pollution in the Upper Rhine
 608 valley aquifer. *Water Science and Technology* **52**: 153-162.

609 Simon KS, Benfield EF. 2001. Leaf and wood breakdown in cave streams. *Journal of the*
 610 *North American Benthological Society* **20**: 550-563.

611 Simon KS, Buikema AL. 1997. Effects of organic pollution on an Appalachian cave: changes
 612 in macroinvertebrate population and food supplies. *American Midland Naturalist* **138**:
 613 387-401.

614 Simon KS, Benfield EF, Macko SA. 2003. Food web structure and the role of epilithic
 615 biofilms in cave streams. *Ecology* **84**: 2395-2406.

616 Sket B. 1977. Gegenseitige beeinflussung der wasserpollution und des höhlenmilieus.
 617 *Proceedins of the 6th International Congress of Speleology*. **5**: 253-262.

618 Sket B. 1999. The nature of biodiversity in hypogean waters and how it is endangered.
 619 *Biodiversity and Conservation* **8**: 1319-1338.

620 Swayne H, Day M, Wetzel MJ. 2004. *Limnodrilus hoffmeisteri* (Annelida: Oligochaeta:
 621 Tubficidae) in Pop's Cave, Wisconsin, USA. *Journal of Cave and Karst Studies* **66**: 28-31.
 622 Waltham AC, Simms MJ, Farrant AR, Goldie HS. 1997. *Karst and Caves of Great Britain*.
 623 Chapman and Hall, London 358pp.
 624 Wetzel MJ, Taylor SJ. 2001. First records of freshwater oligochaetes (Annelida, Clitellata)
 625 from caves in Illinois and Missouri, USA. *Journal of Cave and Karst Studies* **63**: 99-104.
 626 Wood PJ, Gunn J, Perkins J. 2002. The impact of pollution on aquatic invertebrates within a
 627 subterranean ecosystem – out of sight out of mind. *Archiv für Hydrobiologie* **155**: 223-237.
 628 Wood PJ, Gunn J, Smith H, Abas-Kutty A. 2005. Flow permanence and macroinvertebrate
 629 community diversity within groundwater dominated headwater streams and springs.
 630 *Hydrobiologia* **545**: 55-64.
 631
 632

List of Figures

Figure 1. The Peak-Speedwell Cavern system indicating the location of invertebrate sampling sites within Speedwell Cavern (1-6), Peak Cavern control sites (PC1-PC3), polluted sites (PP1-PP3) and other specific locations referred to within the text.

Figure 2. Mean invertebrate community abundance (individuals $m^{-2} \pm 1$ SE) within Peak Cavern (January 1999-December 1999) for: (a) control sites and (b) polluted sites.

Figure 3. Mean invertebrate community abundance (individuals m^{-2}) and 95% confidence intervals for the Peak Cavern benthic invertebrate community (January 1997-December 2003) for: (a) polluted sites; and (b) control sites. * Indicates control site (PC1) not included in the series due to invasion of the site by *Gammarus pulex*.

Figure 4. Invertebrate community abundance for unpolluted control sites within Peak Cavern (January 1999-December 1999): (a) mean abundance of all taxa (individuals $m^{-2} \pm 1$ SE) from control site 2 and 3 (PC2 and PC3); (b) abundance (individuals m^{-2}) of all taxa from control site 1 (PC1); (c) mean abundance of dominant Oligochaeta (*Limnodrilus hoffmeisteri* and *Tubifex tubifex* individuals $m^{-2} \pm 1$ SE) from control site 2 and 3 (PC2 and PC3); (d) abundance of dominant Oligochaeta (*Limnodrilus hoffmeisteri* and *Tubifex tubifex* individuals m^{-2}) from control site 1 (PC1); (e) mean abundance of dominant Copepoda (*Acanthocyclops venustus*, *A. vernalis*, *Diacyclops bicuspidatus* and *Megacyclops viridis* individuals $m^{-2} \pm 1$ SE) from control site 2 and 3 (PC2 and PC3); and (f) abundance of dominant Copepoda (*Acanthocyclops venustus*, *A. vernalis*, *Diacyclops bicuspidatus* and *Megacyclops viridis* individuals m^{-2}) from control site 1 (PC1). Solid line indicates the timing of pollution input and dashed line indicates the first record of *Gammarus pulex* at control site 1 (PC1).

Figure 1

Figure 2

Figure 3

Figure 4

658 Table 1. Summary of scientific papers documenting the impact of organic pollution on aquatic invertebrate communities and fauna within
659 cave ecosystems.
660

Author	Location	Pollution	Impact
Culver et al. 1992	Thompson Cedar Cave, Virginia, USA	Sawdust and Bark from Sawmill Operation	Elimination of stygobitic amphipod and isopod populations. An increase in the abundance of epigeal (stygophilic) Oligochaeta (Tubificidae) and Chironomidae larvae. Limited recovery three-years after the event.
Graening and Brown 2003	Cave Springs Cave, Arkansas, USA	Septic leachate, sewage sludge and cow manure suspected	Elimination of stygobitic amphipods although stygobitic isopods flourished.
Holsinger 1966	Banners Corner Cave, Virginia, USA	Septic leachate (sewage)	An increase in the abundance of stygobitic isopod and Planariidae populations at the same time an increase in abundance of epigeal (stygophilic) fauna occurred.
Panno et al. 2006	Illinois' sinkhole plain, Illinois, USA	Septic leachate (sewage)	Elimination of a stygobitic amphipod (<i>Gammarus acherondytes</i>) from one polluted system and recovery in an adjacent system.
Simon and Buikema 1997	Banners Corner Cave, Virginia, USA	Septic leachate (sewage)	Absence of stygobitic isopods from highly polluted pools, but common occurrence in moderately and slightly polluted waters. Exclusion of stygobitic Amphipods from any polluted waters.
Sket 1977	Various cave systems, Dinaric Karst, Slovenia	Organic enrichment Organic enrichment Organic enrichment	Podpeška jama - Increase in abundance of stygobitic fauna in the absence of epigeal (stygophilic) competitors - which had no access to the site. Jama v Šahnu – elimination of all stygobitic fauna and an increase in abundance of a limited number of epigeal (stygophilic) taxa - primarily Oligochaeta (Tubificidae). Postonjna-Planina cave system – Increase in abundance of epigeal (stygophilic) taxa further within the cave and a corresponding decline of stygobitic taxa.
Wood et al., 2002	Peak Cavern, Derbyshire, UK	Paper pulp and peat	Initial exclusion of all taxa and limited recovery of epigeal (stygophilic) taxa 9-months after detection of pollutant.

661 Table 2. Invertebrate fauna recorded from Speedwell Cavern (1997-2003), Peak Cavern
662 (1997-2003), and during the years when pollution occurred (1999 and 2002) within Peak
663 Cavern for affected and control sites.

	SPEEDWELL CAVERN	PEAK CAVERN	PEAK CAVERN (Polluted sites)		PEAK CAVERN (Control sites)	
	1997-2003	1997-2003	1999	2002	1999	2002 ¹
PLANARIIDAE						
<i>Crenobia alpine</i>	X	X			X	X
<i>Phagocata vitta</i>	X					
GASTROPODA						
<i>Lymnaea peregra</i>		X*				
BIVALVIA						
<i>Pisidium nitidum</i>		X			X	X
<i>Pisidium personatum</i>		X			X	X
OLIGOCHAETA						
<i>Aporrectodea rosea</i>	X				X	X
Enchytraeidae	X	X			X	X
<i>Limnodrilus hoffmeisteri</i>	X	X	X	X	X	X
<i>Lumbriculus variegatus</i>	X	X			X	X
<i>Lumbricus terrestris</i>	X	X	X ^a	X	X	X
<i>Spirosperma ferox</i>	X	X		X	X	X
<i>Stylodrilus</i> sp.	X	X				
<i>Tubifex tubifex</i>	X	X	X	X	X	X
CRUSTACEA						
CLADOCERA						
<i>Alona quadrangularis</i>	X	X			X	X
COPEPODA						
HARPACTICOIDA						
<i>Atheyella crassa</i>		X			X	X
<i>Canthocamptus staphylinus</i>	X	X			X	X
CYCLOPOIDA						
<i>Acanthocyclops venustus</i>	X	X		X	X	X
<i>Acanthocyclops vernalis</i>	X	X	X ^b	X	X	X
<i>Diacyclops bicuspidatus bicuspidatus</i>		X	X ^b	X	X	X
<i>Diacyclops bicuspidatus lubbocki</i>		X			X	
<i>Eucyclops agilis</i>	X	X			X	X
<i>Megacyclops gigas</i>	X	X			X	X
<i>Megacyclops viridis</i>	X	X		X	X	X
<i>Paracyclops fimbriatus</i>		X			X	X
GAMMARIDAE						
<i>Gammarus pulex</i>	X	X	X ^b	X	X	X
EPHEMEROPTERA						
<i>Baetis rhodani</i>	X*					
COLEOPTERA						
<i>Hydroporus ferrugineus</i>	X	X			X	
DIPTERA						
Chironomidae						
Chironominae						
<i>Polypedilum</i> sp.	X	X		X	X	X
Orthocladiinae						
<i>Brillia modesta</i>	X	X			X	X
<i>Parametriocnemus stylatus</i>	X	X		X	X	X
<i>Rheocricotopus fuscipes</i>	X	X		X	X	X
Tanypodinae						
<i>Thienemannimyia</i> gp.	X	X				X
Simuliidae	X*					
Thaumaleidae						
<i>Thaumalea verralli</i>		X			X	X

664 Notes: ¹ Includes control site invaded by *Gammarus pulex* in 2002; * Indicates single specimens of
665 stygoxene (accidental) taxa recorded within the subterranean environment; ^a All specimens of
666 *Lumbricus terrestris* recorded in the 5 months following the detection of pollution were dead and/or

667 decomposing; ^b Taxa recorded for the first time 9-months after the detection of the pollution within
668 Peak Cavern.

669

670

671

672

673

674

675

Table 3. Kruskal-Wallis pair-wise comparison between years for invertebrate community parameters at sites within Peak Cavern affected by pollution (January 1997-December 2003): a) abundance (individuals m⁻¹); b) number of taxa; c) Shannon-Wiener diversity index; and d) Berger-Parker dominance index. n.b. Site invaded by *G. pulex* not included in analysis of 2002 and 2003. NS = not significant, * P <0.05, ** P<0.01 and *** P<0.001.

a)

	1997	1998	1999	2000	2001	2002
1998	NS					
1999	**	**				
2000	NS	NS	*			
2001	NS	NS	*	NS		
2002	**	**	***	***	***	
2003	NS	NS	*	NS	NS	***

b)

	1997	1998	1999	2000	2001	2002
1998	NS					
1999	*	*				
2000	NS	NS	*			
2001	NS	NS	*	NS		
2002	**	*	***	**	**	
2003	NS	NS	*	NS	NS	**

c)

	1997	1998	1999	2000	2001	2002
1998	NS					
1999	*	*				
2000	NS	NS	*			
2001	NS	NS	*	NS		
2002	**	*	*	**	**	
2003	NS	NS	*	NS	NS	**

d)

	1997	1998	1999	2000	2001	2002
1998	NS					
1999	*	*				
2000	NS	NS	*			
2001	NS	NS	*	NS		
2002	**	*	*	**	**	
2003	NS	NS	*	NS	NS	**