

Probabilistic Representation and Fall-Off of Bound States of Relativistic Schrödinger Operators with Spin 1/2

Fumio Hiroshima*, Takashi Ichinose† and József Lőrinczi‡

Abstract

A Feynman-Kac type formula of relativistic Schrödinger operators with unbounded vector potential and spin 1/2 is given in terms of a three-component process consisting of Brownian motion, a Poisson process and a subordinator. This formula is obtained for unbounded magnetic fields and magnetic fields with zeros. From this formula an energy comparison inequality is derived. Spatial decay of bound states is established separately for growing and decaying potentials by using martingale methods.

Keywords: relativistic Schrödinger operators, bound states, spatial decay, Feynman-Kac formulae, Poisson process, subordinate Brownian motion, martingales.

*Faculty of Mathematics, Kyushu University, Fukuoka, 819-0395, Japan. hiroshima@math.kyushu-u.ac.jp.

†Department of Mathematics, Kanazawa University, Kanazawa, 920-11, Japan. ichinose@kenroku.kanazawa-u.ac.jp

‡School of Mathematics, Loughborough University, Loughborough LE11 3TU, United Kingdom. J.Lorinczi@lboro.ac.uk

1 Introduction

In the paper [HIL09] we constructed a Feynman-Kac formula for a generalized Schrödinger operator with spin of the form

$$\Psi(h(a, \sigma)) + V. \quad (1.1)$$

Here V is a real-valued external potential, Ψ is an arbitrary Bernstein function with $\Psi(0) = 0$, and h is a Schrödinger-type operator of the form

$$h(a, \sigma) = \frac{1}{2}(\sigma \cdot (p - a))^2, \quad (1.2)$$

including a vector potential $a = (a_1, a_2, a_3)$ describing a magnetic field, and the Pauli matrices $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ describing spin $1/2$. As we have shown, the Feynman-Kac representation of (1.1) involves three independent stochastic processes, Brownian motion, a Poisson process and a subordinator. Moreover, spin $1/2$ was also extended to higher spins in [HIL09], see also [ARS91].

In this paper we consider a functional integral representation of the strongly continuous one-parameter semigroup generated by the relativistic Schrödinger operator with spin $1/2$ in three-dimensional space,

$$\sqrt{(\sigma \cdot (p - a))^2 + m^2} - m + V. \quad (1.3)$$

Here m is the mass of the relativistic particle, which we regard as a parameter; see [Car78] for standard Schrödinger operators, where $a = 0$. This Hamilton operator is a special case of (1.1) obtained by choosing

$$\Psi(u) = \sqrt{2u + m^2} - m, \quad m \geq 0. \quad (1.4)$$

In this case we have the $\frac{1}{2}$ -stable subordinator about which more details are known than about subordinators related to a general Ψ . Using this extra information, our main goal in this paper is to prove a Feynman-Kac-type formula for (1.3) under weaker conditions than needed for general Ψ , and use it to derive the fall-off properties of bound states. In particular, in contrast to [HIL09] we can cover unbounded magnetic fields in Theorem 3.6 and magnetic fields with zeros in Theorem 3.8.

This paper is organized as follows. Section 2 is devoted to introducing the relativistic Schrödinger operator with spin $1/2$ as a self-adjoint operator on $\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$ and a unitary equivalent representation on $L^2(\mathbb{R}^3 \times \mathbb{Z}_2)$. In Section 3.1 we reassess results in [HIL09] and give a Feynman-Kac formula with bounded magnetic fields. In Section 3.2 we prove a Feynman-Kac formula for unbounded magnetic fields, and in Section 3.3 for magnetic fields having zeros. In Section 4 we derive the decay properties of bound states separately for growing and decaying potentials by using martingale methods.

2 Relativistic Schrödinger operator with spin 1/2

2.1 Definitions

We begin by defining the self-adjoint operator $h(a, \sigma)$ and $\sqrt{2h(a, \sigma) + m^2} - m + V$ rigorously.

The spinless Schrödinger operator h_0 with vector potential a and zero external potential is defined as a self-adjoint operator on $L^2(\mathbb{R}^3)$. Let $D_\mu = p_\mu - a_\mu$, where $p_\mu = -i\partial_{x_\mu}$ is the generalized differential operator. Define the quadratic form q by

$$H^1(\mathbb{R}^3) \times H^1(\mathbb{R}^3) \ni (f, g) \mapsto q(f, g) = \frac{1}{2} \sum_{\mu=1}^3 (D_\mu f, D_\mu g), \quad (2.1)$$

where $H^1(\mathbb{R}^3) = \{f \in L^2(\mathbb{R}^3) \mid D_\mu f \in L^2(\mathbb{R}^3), \mu = 1, 2, 3\}$. If $a \in (L^2_{\text{loc}}(\mathbb{R}^3))^3$, then the quadratic form q is non-negative and closed, and hence there exists a unique self-adjoint operator h_0 satisfying $(h_0 f, g) = q(f, g)$, for $f \in D(h_0)$ and $g \in H^1$, where $D(h_0) = \{f \in Q(q) \mid q(f, \cdot) \in L^2(\mathbb{R}^3)'\}$. Let $C_0^\infty(\mathbb{R}^3) = C_0^\infty$ be the set of infinitely many times differentiable functions with compact support on \mathbb{R}^3 . It can be seen that C_0^∞ is a form core for h_0 under the assumption $a \in (L^2_{\text{loc}}(\mathbb{R}^3))^3$, see [LS81].

Next we introduce a magnetic field $b = (b_1, b_2, b_3)$. Physically it is given by $b = \nabla \times a$, however, in this paper we regard the magnetic field b independent of the vector potential a . We will use the following conditions on the vector potential a .

Assumption 2.1 (Vector potential) The vector potential $a = (a_1, a_2, a_3)$ is a vector-valued function whose components a_μ , $\mu = 1, 2, 3$, are real-valued functions such that $a \in (L^2_{\text{loc}}(\mathbb{R}^3))^3$ and $\nabla \cdot a \in L^1_{\text{loc}}(\mathbb{R}^3)$, where $\nabla \cdot a$ is understood in distributional sense.

Assumption 2.2 (Magnetic field) Suppose that $D(-\Delta) \subset D(b_\mu)$ and for $f \in D(-\Delta)$ the conditions $\|b_\mu f\| \leq \kappa_\mu \|-\Delta f\| + \kappa'_\mu \|f\|$, $\mu = 1, 2, 3$, and $\kappa_1 + \kappa_2 + \kappa_3 < 1$ are satisfied.

Finally we introduce the spin variables. Let $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ be the 2×2 Pauli matrices given by

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

They satisfy the relations $\sigma_\mu \sigma_\nu + \sigma_\nu \sigma_\mu = 2\delta_{\mu\nu} 1$ and $\sigma_\mu \sigma_\nu = i \sum_{\lambda=1}^3 \varepsilon^{\lambda\mu\nu} \sigma_\lambda$, where $\varepsilon^{\lambda\mu\nu}$ is the anti-symmetric Levi-Civita tensor with $\varepsilon^{123} = 1$. Then it can be seen directly that

$$\sigma \otimes b = \sum_{\mu=1}^3 \sigma_\mu \otimes b_\mu = \begin{bmatrix} b_3 & b_1 - ib_2 \\ b_1 + b_2 & -b_3 \end{bmatrix}.$$

Under Assumption 2.2 $\sigma \otimes b$ is relatively bounded with respect to $1 \otimes 2h_0$, as an operator in $\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$, with a relative bound strictly smaller than 1,

$$\|(\sigma \otimes b)f\| \leq (\kappa_1 + \kappa_2 + \kappa_3)\|1 \otimes 2h_0 f\| + C\|f\|, \quad f \in \mathbb{C}^2 \otimes D(h). \quad (2.2)$$

This follows through the diamagnetic inequality $|(f, e^{-th_0}g)| \leq (|f|, e^{-t(-\frac{1}{2}\Delta)}|g|)$ under Assumption 2.1. Thus the self-adjoint operator

$$h = 1 \otimes h_0 - \frac{1}{2}\sigma \otimes b \quad (2.3)$$

in $\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$ is bounded from below under Assumptions 2.1 and 2.2. We choose m so as to guarantee that

$$2h + m^2 = 1 \otimes 2h_0 - \sigma \otimes b + m^2 \geq 0.$$

Note that under a suitable condition h is positive, and in this case we can take $m = 0$. From now on we omit the tensor product \otimes for notational convenience.

We now define the self-adjoint operator H .

Definition 2.3 Under Assumptions 2.1 and 2.2, H is defined by the self-adjoint operator

$$H = \sqrt{2h + m^2} - m \quad (2.4)$$

in $\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$. Here the square root is taken through the spectral resolution of $2h + m^2$.

An example is the operator $\sqrt{(\sigma \cdot (p - a))^2 + m^2} - m$ such that $a \in (L_{\text{loc}}^4(\mathbb{R}^3))^3$, $\nabla \cdot a \in L_{\text{loc}}^2(\mathbb{R}^3)$ and $\nabla \times a \in (L_{\text{loc}}^2(\mathbb{R}^3))^3$. In this case it is seen that

$$(\sigma \cdot (p - a))^2 = (p - a)^2 + \sigma \cdot (\nabla \times a)$$

on $1 \otimes C_0^\infty(\mathbb{R}^3)$.

2.2 Spin variable

In order to construct a functional integral representation of $(f, e^{-t(H+V)}g)$ we make a unitary transform of H on $\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$ to an operator on the space $L^2(\mathbb{R}^3 \times \mathbb{Z}_2)$. This is a space of L^2 -functions of $x \in \mathbb{R}^3$ and an additional two-valued spin variable $\theta \in \mathbb{Z}_2$, where

$$\mathbb{Z}_2 = \{-1, 1\}. \quad (2.5)$$

We define the spin interaction U on $L^2(\mathbb{R}^3 \times \mathbb{Z}_2)$ by

$$U : f(x, \theta) \mapsto U_d(x, \theta)f(x, \theta) + U_{\text{od}}(x, -\theta)f(x, -\theta) \quad (2.6)$$

where $(x, \theta) \in \mathbb{R}^3 \times \mathbb{Z}_2$,

$$U_d(x, \theta) = -\frac{1}{2}\theta b_3(x) \quad (2.7)$$

is the diagonal component, and

$$U_{\text{od}}(x, -\theta) = -\frac{1}{2}(b_1(x) - i\theta b_2(x)) \quad (2.8)$$

is the off-diagonal component. Let

$$h_{\mathbb{Z}_2} = h_0 + U. \quad (2.9)$$

Under Assumption 2.2 U is symmetric, relatively bounded with respect to h_0 with a relative bound strictly smaller than 1 so that $h_{\mathbb{Z}_2}$ and h are unitary equivalent,

$$h_{\mathbb{Z}_2} \cong h \quad (2.10)$$

as seen below. Define the unitary operator $F : L^2(\mathbb{R}^3 \times \mathbb{Z}_2) \rightarrow \mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$ by

$$F : f \mapsto \begin{bmatrix} f(\cdot, +1) \\ f(\cdot, -1) \end{bmatrix}. \quad (2.11)$$

Also, define $\tau_\mu = F^{-1}\sigma_\mu F$. We see that $\tau_1 : f(x, \theta) \mapsto f(x, -\theta)$, $\tau_2 : f(x, \theta) \mapsto -i\theta f(x, -\theta)$ and $\tau_3 : f(x, \theta) \mapsto \theta f(x, \theta)$.

Definition 2.4 Let Assumptions 2.1 and 2.2 hold. Then $H_{\mathbb{Z}_2}$ is defined by

$$H_{\mathbb{Z}_2} = \sqrt{2h_{\mathbb{Z}_2} + m^2} - m. \quad (2.12)$$

In what follows instead of H we study $H_{\mathbb{Z}_2}$, and write H (resp. h) instead of $H_{\mathbb{Z}_2}$ (resp. $h_{\mathbb{Z}_2}$).

2.3 Three independent stochastic processes

In order to construct a path integral representation we will need three independent stochastic processes $(B_t)_{t \geq 0}$, $(N_t)_{t \geq 0}$ and $(T_t)_{t \geq 0}$ which we introduce next. We denote the expectation with respect to path measure W starting at x by \mathbb{E}_W^x .

Let $(B_t)_{t \geq 0}$ be three-dimensional Brownian motion on a probability space $(\Omega_P, \mathcal{F}_P, P^x)$ with initial point $P^x(B_0 = x) = 1$.

Secondly, let $(N_t)_{t \geq 0}$ be a Poisson process on a probability space $(\Omega_N, \mathcal{F}_N, \mu)$ with unit intensity, i.e.,

$$\mu(N_t = n) = \frac{t^n}{n!} e^{-t}, \quad n \in \mathbb{N} \cup \{0\}.$$

Let μ^α be the image measure of the process $(N_t + \alpha)_{t \geq 0}$ for $\alpha \in \mathbb{Z}_2$ and thus $\mathbb{E}_\mu^\alpha[f(N_\cdot)] = \mathbb{E}_\mu^0[f(N_\cdot + \alpha)]$. We define integrals with respect to this process in terms of the sum of evaluations at jumping times, i.e., for g we define

$$\int_{(a,b]} g(s, N_s) dN_s = \sum_{\substack{r \in (a,b] \\ N_r+ \neq N_{r-}}} g(r, N_r) \quad (2.13)$$

and we also write $\int_a^{b+} \cdots dN_s$ for $\int_{(a,b]} \cdots dN_s$. Associated with the Poisson process we also define a \mathbb{Z}_2 -valued stochastic process $(\theta_t)_{t \geq 0}$ on $(\Omega_N, \mathcal{F}_N, \mu^0)$ by

$$\theta_t = (-1)^{N_t}. \quad (2.14)$$

Finally, let $(T_t)_{t \geq 0}$ denote the subordinator starting from 0 at $t = 0$ on a given probability space $(\Omega_\nu, \mathcal{F}_\nu, \nu)$ defined by its Laplace transform

$$\mathbb{E}_\nu^0[e^{-T_t u}] = \exp\left(-t\left(\sqrt{2u + m^2} - m\right)\right). \quad (2.15)$$

Note that $(T_t)_{t \geq 0}$ is a one-dimensional Lévy process with right continuous paths with left limits, almost surely non-decreasing. It can be more explicitly described as the first hitting time process

$$T_t = \inf\{s > 0 \mid B_s^1 + ms = t\},$$

where $(B_t^1)_{t \geq 0}$ is a one-dimensional Brownian motion independent of the three-dimensional Brownian motion B_t above. We also define the measure ν^s , $s \in \mathbb{R}$, by the image measure on $(T_t + s)_{t \geq 0}$, and use the shorthand

$$\mathbb{E}_P^x \mathbb{E}_\mu^\alpha \mathbb{E}_\nu^s = \mathbb{E}_M^{x, \alpha, s}. \quad (2.16)$$

The role of these three stochastic processes is as follows. Clearly, the Schrödinger operator $-\frac{1}{2}\Delta + V$ generates an Itô process which can be described using the Brownian motion $(B_t)_{t \geq 0}$ under V . The Poisson process $(N_t)_{t \geq 0}$ results from the Schrödinger operator with spin. Finally, the subordinator $(T_t)_{t \geq 0}$ appears due to the relativistic Schrödinger operator which generates a Lévy process. A particular combination of these three independent stochastic processes then yields the path integral representation of $e^{-t(H+V)}$ which we will discuss below.

2.4 Generator of Markov process

Consider the $\mathbb{R}^3 \times \mathbb{Z}_2$ -valued joint Brownian and jump process

$$\Omega_P \times \Omega_N \ni (\omega, \omega_1) \mapsto X_t(\omega, \omega_1) = (B_t(\omega), \theta_t(\omega_1)) \in \mathbb{R}^3 \times \mathbb{Z}_2$$

with initial value X_0 . The generator of this Markov process is [HIL09]

$$G_0 = -\frac{1}{2}\Delta + \sigma_F + 1, \quad (2.17)$$

where 1 is the 2×2 identity matrix and σ_F is the fermionic harmonic oscillator defined in terms of the Pauli matrices by

$$\sigma_F = \frac{1}{2}(\sigma_3 + i\sigma_2)(\sigma_3 - i\sigma_2) - 1 = -\sigma_1.$$

Note that $\inf \text{Spec}(G_0) = 0$.

In the relativistic case, the subordinator explained above appears in addition to this. We define the subordinate process $(q_t)_{t \geq 0}$ in terms of the $\mathbb{R}^3 \times \mathbb{Z}_2$ -valued stochastic process

$$\Omega_P \times \Omega_N \times \Omega_\nu \ni (\omega, \omega_1, \omega_2) \mapsto q_t(\omega, \omega_1, \omega_2) = (B_{T_t(\omega_2)}(\omega), \theta_{T_t(\omega_2)}(\omega_1)) \in \mathbb{R}^3 \times \mathbb{Z}_2.$$

In a similar manner to $(X_t)_{t \geq 0}$, we can identify the generator of $(q_t)_{t \geq 0}$.

Proposition 2.5 *The generator of the Markov process $(q_t)_{t \geq 0}$ is*

$$G = \sqrt{-\Delta + 2\sigma_F + 2 + m^2} - m \quad (2.18)$$

and its characteristic function is given by

$$\mathbb{E}_M^{0,0,0}[e^{iZq_t}] = \mathbb{E}_M^{0,0,0}[e^{i\xi B_{T_t}} e^{iz\theta_{T_t}}] = e^{-t(\sqrt{|\xi|^2 + m^2} - m)} \cos z + i e^{-t(\sqrt{|\xi|^2 + 4 + m^2} - m)} \sin z \quad (2.19)$$

for $Z = (\xi, z) \in \mathbb{R}^3 \times \mathbb{R}$.

PROOF. This is obtained through the equalities

$$\begin{aligned} \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} [\overline{f(q_0)} g(q_t)] &= \mathbb{E}_\nu^0 \left[\sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_{P \times \mu}^{x,\alpha} [\overline{f(q_0)} g(q_t)] \right] \\ &= \mathbb{E}_\nu^0 [(f, e^{-T_t(-\frac{1}{2}\Delta + \sigma_F + 1)} g)] = (f, e^{-tG} g). \end{aligned}$$

Hence it follows that (2.18) is the generator of $(q_t)_{t \geq 0}$, while (2.19) is straightforward.

qed

3 Feynman-Kac-type representations

3.1 Bounded magnetic field

In this subsection we briefly discuss some results established in [HIL09] obtained for a general version of the relativistic Schrödinger operator with spin and bounded magnetic field. Write

$$W(x) = \frac{1}{2} \sqrt{b_1(x)^2 + b_2(x)^2}, \quad (3.1)$$

and notice that $|U_{\text{od}}(x, \theta)| = W(x)$.

Proposition 3.1 (Feynman-Kac formula: bounded magnetic field) *Let Assumption 2.1 hold and assume that $b_\mu \in L^\infty$ for $\mu = 1, 2, 3$. Let V be relatively bounded with*

respect to $\sqrt{-\Delta + m^2}$ with a relative bound strictly smaller than 1. Assume, furthermore, that

$$\mathbb{E}_{P \times \nu}^{x,0} \left[\int_0^{T_t} |\log W(B_s)| ds \right] < \infty, \quad \text{a.e. } x \in \mathbb{R}^3. \quad (3.2)$$

Then $H + V$ is self-adjoint on $D(H)$ and

$$(f, e^{-t(H+V)}g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{S}} \right], \quad (3.3)$$

where the exponent $\mathcal{S} = \mathcal{S}_V + \mathcal{S}_A + \mathcal{S}_S$ is given by

$$\mathcal{S}_V = - \int_0^t V(B_{T_s}) ds, \quad (3.4)$$

$$\mathcal{S}_A = -i \int_0^{T_t} a(B_s) \circ dB_s, \quad (3.5)$$

$$\mathcal{S}_S = - \int_0^{T_t} U_d(B_s, \theta_s) ds + \int_0^{T_t+} \log(-U_{od}(B_s, -\theta_{s-})) dN_s. \quad (3.6)$$

PROOF. Since $\|Vf\| \leq \kappa \|\sqrt{-\Delta + m^2}f\| + \kappa' \|f\|$ with constants $\kappa < 1$ and κ' , and b_μ is bounded, we have $\|Vf\| \leq \kappa \|Hf\| + C \|f\|$ with a constant C . Hence self-adjointness follows by the Kato-Rellich theorem. (3.3) follows from [HIL09, Theorem 5.9]. **qed**

We note that \mathcal{S}_A and \mathcal{S}_S in Proposition 3.1 stand for the integrals $-i \int_0^r a(B_s) \circ dB_s$ and $-\int_0^r U_d(B_s, \theta_s) ds + \int_0^{r+} \log(-U_{od}(B_s, -\theta_{s-})) dN_s$ evaluated at $r = T_t$, respectively.

A Feynman-Kac formula without spin is an immediate corollary. This was first established in [CMS90] without a vector potential; we give a version including a vector potential. Let

$$H_{\text{spinless}} = \sqrt{2h_0 + m^2} - m. \quad (3.7)$$

Corollary 3.2 *Let Assumption 2.1 hold, and assume that $V = V_+ - V_-$ satisfies that $V_+ \in L_{\text{loc}}^1(\mathbb{R}^3)$ and V_- is relatively form bounded with respect to $\sqrt{-\Delta + m^2}$ with a relative bound strictly less than 1. Then*

$$(f, e^{-t(H_{\text{spinless}} + V_+ - V_-)}g) = \int_{\mathbb{R}^3} dx \mathbb{E}_{P \times \nu}^{x,0} \left[\overline{f(B_0)} g(B_{T_t}) e^{\mathcal{S}_V + \mathcal{S}_A} \right]. \quad (3.8)$$

In particular, when $a = 0$,

$$(f, e^{-t(\sqrt{-\Delta + m^2} - m + V_+ - V_-)}g) = \int_{\mathbb{R}^3} dx \mathbb{E}_{P \times \nu}^{x,0} \left[\overline{f(X_0)} g(X_t) e^{-\int_0^t V(X_s) ds} \right]. \quad (3.9)$$

By Corollary 3.2 we have the following energy comparison inequality. Let

$$H_0 = \sqrt{-\Delta + m^2} - m. \quad (3.10)$$

Corollary 3.3 *Under the assumptions of Corollary 3.2 we have*

- (1) $|(f, e^{-t(H_{\text{spinless}} + V_+ - V_-)}g)| \leq (|f|, e^{-t(H_0 + V_+ - V_-)}|g|)$
- (2) $\inf \text{Spec}(H_0 + V_+ - V_-) \leq \inf \text{Spec}(H_{\text{spinless}} + V_+ - V_-).$

3.2 Unbounded magnetic field

We extend the Feynman-Kac formula above (Proposition 3.1) to the case of magnetic fields b that are possibly unbounded and satisfy Assumption 2.2. This extension is not straightforward, and we need several lemmas.

Define the truncated magnetic field $b^{(N)}$ by

$$b_\mu^{(N)}(x) = \begin{cases} b_\mu(x) & \text{if } |b_\mu(x)| \leq N \\ N & \text{if } b_\mu(x) > N \\ -N & \text{if } b_\mu(x) < -N. \end{cases}$$

Then the Feynman-Kac formula for the Hamiltonian with the truncated magnetic field is readily given by Proposition 3.1 in which b is replaced by $b^{(N)}$. Let H_N be defined by H with b replaced by $b^{(N)}$.

Lemma 3.4 *Under Assumptions 2.1 and 2.2 the semigroup e^{-tH_N} is strongly convergent to e^{-tH} as $N \rightarrow \infty$.*

PROOF. Let h_N be h with b replaced by $b^{(N)}$. We see that $h_N \rightarrow h$ as $N \rightarrow \infty$ on the common domain $D(h_N) = D(h)$. Then $e^{-th_N} \rightarrow e^{-th}$ strongly as $N \rightarrow \infty$. Thus it is immediate to see that

$$(f, e^{-tH_N}g) = \mathbb{E}_\nu^0[(f, e^{-T_t h_N}g)] \rightarrow \mathbb{E}_\nu^0[(f, e^{-T_t h}g)] = (f, e^{-tH}g), \quad (3.11)$$

which implies strong convergence. qed

Lemma 3.5 *Let $f, g \in L^2(\mathbb{R}^3 \times \mathbb{Z}_2)$, and set*

$$\rho = f(q_0)g(q_t)e^{\int_0^{T_t} \frac{1}{2}|b_3(B_s)|ds}e^{\int_0^{T_t} \log W(B_s)dN_s}e^{T_t}.$$

Then under Assumption 2.2 it follows that $\sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0}[|\rho|] < \infty$.

PROOF. Define the spin operator $|U|$ and $|U|_N$ by

$$|U| : f(x, \theta) \mapsto -\frac{1}{2}|b_3(x)|f(x, \theta) - W(x)f(x, -\theta), \quad (3.12)$$

$$|U|_N : f(x, \theta) \mapsto -\frac{1}{2}|b_3^{(N)}(x)|f(x, \theta) - W^{(N)}(x)f(x, -\theta), \quad (3.13)$$

where $W^{(N)}$ is W with b replaced by $b^{(N)}$, and define

$$\widehat{H} = \sqrt{-\Delta + 2|U| + m^2} - m. \quad (3.14)$$

Also, we define \widehat{H}_N by \widehat{H} with $|U|$ replaced by $|U|_N$. Let $f, g \in L^2(\mathbb{R}^3)$ be non-negative. For \widehat{H}_N we have the Feynman-Kac formula

$$(f, e^{-t\widehat{H}_N} g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t \overline{f(q_0)} g(q_t)} e^{\widehat{\mathcal{S}}_S^N} \right], \quad (3.15)$$

where

$$\widehat{\mathcal{S}}_S^N = \int_0^{T_t} \frac{1}{2} |b_3^{(N)}(B_s)| ds + \int_0^{T_t+} \log W^{(N)}(B_s) dN_s. \quad (3.16)$$

By the monotone convergence theorem for forms we see that $e^{-t(-\Delta+2|U|_N)} \rightarrow e^{-t(-\Delta+2|U|)}$ strongly as $N \rightarrow \infty$, and thus $e^{-t\widehat{H}_N} \rightarrow e^{-t\widehat{H}}$ strongly as $N \rightarrow \infty$ is shown in the same way as (3.11). Then the monotone convergence theorem for integrals implies that ρ is integrable and the Feynman-Kac formula (3.15) with $b^{(N)}$ replaced by b also holds. **qed**

Now we can state the first main theorem.

Theorem 3.6 (Feynman-Kac formula: unbounded magnetic field) *Let Assumptions 2.1 and 2.2 as well as condition (3.2) hold, and suppose that V is relatively bounded with respect to $\sqrt{-\Delta + m^2}$ with a relative bound strictly less than 1. Then $H + V$ is self-adjoint on $D(H)$ and*

$$(f, e^{-t(H+V)} g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t \overline{f(q_0)} g(q_t)} e^{\mathcal{S}} \right]. \quad (3.17)$$

PROOF. We divide the proof in five steps.

Step 1: Suppose that $V = 0$. Then the theorem holds.

Proof: Recall that H_N is defined by H with b replaced by $b^{(N)}$. Then the Feynman-Kac formula holds with \mathcal{S}_S replaced by \mathcal{S}_S^N , where \mathcal{S}_S^N is defined by \mathcal{S}_S with b replaced by $b^{(N)}$:

$$(f, e^{-tH_N} g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t \overline{f(q_0)} g(q_t)} e^{\mathcal{S}_S^N + \mathcal{S}_A} \right]. \quad (3.18)$$

The left hand side above converges to $(f, e^{-tH}g)$ as $N \rightarrow \infty$ by Lemma 3.4. On the other hand, we have

$$e^{T_t} |f(q_0)g(q_t)| |e^{\mathcal{I}_S^N + \mathcal{I}_A}| \leq e^{T_t} |f(q_0)g(q_t)| e^{\int_0^{T_t} \frac{1}{2} |b_3(B_s)| ds} e^{\int_0^{T_t} \log W(B_s) dN_s}$$

so that the right hand side of (3.18) is integrable by Lemma 3.5, and therefore the Lebesgue dominated convergence theorem yields

$$\lim_{N \rightarrow \infty} \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} f(q_0)g(q_t) e^{\mathcal{I}_S^N + \mathcal{I}_A} \right] = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} f(q_0)g(q_t) e^{\mathcal{I}_S + \mathcal{I}_A} \right].$$

Hence the theorem follows for $V = 0$.

Step 2: V is relatively bounded with respect to H with a relative bound strictly smaller than 1. In particular, $H + V$ is self-adjoint on $D(H)$.

Proof: Let $b_0 = (\sqrt{b_1^2 + b_2^2}, 0, b_3)$ and H_{b_0} be defined by H with $a = 0$ and b replaced by b_0 , i.e., $H_{b_0} = \sqrt{-\Delta + \sigma \cdot b_0 + m^2} - m$. Set $\sigma \cdot b_0 = U_{b_0}$. Then we have

$$\|\sqrt{-\Delta + m^2}f\|^2 = \|(H_{b_0} + m)f\|^2 + (f, -U_{b_0}f).$$

Since $|(f, U_{b_0}f)| \leq \kappa' \|f\|^2$ with a constant κ' , and $\|Vf\| \leq \kappa \|\sqrt{-\Delta + m^2}f\| + \kappa'' \|f\|$ with constants $\kappa < 1$ and κ'' , we have $\|Vf\| \leq A \|H_{b_0}f\| + C \|f\|$ with some C and $A < 1$. From the Feynman-Kac formula established in Step 1 the diamagnetic inequality,

$$|(f, e^{-tH}g)| \leq (|f|, e^{-tH_{b_0}}|g|) \quad (3.19)$$

follows. From (3.19) we have $\|H_{b_0}f\| \leq \|Hf\| + c\|f\|$, and thus

$$\|Vf\| \leq A \|Hf\| + C' \|f\|$$

with a constant C' . Hence self-adjointness follows by the Kato-Rellich theorem.

Step 3: Suppose $V \in L^\infty(\mathbb{R}^3) \cap C(\mathbb{R}^3)$. Then the statement holds.

Proof: By the Trotter product formula and the Markov property of $(q_t)_{t \geq 0}$ we have that

$$\begin{aligned} (f, e^{-t(H+V)}g) &= \lim_{n \rightarrow \infty} (f, (e^{-(t/n)H} e^{-(t/n)V})^n g) \\ &= \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} f(q_0)g(q_t) e^{-\sum_{j=1}^n (t/n)V(B_{T_{tj/n}})} e^{\mathcal{I}_S + \mathcal{I}_A} \right]. \end{aligned}$$

Note that $s \mapsto V(B_{T_s})$ is continuous in $s \in [0, t]$ except for at most finitely many points. Thus

$$-\sum_{j=1}^n (t/n)V(B_{T_{tj/n}}(\omega)) \xrightarrow{n \rightarrow \infty} -\int_0^t V(B_{T_s}(\omega))ds$$

for almost every $(\omega, \omega_2) \in \Omega_P \times \Omega_\nu$ as a Riemann integral. Then the theorem follows for $V \in L^\infty(\mathbb{R}^3) \cap C(\mathbb{R}^3)$.

Step 4: Suppose $V \in L^\infty(\mathbb{R}^3)$. Then the statement holds.

Proof: Let $V_n = \phi(\cdot/n)(V * j_n)$, where $j_n(x) = n^3 \phi(xn)$ with $\phi \in C_0^\infty$ such that $0 \leq \phi \leq 1$, $\int_{\mathbb{R}^3} \phi(x) dx = 1$ and $\phi(0) = 1$. Then $V_n(x) \rightarrow V(x)$ for $x \notin \mathcal{N}$, where \mathcal{N} is a set of Lebesgue measure zero. Notice that

$$\mathbb{E}_{P \times \nu}^{x,0}[1_{\mathcal{N}}(B_{T_s})] = \int_{\mathbb{R}^3} 1_{\mathcal{N}}(x+y) P_s(y) dy = 0$$

for $x \in \mathcal{N}$, where

$$P_s(x) = 2 \left(\frac{m}{2\pi} \right)^2 \frac{s}{s^2 + |x|^2} K_2 \left(m \sqrt{|x|^2 + s^2} \right)$$

is the distribution of the random variable B_{T_s} and

$$K_2(x) = \frac{1}{2} \int_0^\infty \xi e^{-\frac{1}{2}(\xi + \xi^{-1})x} d\xi$$

is the modified Bessel function of the third kind. Hence

$$0 = \int_0^t \mathbb{E}_{P \times \nu}^{x,0}[1_{\mathcal{N}}(B_{T_s})] ds = \mathbb{E}_{P \times \nu}^{x,0} \left[\int_0^t 1_{\mathcal{N}}(B_{T_s}) ds \right].$$

Then the Lebesgue measure of $\{s \in [0, \infty) \mid B_{T_s(\omega_2)}(\omega) \in \mathcal{N}\}$ is zero for almost every path $(\omega, \omega_2) \in \Omega_P \times \Omega_\nu$. Therefore $\int_0^t V_n(B_{T_s}) ds \rightarrow \int_0^t V(B_{T_s}) ds$ as $n \rightarrow \infty$ for almost every path $(\omega, \omega_2) \in \Omega_P \times \Omega_\nu$. Moreover,

$$\begin{aligned} & \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{I}_A + \mathcal{I}_S} e^{-\int_0^t V_n(B_s) ds} \right] \\ & \xrightarrow{n \rightarrow \infty} \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{I}_A + \mathcal{I}_S} e^{-\int_0^t V(B_s) ds} \right]. \end{aligned}$$

On the other hand, $e^{-t(H+V_n)} \rightarrow e^{-t(H+V)}$ strongly as $n \rightarrow \infty$, since $H + V_n$ converges to $H + V$ on the common domain $D(H)$. Then the theorem follows for $V \in L^\infty(\mathbb{R}^3)$.

Step 5: We complete the proof of Theorem 3.6. Let $V = V_+ - V_-$ and $V_{mn} = V_{+m} - V_{-n}$, with V_+ , V_- denoting the positive and negative parts of V , respectively, and $V_{+m}(x) = V_+(x)$ if $V_+(x) \leq m$, and $V_{+m}(x) = m$ if $V_+(x) \geq m$, similarly $V_{-n}(x) = V_-(x)$ if $V_-(x) \leq n$ and $V_{-n}(x) = n$ if $V_-(x) \geq n$. Then by the monotone convergence theorem for forms, we

have $e^{-t(H+V_{mn})}$ strongly converges to $e^{-t(H+V_{m\infty})}$ as $n \rightarrow \infty$, and furthermore $e^{-t(H+V_{m\infty})}$ strongly converges to $e^{-t(H+V)}$ as $m \rightarrow \infty$. Hence

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} e^{-t(H+V_{mn})} = e^{-t(H+V)}.$$

On the other hand, by the monotone convergence theorem for integrals the right hand side converges. This completes the proof of the theorem. **qed**

3.3 Magnetic field with zeros

Next we consider the case when the off-diagonal component $U_{\text{od}}(x, -\theta)$ vanishes for some $x \in \mathbb{R}^3$. In this case it is not clear whether $\int_0^{t+} |\log W(B_s)| dN_s < \infty$ holds almost surely. An example when this is not the case is obtained by choosing $b \in (C_0^\infty)^3$.

Let

$$\delta_\varepsilon(z) = \begin{cases} 1, & |z| < \varepsilon, \\ 0, & |z| \geq \varepsilon, \end{cases}$$

for $z \in \mathbb{C}$ and write

$$\chi_\varepsilon(z) = z + \varepsilon \delta_\varepsilon(z), \quad z \in \mathbb{C}. \quad (3.20)$$

We see that

$$|\chi_\varepsilon(U_{\text{od}}(x, -\theta))| > \varepsilon, \quad (x, \theta) \in \mathbb{R}^3 \times \mathbb{Z}_2.$$

Define h_ε by h with the off-diagonal part replaced by $\chi_\varepsilon(U_{\text{od}}(x, -\theta))$, i.e.,

$$h_\varepsilon f(x, \theta) = (h_0 + U_{\text{d}}(x, \theta)) f(x, \theta) + \chi_\varepsilon(U_{\text{od}}(x, -\theta)) f(x, -\theta), \quad (x, \theta) \in \mathbb{R}^3 \times \mathbb{Z}_2.$$

Also, define H_ε by H with U_{od} replaced by $\chi_\varepsilon(U_{\text{od}}(x, -\theta))$.

We note that for every $(x, \omega, \omega_1, \omega_2) \in \mathbb{R}^3 \times \Omega_P \times \Omega_N \times \Omega_\nu$ there exists a number $n = n(\omega_1, \omega_2)$ and random jump times $r_1(\omega_1), \dots, r_n(\omega_1)$ of $s \mapsto N_s$ for $0 \leq s \leq T_t(\omega_2)$ such that

$$\int_0^{T_t(\omega_2)+} \log W(x + B_s(\omega)) dN_s = \sum_{j=1}^{n(\omega_1, \omega_2)} \log W(x + B_{r_j(\omega_1)}(\omega)).$$

Consider

$$\mathscr{W} = \left\{ (x, \omega, \omega_1, \omega_2) \in \mathbb{R}^3 \times \Omega_P \times \Omega_N \times \Omega_\nu \left| \int_0^{T_t+} \log W(x + B_s) dN_s > -\infty \right. \right\}. \quad (3.21)$$

Notice that by the definition $(x, \omega, \omega_1, \omega_2) \in \mathscr{W}^c$ if and only if there exists r such that

$$(1) \quad 0 < r \leq t \leq T_t(\omega_2),$$

(2) $s \mapsto N_s$ is discontinuous at $s = r$,

(3) $b_1(B_r(\omega)) = b_2(B_r(\omega)) = 0$.

Lemma 3.7 *For every $(x, \omega, \omega_1, \omega_2) \in \mathcal{W}^c$ we have*

$$\lim_{\varepsilon \rightarrow 0} \left| e^{\int_0^{T_t^+} \log(-\chi_\varepsilon(U_{\text{od}}(B_s, -\theta_{s-}))) dN_s} \right| = 0.$$

PROOF. We have $|e^{\int_0^{T_t^+} \log(-\chi_\varepsilon(U_{\text{od}}(B_s, -\theta_{s-}))) dN_s}| \leq e^{\int_0^{T_t^+} \log(W(B_s) + \varepsilon) dN_s}$. Observe that

$$\int_0^{T_t^+} \log(W(B_s) + \varepsilon) dN_s = \sum_{j=1}^n \log(W(B_{r_j}) + \varepsilon), \quad r_1, \dots, r_n \in (0, T_t].$$

Since $(x, \omega, \omega_1, \omega_2) \in \mathcal{W}^c$, there exists an r_i such that $b_1(B_{r_i}(\omega)) = b_2(B_{r_i}(\omega)) = 0$. Then

$$\int_0^{T_t^+} \log(W(B_s) + \varepsilon) dN_s = \sum_{j \neq i}^n \log(W(B_{r_j}) + \varepsilon) + \log \varepsilon,$$

and $e^{\int_0^{T_t^+} \log(W(B_s) + \varepsilon) dN_s} \leq e^{\sum_{j \neq i}^n \log(W(B_{r_j}) + \varepsilon)} e^{\log \varepsilon}$. Thus $\lim_{\varepsilon \rightarrow 0} |e^{\int_0^{T_t^+} \log(W(B_s) + \varepsilon) dN_s}| = 0$, and the lemma follows. qed

Theorem 3.8 (Feynman-Kac formula: magnetic field with zeros) *Let Assumptions 2.1 and 2.2 hold, and suppose that V is relatively bounded with respect to $\sqrt{-\Delta} + m^2$ with a relative bound strictly less than 1. Let \mathcal{W} be given by (3.21). Then*

$$(f, e^{-t(H+V)}g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{S}} 1_{\mathcal{W}} \right]. \quad (3.22)$$

PROOF. Put $V = 0$ and fix $\varepsilon > 0$. We can show that the functional integral representation of H_ε is given by (3.17) with \mathcal{S} replaced by $\mathcal{S}_A + \mathcal{S}_S(\varepsilon)$ with

$$\mathcal{S}_S(\varepsilon) = - \int_0^{T_t} U_d(B_s, \theta_s) ds + \int_0^{T_t^+} \log(-\chi_\varepsilon(U_{\text{od}}(B_s, -\theta_{s-}))) dN_s. \quad (3.23)$$

That is,

$$(f, e^{-tH_\varepsilon}g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x,\alpha,0} \left[e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{S}_A + \mathcal{S}_S(\varepsilon)} \right]. \quad (3.24)$$

Take the limit $\varepsilon \downarrow 0$ on both sides above. This gives

$$\lim_{\varepsilon \downarrow 0} \exp(-tH_\varepsilon) = \exp(-tH) \quad (3.25)$$

in strong sense, obtained in the same way as Lemma 3.4. On the other hand, by the Lebesgue dominated convergence theorem it follows that

$$\lim_{\varepsilon \downarrow 0} \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x, \alpha, 0} \left[e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{S}_A + \mathcal{S}_S(\varepsilon)} \right] = \int_{\mathbb{R}^3} dx \mathbb{E}_M^{x, \alpha, 0} \left[\lim_{\varepsilon \downarrow 0} e^{T_t} \overline{f(q_0)} g(q_t) e^{\mathcal{S}_A + \mathcal{S}_S(\varepsilon)} \right].$$

By Lemma 3.7 we find that $\lim_{\varepsilon \rightarrow 0} \mathcal{S}_S(\varepsilon) = 0$ on \mathcal{W} and hence

$$\lim_{\varepsilon \downarrow 0} e^{\mathcal{S}_A + \mathcal{S}_S(\varepsilon)} = \lim_{\varepsilon \downarrow 0} e^{\mathcal{S}_A + \mathcal{S}_S(\varepsilon)} 1_{\mathcal{W}} + \lim_{\varepsilon \downarrow 0} e^{\mathcal{S}_A + \mathcal{S}_S(\varepsilon)} 1_{\mathcal{W}^c} = e^{\mathcal{S}_A + \mathcal{S}_S} 1_{\mathcal{W}}.$$

Next suppose that $V \in L^\infty(\mathbb{R}^3) \cap C(\mathbb{R}^3)$. In this case we can show the theorem in the same way as in Step 3 in the proof of Theorem 3.6. Furthermore, the theorem holds for the required V in the same way as in Steps 4 and 5 above. **qed**

A diamagnetic inequality follows immediately from Theorem 3.8. Recall that H_{b_0} is defined by H with b replaced by $b_0 = (\sqrt{b_1^2 + b_2^2}, 0, b_3)$ and a by zero, respectively.

Corollary 3.9 (Energy comparison inequality) *Under the assumptions of Theorem 3.8 we have that*

$$|(f, e^{-t(H+V)} g)| \leq (|f|, e^{-t(H_{b_0}+V)} |g|). \quad (3.26)$$

In particular, it follows that $\inf \text{Spec}(H_{b_0} + V) \leq \inf \text{Spec}(H + V)$.

4 Fall-off of bound states

4.1 Martingale properties: non-relativistic case

In this section we prove the decay properties of bound states of relativistic Schrödinger operators with spin by means of the Feynman-Kac formula derived in the previous section. For simplicity we assume throughout that

$$\mathbb{E}_P^x \left[\int_0^t |\log W(B_s)| ds \right] < \infty, \quad \text{a.e. } x \in \mathbb{R}^3, \quad (4.1)$$

and

$$\mathbb{E}_{P \times \nu}^{x, 0} \left[\int_0^{T_t} |\log W(B_s)| ds \right] < \infty, \quad \text{a.e. } x \in \mathbb{R}^3, \quad (4.2)$$

i.e., the measure of \mathcal{W}^c in (3.21) is zero.

We first consider the non-relativistic case. Let H_{NR} be the Hamiltonian defined by

$$H_{\text{NR}} = h + V, \quad (4.3)$$

where h is given by (2.3). Let \mathcal{S}_{NR} be defined by the exponent \mathcal{S} with the subordinator T_t replaced by the non-random time t . If Assumptions 2.1 and 2.2 hold and V is relatively

bounded with respect to $-\Delta$ with a relative bound strictly smaller than 1, then $h + V$ is self-adjoint on $D(-\Delta)$. Then the Feynman-Kac formula of $(f, e^{-t(h+V)}g)$ is

$$(f, e^{-t(h+V)}g) = \sum_{\alpha=0,1} \int_{\mathbb{R}^3} dx \mathbb{E}_{P \times \mu}^{x,\alpha} \left[e^{\overline{t f(B_0, \theta_0)}} g(B_t, \theta_t) e^{\mathcal{K}_{\text{NR}}} \right], \quad (4.4)$$

where the exponent $\mathcal{K}_{\text{NR}} = \mathcal{K}_{\text{NRV}} + \mathcal{K}_{\text{NRA}} + \mathcal{K}_{\text{NRS}}$ is given by

$$\begin{aligned} \mathcal{K}_{\text{NRV}} &= - \int_0^t V(B_s) ds, \\ \mathcal{K}_{\text{NRA}} &= -i \int_0^t a(B_s) \circ dB_s, \\ \mathcal{K}_{\text{NRS}} &= - \int_0^t U_d(B_s, \theta_s) ds + \int_0^{t+} \log(-U_{\text{od}}(B_s, -\theta_{s-})) dN_s. \end{aligned}$$

Let φ_g be a bound state such that $H_{\text{NR}}\varphi_g = E\varphi_g$ with $E \in \mathbb{R}$. We consider the spatial decay of $|\varphi_g(x, (-1)^\alpha)|$, i.e., its behavior for large $|x|$.

Let $\mathcal{K}_{\text{NR}}(x, \alpha) = \mathcal{K}_{\text{NRV}}(x) + \mathcal{K}_{\text{NRA}}(x) + \mathcal{K}_{\text{NRS}}(x, \alpha)$ be given by \mathcal{K}_{NR} with B_s and N_s replaced by $B_s + x$ and $N_s + \alpha$, respectively:

$$\begin{aligned} \mathcal{K}_{\text{NRV}}(x) &= - \int_0^t V(B_s + x) ds, \\ \mathcal{K}_{\text{NRA}}(x) &= -i \int_0^t a(B_s + x) \circ dB_s, \\ \mathcal{K}_{\text{NRS}}(x, \alpha) &= - \int_0^t U_d(B_s + x, (-1)^\alpha \theta_s) ds + \int_0^{t+} \log(-U_{\text{od}}(B_s + x, -(-1)^\alpha \theta_{s-})) dN_s. \end{aligned}$$

Define the stochastic process $(M_t(x, \alpha))_{t \geq 0}$ by

$$M_t(x, \alpha) = e^{t(E+1)} e^{\mathcal{K}_{\text{NR}}(x, \alpha)} \varphi_g(B_t + x, (-1)^\alpha \theta_t), \quad t \geq 0,$$

and the filtration

$$\mathcal{M}_t = \sigma((B_r, \theta_r), 0 \leq r \leq t), \quad t \geq 0.$$

Note that $e^{-t(H_{\text{NR}} - E)}\varphi_g = \varphi_g$ and then

$$\mathbb{E}_{P \times \mu}^{x,\alpha}[M_t(0, 0)] = \mathbb{E}_{P \times \mu}^{0,0}[M_t(x, \alpha)] = \varphi_g(x, (-1)^\alpha) \quad (4.5)$$

by (4.4).

Lemma 4.1 *The stochastic process $(M_t(x, \alpha))_{t \geq 0}$ is a martingale with respect to $(\mathcal{M}_t)_{t \geq 0}$, i.e., $\mathbb{E}_{P \times \mu}^{0,0}[M_t(x, \alpha) | \mathcal{M}_s] = M_s(x, \alpha)$ for $t \geq s$.*

PROOF. We prove the case when $(x, \alpha) = (0, 0)$ for notational simplicity, the proof for $(x, \alpha) \neq (0, 0)$ is similar. Let $\mathcal{K}_{\text{NR}}([u, v])$ be defined by \mathcal{K}_{NR} with the integration domain in $\int_0^t \cdots$ replaced by $\int_u^v \cdots$. Write $M_t = M_t(0, 0)$. We see that

$$\mathbb{E}_{P \times \mu}^{0,0}[M_t | \mathcal{M}_s] = e^{t(E+1)} e^{\mathcal{K}_{\text{NR}}([0,s])} \mathbb{E}_{P \times \mu}^{0,0} [e^{\mathcal{K}_{\text{NR}}([s,t])} \varphi_g(B_t, \theta_t) | \mathcal{M}_s].$$

By the Markov property of the $\mathbb{R}^3 \times \mathbb{Z}_2$ -valued stochastic process $(B_t, N_t)_{t \geq 0}$ we have

$$\begin{aligned} & \mathbb{E}_{P \times \mu}^{0,0} [e^{\mathcal{K}_{\text{NR}}([s,t])} \varphi_g(B_t, \theta_t) | \mathcal{M}_s] \\ &= \mathbb{E}_{P \times \mu}^{B_s, N_s} \left[e^{-\int_0^{t-s} V(B_r) dr} e^{-i \int_0^{t-s} a(B_r) \circ d B_r} e^{\int_0^{t-s} U_d(B_r, \theta_r) dr} e^K \varphi_g(B_{t-s}, \theta_{t-s}) \right]. \end{aligned} \quad (4.6)$$

The off-diagonal part K in (4.6) is

$$\begin{aligned} K &= \sum_{\substack{s < u \leq t \\ N_{(u-s)+} \neq N_{(u-s)-}}} \log(-U_{\text{od}}(B_{u-s}, -\theta_{(u-s)-})) \\ &= \sum_{\substack{0 < r \leq t-s \\ N_r+ \neq N_r-}} \log(-U_{\text{od}}(B_r, -\theta_{r-})) = \int_0^{(t-s)+} \log(-U_{\text{od}}(B_r, -\theta_{r-})) dN_r. \end{aligned}$$

Hence we conclude that

$$\mathbb{E}_{P \times \mu}^{0,0} [e^{\mathcal{K}_{\text{NR}}([s,t])} \varphi_g(B_t, \theta_t) | \mathcal{M}_s] = \mathbb{E}_{P \times \mu}^{B_s, N_s} [e^{\mathcal{K}_{\text{NR}}([0,t-s])} \varphi_g(B_{t-s}, \theta_{t-s})],$$

which implies that

$$\mathbb{E}_{P \times \mu}^{0,0} [M_t | \mathcal{M}_s] = e^{s(E+1)} e^{\mathcal{K}_{\text{NR}}([0,s])} \mathbb{E}_{P \times \mu}^{B_s, N_s} [M_{t-s}] = M_s.$$

Thus the lemma follows. qed

4.2 Martingale properties: relativistic case

Next we discuss the relativistic case $H + V$. Let φ_g be a bound state of $H + V$ such that

$$(H + V)\varphi_g = E\varphi_g \quad (4.7)$$

for $E \in \mathbb{R}$. We use the same notation φ_g as for the non-relativistic case. Consider the stochastic process

$$Y_t = e^{tE} e^{T_t} e^{\mathcal{J}} \varphi_g(q_t), \quad t \geq 0. \quad (4.8)$$

Furthermore, we define

$$Y_t(x, \alpha) = e^{tE} e^{T_t} e^{\mathcal{J}(x, \alpha)} \varphi_g(q_t(x, \alpha)), \quad t \geq 0, \quad (4.9)$$

where $q_t(x, \alpha) = (B_{T_t} + x, (-1)^\alpha \theta_{T_t})$ and $\mathcal{S}(x, \alpha) = \mathcal{S}_V(x) + \mathcal{S}_A(x) + \mathcal{S}_S(x, \alpha)$ is given by

$$\mathcal{S}_V = - \int_0^t V(B_{T_s} + x) ds, \quad (4.10)$$

$$\mathcal{S}_A = -i \int_0^{T_t} a(B_s + x) \circ dB_s, \quad (4.11)$$

$$\mathcal{S}_S = - \int_0^{T_t} U_d(B_s + x, (-1)^\alpha \theta_s) ds + \int_0^{T_t+} \log(-U_{od}(B_s + x, -(-1)^\alpha \theta_{s-})) dN_s. \quad (4.12)$$

Then

$$\mathbb{E}_M^{x, \alpha, 0}[Y_t] = \mathbb{E}_M^{0, 0, 0}[Y_t(x, \alpha)] = \varphi_g(x, (-1)^\alpha). \quad (4.13)$$

We introduce a filtration under which $(Y_t)_{t \geq 0}$ is a martingale. Define $Y_t(\omega)$ and $Y_t(x, \alpha, \omega)$ for every $\omega \in \Omega_\nu$ by Y_t and $Y_t(x, \alpha)$, respectively, and with subordinator T_t replaced by the number $T_t(\omega) \geq 0$. Let

$$\mathcal{F}_t^{(1)}(\omega) = \sigma((B_r, N_r), 0 \leq r \leq T_t(\omega)) \in \mathcal{F}_P \times \mathcal{F}_\mu \quad (4.14)$$

for $\omega \in \Omega_\nu$, and define

$$\mathcal{F}_t^{(1)} = \left\{ \bigcup_{\omega \in \Omega_\nu} (A(\omega), \omega) \mid A(\omega) \in \mathcal{F}_t^{(1)}(\omega) \right\} \subset \mathcal{F}_P \times \mathcal{F}_\mu \times \mathcal{F}_\nu. \quad (4.15)$$

We also define

$$\mathcal{F}_t^{(2)} = \left\{ \bigcup_{\omega \in \Omega_P \times \Omega_N} (\omega, B(\omega)) \mid B(\omega) \in \sigma(T_r, 0 \leq r \leq t) \right\} \subset \mathcal{F}_P \times \mathcal{F}_\mu \times \mathcal{F}_\nu. \quad (4.16)$$

We see that $\mathcal{F}_t^{(1)}$ and $\mathcal{F}_t^{(2)}$ are sub- σ -fields of $\mathcal{F}_P \times \mathcal{F}_\mu \times \mathcal{F}_\nu$. Write

$$\mathcal{F}_t = \mathcal{F}_t^{(1)} \cap \mathcal{F}_t^{(2)}, \quad t \geq 0. \quad (4.17)$$

The conditional expectation $\mathbb{E}_M^{0, 0, 0}[Y_t(x, \alpha) | \mathcal{F}_t^{(1)}] = \mathbb{E}_M^{0, 0, 0}[Y_t(x, \alpha) | \mathcal{F}_t^{(1)}](\cdot, \cdot, \cdot)$ is a stochastic process on $\Omega_P \times \Omega_N \times \Omega_\nu$.

Lemma 4.2 *We have $\mathbb{E}_M^{0, 0, 0}[Y_t(x, \alpha) | \mathcal{F}_t^{(1)}](\cdot, \cdot, \omega) = \mathbb{E}_{P \times \mu}^{0, 0}[Y_t(x, \alpha, \omega) | \mathcal{F}_t^{(1)}(\omega)](\cdot, \cdot)$ for all $\omega \in \Omega_\nu$.*

PROOF. Let $A = \bigcup_{\omega \in \Omega_\nu} (A(\omega), \omega)$ with $A(\omega) \in \mathcal{F}_t^{(1)}(\omega)$. Then

$$\begin{aligned} \mathbb{E}_M^{0, 0, 0}[1_A Y_t(x, \alpha)] &= \int_{\Omega_\nu} d\nu(\omega) \mathbb{E}_{P \times \mu}^{0, 0}[1_{A(\omega)} Y_t(x, \alpha, \omega)] \\ &= \int_{\Omega_\nu} d\nu(\omega) \mathbb{E}_{P \times \mu}^{0, 0} \left[1_{A(\omega)}(\cdot, \cdot) \mathbb{E}_{P \times \mu}^{0, 0} [Y_t(x, \alpha, \omega) | \mathcal{F}_t^{(1)}(\omega)](\cdot, \cdot) \right]. \end{aligned}$$

On the other hand, we have

$$\begin{aligned}\mathbb{E}_M^{0,0,0}[1_A Y_t(x, \alpha)] &= \mathbb{E}_M^{0,0,0} \left[1_A \mathbb{E}_M^{0,0,0}[Y_t(x, \alpha) | \mathcal{F}_t^{(1)}] \right] \\ &= \int_{\Omega_\nu} d\nu(\omega) \mathbb{E}_{P \times \mu}^{0,0} \left[1_{A(\omega)}(\cdot, \cdot) \mathbb{E}_M^{0,0,0} \left[Y_t(x, \alpha) | \mathcal{F}_t^{(1)} \right] (\cdot, \cdot, \omega) \right].\end{aligned}$$

A comparison of the two sides above completes the proof. qed

Lemma 4.3 *The stochastic process $(Y_t(x, \alpha))_{t \geq 0}$ is a martingale with respect to $(\mathcal{F}_t)_{t \geq 0}$, i.e., $\mathbb{E}_M^{0,0,0}[Y_t(x, \alpha) | \mathcal{F}_s] = Y_s(x, \alpha)$ for $t \geq s$.*

PROOF. We prove the case when $(x, \alpha) = (0, 0)$ to keep the notation simple, the proof for $(x, \alpha) \neq (0, 0)$ is again similar.

Note that $\mathbb{E}_M^{0,0,0}[Y_t | \mathcal{F}_s] = \mathbb{E}_M^{0,0,0}[Y_t | \mathcal{F}_s^{(1)} \cap \mathcal{F}_s^{(2)}] = \mathbb{E}_M^{0,0,0}[\mathbb{E}_M^{0,0,0}[Y_t | \mathcal{F}_s^{(1)}] | \mathcal{F}_s^{(2)}]$. We first compute $\mathbb{E}_{P \times \mu}^{0,0}[Y_t(\omega) | \mathcal{F}_s^{(1)}(\omega)]$. Write

$$\begin{aligned}\mathcal{S}([u, v]) &= - \int_u^v V(B_{T_r}) dr - i \int_{T_u}^{T_v} a(B_r) \circ dB_r \\ &\quad - \int_{T_u}^{T_v} U_d(B_r, \theta_r) dr + \int_{T_u}^{T_v+} \log(-U_{od}(B_r, -\theta_{r-})) dN_r\end{aligned}$$

and, for every $\omega \in \Omega_\nu$

$$\begin{aligned}\mathcal{S}([u, v], \omega) &= - \int_u^v V(B_{T_r(\omega)}) dr - i \int_{T_u(\omega)}^{T_v(\omega)} a(B_r) \circ dB_r \\ &\quad - \int_{T_u(\omega)}^{T_v(\omega)} U_d(B_r, \theta_r) dr + \int_{T_u(\omega)}^{T_v(\omega)+} \log(-U_{od}(B_r, -\theta_{r-})) dN_r\end{aligned}$$

and $q_t(\omega) = (B_{T_t(\omega)}, \theta_{T_t(\omega)})$, $t \geq 0$. Since $T_t(\omega)$ is non-random, we see in a similar way to the non-relativistic case that

$$\begin{aligned}\mathbb{E}_{P \times \mu}^{0,0}[Y_t(\omega) | \mathcal{F}_s^{(1)}(\omega)] &= e^{tE} e^{T_t(\omega)} e^{\mathcal{S}([0, s], \omega)} \mathbb{E}_{P \times \mu}^{0,0}[e^{\mathcal{S}([s, t], \omega)} \varphi_g(q_t(\omega)) | \mathcal{F}_s^{(1)}(\omega)] \\ &= e^{tE} e^{T_t(\omega)} e^{\mathcal{S}([0, s], \omega)} \mathbb{E}_{P \times \mu}^{B_{T_s(\omega)}, N_{T_s(\omega)}} \left[e^{-\int_s^t V(B_{T_r(\omega)-T_s(\omega)}) dr} e^{-i \int_{T_s(\omega)}^{T_t(\omega)} a(B_{r-T_s(\omega)}) \circ dB_r} \right. \\ &\quad \times e^{-\int_{T_s(\omega)}^{T_t(\omega)} U_d(B_{r-T_s(\omega)}, \theta_{r-T_s(\omega)}) dr} e^{\int_{T_s(\omega)}^{T_t(\omega)+} \log(-U_{od}(B_{r-T_s(\omega)}, -\theta_{(r-T_s(\omega))}) dN_r} \\ &\quad \left. \times \varphi_g(B_{T_t(\omega)-T_s(\omega)}, \theta_{T_t(\omega)-T_s(\omega)}) \right].\end{aligned}$$

Hence by Lemma 4.2 we have

$$\mathbb{E}_M^{0,0,0}[Y_t|\mathcal{F}_s^{(1)}] = e^{tE} e^{T_s} e^{\mathcal{S}([0,s])} Z_{t,s},$$

where

$$\begin{aligned} Z_{t,s} = & e^{T_t-T_s} \mathbb{E}_{P \times \mu}^{B_{T_s}, N_{T_s}} \left[e^{-\int_s^t V(B_{T_r-T_s}) dr} e^{-i \int_{T_s}^{T_t} a(B_{r-T_s}) \odot dB_r} \right. \\ & \left. \times e^{-\int_{T_s}^{T_t} U_d(B_{r-T_s}, \theta_{r-T_s}) dr} e^{\int_{T_s}^{T_t+} \log(-U_{\text{od}}(B_{r-T_s}, -\theta_{(r-T_s)-})) dN_r} \varphi_g(B_{T_t-T_s}, \theta_{T_t-T_s}) \right]. \end{aligned}$$

Here $Z_{t,s}$ is given by

$$\begin{aligned} & e^{u-v} \mathbb{E}_{P \times \mu}^{B_u, N_v} \left[e^{-\int_s^t V(B_{T_r-u}) dr} e^{-i \int_v^u a(B_{r-v}) \odot dB_r} \right. \\ & \left. \times e^{-\int_v^u U_d(B_{r-v}, \theta_{r-v}) dr} e^{\int_v^{u+} \log(-U_{\text{od}}(B_{r-v}, -\theta_{(r-v)-})) dN_r} \varphi_g(B_{u-v}, \theta_{u-v}) \right] \end{aligned}$$

evaluated at $u = T_t$ and $v = T_s$. Take the conditional expectation of the right hand side above with respect to $\mathcal{F}_s^{(2)}$. We note that

$$\mathbb{E}_M^{0,0,0}[f|\mathcal{F}_s^{(2)}](\omega_1, \omega_2, \cdot) = \mathbb{E}_\nu^0[f(\omega_1, \omega_2, \cdot)|\mathcal{N}_s](\cdot), \quad (4.18)$$

where $\mathcal{N}_s = \sigma(T_r, 0 \leq r \leq s)$. Since $e^{tE} e^{T_s} e^{\mathcal{S}([0,s])}$ is measurable with respect to $\mathcal{F}_s^{(2)}$, by (4.18) we consider the conditional expectation of $Z_{t,s}$ giving

$$\begin{aligned} & \mathbb{E}_M^{0,0,0}[Z_{t,s}|\mathcal{F}_s^{(2)}] \\ & = \mathbb{E}_\nu^0 \left[e^{T_t-T_s} \mathbb{E}_{P \times \mu}^{B_{T_s}, N_{T_s}} \left[e^{-\int_s^t V(B_{T_r-T_s}) dr} e^{-i \int_0^{T_t-T_s} a(B_r) \odot dB_r} \right. \right. \\ & \quad \left. \left. \times e^{-\int_0^{T_t-T_s} U_d(B_r, \theta_r) dr} e^{\int_0^{(T_t-T_s)+} \log(-U_{\text{od}}(B_r, -\theta_{r-})) dN_r} \varphi_g(B_{T_t-T_s}, \theta_{T_t-T_s}) \right] \middle| \mathcal{N}_s \right], \end{aligned}$$

where we used the Markov property of $((B_t, N_t))_{t \geq 0}$. By the Markov property of $(T_t)_{t \geq 0}$ we have

$$\begin{aligned} & = \mathbb{E}_\nu^{T_s} \left[e^{T_t-s-T_0} \mathbb{E}_{P \times \mu}^{B_{T_0}, N_{T_0}} \left[e^{-\int_s^t V(B_{T_r-s-T_0}) dr} e^{-i \int_0^{T_t-s-T_0} a(B_r) \odot dB_r} \right. \right. \\ & \quad \left. \left. \times e^{-\int_0^{T_t-s-T_0} U_d(B_r, \theta_r) dr} e^{\int_0^{(T_t-s-T_0)+} \log(-U_{\text{od}}(B_r, -\theta_{r-})) dN_r} \varphi_g(B_{T_t-s-T_0}, \theta_{T_t-s-T_0}) \right] \right]. \end{aligned}$$

Since $\mathbb{E}_\nu^u[f(T)] = \mathbb{E}_\nu^0[f(T+u)]$, we see that

$$\begin{aligned} & = \mathbb{E}_\nu^0 \left[e^{T_t-s-T_0} \mathbb{E}_{P \times \mu}^{B_{T_0+u}, N_{T_0+u}} \left[e^{-\int_s^t V(B_{T_r-s-T_0}) dr} e^{-i \int_0^{T_t-s-T_0} a(B_r) \odot dB_r} \right. \right. \\ & \quad \left. \left. \times e^{-\int_0^{T_t-s-T_0} U_d(B_r, \theta_r) dr} e^{\int_0^{(T_t-s-T_0)+} \log(-U_{\text{od}}(B_r, -\theta_{r-})) dN_r} \varphi_g(B_{T_t-s-T_0}, \theta_{T_t-s-T_0}) \right] \right] \Big|_{u=T_s} \\ & = \mathbb{E}_M^{B_{T_s}, N_{T_s}, 0} \left[e^{T_t-s} e^{-\int_0^{t-s} V(B_{T_r}) dr} e^{-i \int_0^{T_t-s} a(B_r) \odot dB_r} \right. \\ & \quad \left. \times e^{-\int_0^{T_t-s} U_d(B_r, \theta_r) dr} e^{\int_0^{T_t-s+} \log(-U_{\text{od}}(B_r, -\theta_{r-})) dN_r} \varphi_g(q_{t-s}) \right] \\ & = (e^{-(t-s)H_{\mathbb{Z}_2}} \varphi_g)(q_s). \end{aligned}$$

Hence we conclude that

$$\mathbb{E}_M^{0,0,0}[Y_t|\mathcal{F}_s] = e^{sE} e^{T_s} e^{\mathcal{S}([0,s])} (e^{-(t-s)(H_{\mathbb{Z}_2}-E)} \varphi_g)(q_s) = Y_s$$

and the lemma follows. qed

4.3 Upper estimates on bound states

We will use the following conditions.

Assumption 4.4 *The following properties hold:*

- (1) $b_3 \in L^\infty$ and $W = \sqrt{b_1^2 + b_2^2} \in L^\infty$.
- (2) With $m_* = \|b_3\|_\infty + \|W\|_\infty$, we have $m_* < m^2/2$.
- (3) V is of relativistic Kato-class, i.e.,

$$\lim_{t \downarrow 0} \sup_{x \in \mathbb{R}^3} \mathbb{E}_{P \times \nu}^{x,0} \left[\int_0^t V(B_{T_r}) dr \right] = 0. \quad (4.19)$$

Lemma 4.5 *If Assumption 4.4 holds, then $\varphi_g \in L^\infty(\mathbb{R}^3)$ and*

$$|\varphi_g(x, (-1)^\alpha)| \leq \mathbb{E}_{P \times \nu}^{0,0} \left[e^{(t \wedge \tau)E} e^{-\int_0^{t \wedge \tau} V(B_{T_r} + x) dr} e^{\frac{1}{2} T_{t \wedge \tau} m_*} \right] \|\varphi_g\| \quad (4.20)$$

for every stopping time τ with respect to $(\mathcal{F}_s)_{s \geq 0}$ and $t \geq 0$.

PROOF. Notice that $\varphi_g(x, (-1)^\alpha) = \mathbb{E}_M^{x,\alpha,0}[Y_t]$ for every t . Then Schwarz inequality yields that

$$\begin{aligned} & |\varphi_g(x, (-1)^\alpha)| \\ & \leq e^{tE} \mathbb{E}_M^{x,\alpha,0} \left[e^{2T_t} e^{-2 \int_0^t V(B_{T_r}) dr} e^{\int_0^{T_t} |b_3(B_r)| dr} e^{\int_0^{T_t} \log W(B_r) dN_r} \right]^{1/2} \mathbb{E}_M^{x,\alpha,0} [|\varphi_g(q_t)|]^{1/2} \\ & \leq e^{tE} \left(\mathbb{E}_{P \times \nu}^{0,0} [e^{-2 \int_0^t V(B_{T_r} + x) dr} e^{T_t m_*}] \right)^{1/2} \left(\mathbb{E}_M^{x,\alpha,0} [|\varphi_g(q_t)|^2] \right)^{1/2}. \end{aligned}$$

Here we used that $\mathbb{E}_\mu^0[e^{N_{T_t} \log W}] = e^{T_t(W-1)}$. Note that

$$\mathbb{E}_M^{x,\alpha,0} [|\varphi_g(q_t)|^2] = \int_0^\infty ds p_t(s) \int_{\mathbb{R}^3} \Pi_s(y) dy \sum_{n=0}^\infty |\varphi_g(x+y, (-1)^{\alpha+n})|^2 \frac{s^n}{n!} e^{-s},$$

where $p_t(s) = \frac{te^{tm}}{\sqrt{2\pi s^3}} e^{-\frac{1}{2}(\frac{t^2}{s} + m^2 s)} 1_{[0,\infty)}(s)$ denotes the distribution of subordinator T_t .

Since

$$\sum_{n=0}^{\infty} |\varphi_g(x+y, (-1)^{\alpha+n})|^2 \frac{s^n}{n!} e^{-s} \leq |\varphi_g(x+y, 1)|^2 + |\varphi_g(x+y, -1)|^2,$$

we obtain

$$\begin{aligned} & \mathbb{E}_M^{x,\alpha,0}[|\varphi_g(q_t)|^2] \\ & \leq \int_{\mathbb{R}^3} dy \int_0^\infty ds p_t(s) \Pi_s(y) (|\varphi_g(x+y, 1)|^2 + |\varphi_g(x+y, -1)|^2) \\ & = \int_{\mathbb{R}^3} dy (|\varphi_g(x+y, 1)|^2 + |\varphi_g(x+y, -1)|^2) \pi^{-2} \frac{te^{mt}}{(|y|^2 + t^2)^2} \int_0^\infty \xi e^{-(\xi + \frac{m^2(|y|^2 + t^2)/4}{\xi})} d\xi \\ & \leq C_t \|\varphi_g\|_{L^2(\mathbb{R}^3 \times \mathbb{Z}_2)}^2 \end{aligned}$$

with a constant C_t . Furthermore, let $m^2/(2m_*) > q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$. Then we get

$$\mathbb{E}_{P \times \nu}^{x,0}[e^{-2 \int_0^t V(B_{T_r}) dr} e^{T_t m_*}] \leq \left(\mathbb{E}_{P \times \nu}^{x,0}[e^{-2p \int_0^t V(B_{T_r}) dr}] \right)^{1/p} \left(\mathbb{E}_{P \times \nu}^{x,0}[e^{q T_t m_*}] \right)^{1/q}.$$

The first term at the right hand side above satisfies that

$$\sup_{x \in \mathbb{R}^3} \left(\mathbb{E}_{P \times \nu}^{x,0} \left[e^{-2p \int_0^t V(B_{T_r}) dr} \right] \right)^{1/p} < \infty \quad (4.21)$$

since V is of relativistic Kato class, and

$$\mathbb{E}_{P \times \nu}^{x,0}[e^{q T_t m_*}] = \mathbb{E}_\nu^0[e^{q T_t m_*}] = \int_0^\infty e^{q s m_*} \frac{te^{mt}}{\sqrt{2\pi s^3}} e^{-\frac{1}{2}(\frac{t^2}{s} + m^2 s)} ds = e^{+t(m - \sqrt{m^2 - 2q m_*})} < \infty. \quad (4.22)$$

Hence $\varphi_g \in L^\infty(\mathbb{R}^3)$. Notice that by the martingale property of $Y_t(x, \alpha)$,

$$\varphi_g(x, (-1)^\alpha) = \mathbb{E}_M^{0,0,0}[Y_{t \wedge \tau}(x, \alpha)] \quad (4.23)$$

for every stopping time τ and $t \geq 0$. (4.20) follows from (4.23) and

$$|\varphi_g(x, (-1)^\alpha)| \leq \mathbb{E}_{P \times \nu}^{0,0} \left[e^{(t \wedge \tau)E} e^{-\int_0^{t \wedge \tau} V(B_{T_r} + x) dr} e^{T_{t \wedge \tau} m_*/2} \right] \|\varphi_g\|.$$

qed

4.4 Decay of bound states: the case $V \rightarrow \infty$

In this subsection we show the spatial exponential decay of bound states of $H + V$ at infinity.

Lemma 4.6 *Let $\tau_R = \inf\{t \mid |B_{T_t}| > R\}$. Then τ_R is a stopping time with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$.*

PROOF. It suffices to show that $\{\tau_R \leq t\} \in \mathcal{F}_t$. Notice that

$$\{\tau_R \leq t\} = \bigcup_{\omega \in \Omega_\nu} (A(\omega), \omega),$$

where $A(\omega) = \{\omega' \in \Omega_P \mid \sup_{0 \leq s \leq t} |B_{T_s}(\omega')| > R\} \in \mathcal{F}_t^{(1)}(\omega)$. Thus $\{\tau_R \leq t\} \in \mathcal{F}_t^{(1)}$. Moreover

$$\{\tau_R \leq t\} = \bigcup_{\omega \in \Omega_P} (\omega, B(\omega)),$$

where $B(\omega) = \{\omega' \in \Omega_\nu \mid \sup_{0 \leq s \leq t} |B_{T_s}(\omega')(\omega)| > R\}$. Therefore $\{\tau_R \leq t\} \in \mathcal{F}_t^{(2)}$ and hence $\{\tau_R \leq t\} \in \mathcal{F}_t$. **qed**

Theorem 4.7 *If Assumption 4.4 holds and*

$$\lim_{|x| \rightarrow \infty} V(x) = \infty, \quad (4.24)$$

then for every $a > 0$ there exists $b > 0$ such that

$$|\varphi_g(x, (-1)^\alpha)| \leq b e^{-a|x|}. \quad (4.25)$$

PROOF. We have by Lemma 4.5 that

$$|\varphi_g(x, (-1)^\alpha)| \leq \left(\mathbb{E}_{P \times \nu}^{0,0} \left[e^{2(t \wedge \tau_R)E} e^{-2 \int_0^{t \wedge \tau_R} V(B_{T_r} + x) dr} \right] \right)^{1/2} \left(\mathbb{E}_{P \times \nu}^{0,0} [e^{m_* T_{t \wedge \tau_R}}] \right)^{1/2} \|\varphi_g\|.$$

Let $W(x) = W_R(x) = \inf\{V(y) \mid |x - y| < R\}$, and notice that

$$\lim_{|x| \rightarrow \infty} W(x) - E = \infty. \quad (4.26)$$

In particular, we may assume that $W(x) - E > 0$. This gives

$$\begin{aligned} & \left(\mathbb{E}_{P \times \nu}^{0,0} \left[e^{2(t \wedge \tau_R)E} e^{-2 \int_0^{t \wedge \tau_R} V(B_{T_r} + x) dr} \right] \right)^{1/2} \\ & \leq \left(\mathbb{E}_{P \times \nu}^{0,0} [e^{-2(t \wedge \tau_R)(W(x) - E)}] \right)^{1/2} \\ & \leq \left(\mathbb{E}_{P \times \nu}^{0,0} [1_{\{\tau_R < t\}} e^{-2(t \wedge \tau_R)(W(x) - E)}] \right)^{1/2} + \left(\mathbb{E}_{P \times \nu}^{0,0} [1_{\{\tau_R \geq t\}} e^{-2(t \wedge \tau_R)(W(x) - E)}] \right)^{1/2} \\ & \leq \left(\mathbb{E}_{P \times \nu}^{0,0} [1_{\{\tau_R < t\}}] \right)^{1/2} + e^{-t(W(x) - E)}. \end{aligned}$$

We see that

$$\mathbb{E}_{P \times \nu}^{0,0} [1_{\{\tau_R < t\}}] = \mathbb{E}_{P \times \nu}^{0,0} [1_{\{\sup_{0 \leq s \leq t} |B_{T_s}| - R \geq 0\}}] \leq \mathbb{E}_{P \times \nu}^{0,0} [e^{\alpha(\sup_{0 \leq s \leq t} |B_{T_s}| - R)}] \quad (4.27)$$

for any $\alpha \geq 0$. It can be shown that $(\mathbb{E}_{P \times \nu}^{0,0} [e^{\alpha \sup_{0 \leq s \leq t} |B_{T_s}|}])^{1/2} \leq C_1 e^{C_2 t}$ for sufficiently small α , see [CMS90, Proposition II.5]. Hence $(\mathbb{E}_{P \times \nu}^{0,0} [1_{\{\tau_R \leq t\}}])^{1/2} \leq e^{-\alpha R/2} C_1 e^{C_2 t}$, and

$$\left(\mathbb{E}_{P \times \nu}^{0,0} \left[e^{2(t \wedge \tau_R)E} e^{-2 \int_0^{t \wedge \tau_R} V(B_{T_r} + x) dr} \right] \right)^{1/2} \leq e^{-t(W(x) - E)} + e^{-\alpha R/2} C_1 e^{C_2 t}. \quad (4.28)$$

We also see that

$$\begin{aligned} \mathbb{E}_{P \times \nu}^{0,0} [e^{m_* T_{t \wedge \tau_R}}] &\leq \mathbb{E}_{P \times \nu}^{0,0} [1_{\{t < \tau_R\}} e^{m_* T_{t \wedge \tau_R}}] + \mathbb{E}_{P \times \nu}^{0,0} [1_{\{t \geq \tau_R\}} e^{m_* T_{t \wedge \tau_R}}] \\ &\leq \mathbb{E}_{P \times \nu}^{0,0} [e^{m_* T_t}] + \mathbb{E}_{P \times \nu}^{0,0} [1_{\{t \geq \tau_R\}} e^{m_* T_{\tau_R}}] \\ &\leq 2\mathbb{E}_{\nu}^0 [e^{m_* T_t}], \end{aligned}$$

where we used that $T_{\tau_R} \leq T_t$ for $\tau_R \leq t$. Thus we have

$$\mathbb{E}_{P \times \nu}^{0,0} [e^{m_* T_{t \wedge \tau_R}}]^{1/2} \leq \sqrt{2} e^{t(m - \sqrt{m^2 - 2m_*})/2} \quad (4.29)$$

by (4.22). Hence by (4.28) and (4.29),

$$|\varphi_g(x, (-1)^\alpha)| \leq \sqrt{2} (e^{-t(W(x) - E)} + e^{-\alpha R/2} C_1 e^{C_2 t}) e^{t(m - \sqrt{m^2 - 2m_*})/2} \|\varphi_g\|. \quad (4.30)$$

Notice that by inserting $R = p|x|$ with any $0 < p < 1$, $W(x) - E = W_R(x) - E = W_{p|x|}(x) - E \rightarrow \infty$ as $|x| \rightarrow \infty$. Thus substituting $t = \delta|x|$ for sufficiently small $\delta > 0$ and $R = p|x|$ with some $0 < p < 1$ in (4.30), the theorem follows. \square

4.5 Decay of bound states: the case $V \rightarrow 0$

In this subsection we consider the case of potentials decaying to zero as $|x| \rightarrow \infty$.

Theorem 4.8 *Let Assumption 4.4 hold and suppose that*

$$\lim_{|x| \rightarrow \infty} V(x) = 0. \quad (4.31)$$

Also, assume that

$$m - \sqrt{m^2 - 2m_*} < -2E. \quad (4.32)$$

Then there exist $a, b > 0$ such that

$$|\varphi_g(x, (-1)^\alpha)| \leq b e^{-a|x|}. \quad (4.33)$$

PROOF. Define $\tau_R = \tau_R(x) = \inf\{t \geq 0 \mid |B_{T_t} + x| \leq R\}$. Then τ_R is a stopping time, which can be seen in the same way as in Lemma 4.6. Thus

$$\begin{aligned} |\varphi_g(x, (-1)^\alpha)| &\leq \left(\mathbb{E}_{P \times \nu}^{0,0} \left[e^{2(t \wedge \tau_R)E} e^{-2 \int_0^{t \wedge \tau_R} V(B_{T_r} + x) dr} \right] \right)^{1/2} \left(\mathbb{E}_{P \times \nu}^{0,0} [e^{m_* T_{t \wedge \tau_R}}] \right)^{1/2} \|\varphi_g\| \\ &= \left(\mathbb{E}_{P \times \nu}^{x,0} \left[e^{2(t \wedge \tau_R(0))E} e^{-2 \int_0^{t \wedge \tau_R(0)} V(B_{T_r}) dr} \right] \right)^{1/2} \left(\mathbb{E}_{P \times \nu}^{x,0} [e^{m_* T_{t \wedge \tau_R(0)}}] \right)^{1/2} \|\varphi_g\|. \end{aligned}$$

We rewrite $\tau_R(0)$ by τ_R . Let $\varepsilon > 0$ be arbitrary. Then for sufficiently large R it follows that $\sup_{|x| > R} |V(x)| < \varepsilon$ by (4.31), and we see that $|\int_0^{t \wedge \tau_R} V(B_{T_r}) dr| \leq (t \wedge \tau_R)\varepsilon$. This gives

$$|\varphi_g(x, (-1)^\alpha)| \leq \left(\mathbb{E}_{P \times \nu}^{x,0} [e^{2(t \wedge \tau_R)(E+\varepsilon)}] \right)^{1/2} \left(\mathbb{E}_{P \times \nu}^{x,0} [e^{m_* T_{t \wedge \tau_R}}] \right)^{1/2} \|\varphi_g\|.$$

Thus

$$\begin{aligned} \mathbb{E}_{P \times \nu}^{x,0} [e^{2(t \wedge \tau_R)(E+\varepsilon)}] &= \mathbb{E}_{P \times \nu}^{x,0} [1_{\{t \leq \tau_R\}} e^{2t(E+\varepsilon)}] + \mathbb{E}_{P \times \nu}^{x,0} [1_{\{t > \tau_R\}} e^{2\tau_R(E+\varepsilon)}] \\ &\leq e^{2t(E+\varepsilon)} + C_1 e^{-m_\varepsilon |x|} \end{aligned}$$

by making use of [CMS90, (II.29)(II.22) and (IV.3)] as above, where

$$m_\varepsilon = \begin{cases} m & \text{if } 2|E| > m \\ 2\sqrt{m|E| - |E|^2} & \text{if } 2|E| \leq m. \end{cases}$$

Also, notice that

$$\mathbb{E}_{P \times \nu}^{x,0} [e^{m_* T_{t \wedge \tau_R}}] \leq 2e^{t(m - \sqrt{m^2 - 2m_*})}.$$

Therefore

$$|\varphi_g(x, (-1)^\alpha)| \leq (e^{t(E+\varepsilon)} + C_1 e^{-m_\varepsilon |x|/2}) \sqrt{2} e^{t(m - \sqrt{m^2 - 2m_*})/2}. \quad (4.34)$$

On inserting $t = \delta|x|$ with sufficiently small δ , the theorem follows from (4.32). **qed**

Acknowledgments: FH acknowledges support of Grant-in-Aid for Science Research (B) 20340032 from JSPS and Grant-in-Aid for Challenging Exploratory Research 22654018 from JSPS, and thanks the hospitality of ICMS Edinburgh, where part of this work was done. JL also thanks ICMS Edinburgh for a Research-in-Group grant allowing to complete the project of this paper. We thank Toshimitsu Takaesu for pointing out typing errors in the first version of the manuscript.

References

- [ARS91] De Angelis, G. F., Rinaldi, A. and Serva, M.: Imaginary-time path integral for a relativistic spin-(1/2) particle in a magnetic field, *Europhys. Lett.* **14** (1991), 95–100.

- [Car78] Carmona, R.: Pointwise bounds for Schrödinger eigenstates *Commun. Math. Phys.* **62** (1978), 97–106.
- [CMS90] Carmona, R., Masters, W. C. and Simon, B.: Relativistic Schrödinger operators: asymptotic behavior of the eigenfunctions, *J. Funct. Anal.* **91** (1990), 117–142.
- [HIL09] Hiroshima, F., Ichinose, T. and Lőrinczi, J.: Path integral representation for Schrödinger operator with Bernstein function of the Laplacian, *Rev. Math. Phys.* **24** (2012), 1250013.
- [HL08] Hiroshima, F. and Lőrinczi, J.: Functional integral representation of the Pauli-Fierz model with spin 1/2, *J. Funct. Anal.* **254** (2008), 2127–2185.
- [LS81] Leinfelder, H. and Simader, C.G.: Schrödinger operators with singular magnetic potentials, *Math. Z.* **176** (1981), 1–19.
- [LHB09] Lőrinczi, J., Hiroshima, F. and Betz, V.: *Feynman-Kac-Type Theorems and Gibbs Measures on Path Space. With Applications to Rigorous Quantum Field Theory*, De Gruyter Studies in Mathematics **34**, 2011.