

## University Library

Author/Filing Title ..... TALIB .....

Class Mark ..... T .....

Please note that fines are charged on ALL  
overdue items.

**FOR REFERENCE ONLY**

040286039X



|  
|  
|  
|  
|  
|  
|  
|  
|

# **New Classes Of Ligands For Asymmetric Synthesis**

**A thesis presented by**

**Salem Ali Talib**

**In Partial Fulfilment of the Requirement for the Award of**


**Doctor of Philosophy**

*Supervised by*

**Professor Philip C. B. Page**



Department of Chemistry  
Ashby Road  
Loughborough  
Leics LE11 3TU

 <b>Loughborough University</b> Public Library
Date <i>May 21</i>
Class
Acc No. <i>040286039</i>

## Abstract

This thesis is divided into six chapters.

The first is a review of general asymmetric synthesis, and considers in detail the palladium catalysed allylic substitution reaction, the 1,4-conjugate addition and the Heck reaction.

The second deals with our general synthetic approach to the chiral ligands. Here we synthesized a range of ligands derived from chiral amino alcohols condensed with aryl and aliphatic ketones. We also synthesized chiral sulfur imine and phosphorus imine ligands.

The third deals with applications of the ligands in the above reactions, and discusses the most successful ligands. In the palladium catalysed allylic displacement reaction, the sulfur imine ligands were the most successful ligands with ee of 96%. In the case of the 1,4 conjugate addition of diethylzinc to cyclic and acyclic enones, we were able to achieve excellent results using the phosphorus imine and the S<sub>N</sub> ligands derived from pseudoephedrine and ketones, ee of >99% were obtained.

Chapter four deals with asymmetric sulfoxidation and the effect of electron donating and withdrawing groups on the sulfoxidation. Here we demonstrated the inductive influence of the substituent on the ee of the sulfoxide.

Chapter five deals with the conclusion.

The sixth part of this thesis deals with the experimental procedures undertaken in this work.

### **Acknowledgement**

I would like to thank my supervisor, Professor P C B Page, for his advice and constant help. I would also like to thank Professor Harry Heaney for his invaluable contribution and advice throughout.

My thanks also go to my Industrial supervisor, Dr J Liddle, for his contribution and assistance.

I would also like to thank Dr G Weaver for his help in proof reading this thesis.

Last but not least I would like to thank GSK and EPSRC for their financial contribution.

## Contents

	<i>Page</i>
Abstract	i
Acknowledgements	ii
Contents	iii
List of Abbreviations	iv
<b>Chapter 1: Introduction</b>	<b>1</b>
1.1 Introduction	2
1.2 References	25
<b>Chapter 2: Design of Ligands</b>	<b>32</b>
<b>Chapter 3: Testing the Ligands; “The Moment of Truth”</b>	<b>71</b>
3.1 Palladium Allylic Displacement Reaction	72
<b>Chapter 4: Effect of Substituent Groups on Enantioselectivity of Sulfoxidation</b>	<b>122</b>
<b>Chapter 5: Conclusion</b>	<b>137</b>
<b>General Experimental Procedures</b>	<b>139</b>
<b>References</b>	<b>269</b>
<b>Appendices</b>	<b>275</b>

## List of Abbreviations

<b>Ac</b>	acetyl
<b>Ar</b>	aryl
<b>AIBN</b>	azobis-isobutyronitrile
<b>ATP</b>	adenosine 5'-triphosphate (and related compounds)
<b>9-BBN</b>	9-borabicyclo[3.3.1]nonane
<b>BINAP</b>	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
<b>BINOL</b>	1,1'-bis-2-naphthol
<b>Bn</b>	benzyl
<b>Boc</b>	<i>tert</i> -butoxycarbonyl
<b>Bz</b>	benzoyl
<b>CAN</b>	cerium(IV) ammonium nitrate
<b>Cbz</b>	carbobenzyloxy
<b>COD</b>	cycloocta-1,5-diene
<b>COSY</b>	correlation spectroscopy
<b>Cp</b>	cyclopentadienyl ligand
<b>CSA</b>	camphorsulfonic acid
<b>Cy</b>	cyclohexyl
<b>DABCO</b>	1,4-diazabicyclo[2.2.2]octane
<b>DAST</b>	diethylaminosulfur trifluoride
<b>DBA</b>	(or dba as a ligand) dibenzylideneacetone

<b>DBU</b>	1,8-diazabicyclo[5.4.0]undec-7-ene
<b>DCC</b>	1,3-dicyclohexylcarbodiimide
<b>DCE</b>	dichloroethane
<b>DCM</b>	dichloromethane
<b>de</b>	diastereomeric excess
<b>DEAD</b>	diethyl azodicarboxylate
<b>(+)-DET</b>	(+)-(R,R)-diethyl tartrate
<b>DIBAL</b>	diisobutylaluminium hydride
<b>DMA</b>	N,N-dimethylacetamide
<b>DMAD</b>	dimethyl acetylenedicarboxylate
<b>DMAP</b>	4-(N,N-dimethylamino)pyridine
<b>DME</b>	1,2-dimethoxyethane
<b>DMF</b>	N,N-dimethylformamide
<b>DMPU</b>	1,3-dimethyl-3,4,5,6-tetrahydropyrimidin-2(1H)-one
<b>DMS</b>	dimethyl sulfide
<b>DMSO</b>	dimethyl sulfoxide
<b>DNA</b>	deoxyribonucleic acid
<b>EDCI</b>	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide
<b>EDTA</b>	ethylenediaminetetraacetic acid
<b>ee</b>	enantiomeric excess
<b>Et</b>	ethyl
<b>FAB</b>	fast atom bombardment

<b>Fc</b>	ferrocene
<b>Fmoc</b>	fluoren-9-ylmethoxycarbonyl
<b>HMPA</b>	hexamethylphosphoramide
<b>HMPT</b>	hexamethylphosphorus triamide
<b>HPLC</b>	high-performance liquid chromatography
<b>Hz</b>	Hertz
<b>IR</b>	infrared
<b>J</b>	coupling constant in NMR spectroscopy
<b>LAH</b>	lithium aluminium hydride
<b>LDA</b>	lithium diisopropylamide
<b>Li(K)HMDS</b>	lithium (potassium) hexamethyldisilazide
<b>MCPBA</b>	<i>m</i> -chloroperbenzoic acid
<b>Me</b>	methyl
<b>MEM</b>	(2-methoxyethoxy)methyl
<b>Mes</b>	2,4,6-trimethylphenyl (mesityl) (not methylsulfonyl)
<b>MOM</b>	methoxymethyl
<b>Ms</b>	methanesulfonyl (mesyl)
<b>Naph</b>	Naphthyl
<b>NBS</b>	<i>N</i> -bromosuccinimide
<b>NCS</b>	<i>N</i> -chlorosuccinimide
<b>NIS</b>	<i>N</i> -iodosuccinimide
<b>NMO</b>	4-methylmorpholine <i>N</i> -oxide

<b>nOe</b>	nuclear Overhauser enhancement effect/ nuclear Overhauser effect
<b>Nu</b>	nucleophile
<b>PBN</b>	<i>N-tert-butyl-<math>\alpha</math>-phenylnitrone</i> ( <i>N</i> -benzylidene- <i>tert</i> - butyl amine N-oxide
<b>PCC</b>	pyridinium chlorochromate
<b>PDC</b>	pyridinium dichromate
<b>PEG</b>	poly(ethylene glycol)
<b>Ph</b>	phenyl
<b>PPTS</b>	pyridinium toluene- <i>p</i> -sulfonate
<b>P r</b>	propyl
<b>P r i</b>	<i>isopropyl</i>
<b>PTSA</b>	toluene- <i>p</i> -sulfonic acid
<b>QSAR</b>	quantitative structure activity relationship
<b>RNA</b>	ribonucleic acid
<b>RT</b>	room temperature
<b>SPMB</b>	(4-methoxyphenyl) methane thiol
<b>TBAF</b>	tetrabutylammonium fluoride
<b>TBDMS</b>	<i>tert</i> -butyldimethylsilyl
<b>TBDPS</b>	<i>tert</i> -butyldiphenylsilyl
<b>TCNE</b>	tetracyanoethylene
<b>TCNQ</b>	tetracyanoquinodimethane
<b>TEMPO</b>	tetramethylpiperidine- <i>N</i> -oxide

<b>Tf</b>	trifluoromethylsulfonyl (triflyl)
<b>TFA</b>	trifluoroacetic acid
<b>TFAA</b>	trifluoroacetic anhydride
<b>THF</b>	tetrahydrofuran
<b>THP</b>	tetrahydropyran
<b>TIPS</b>	triisopropylsilyl
<b>TMEDA</b>	<i>N,N,N',N'</i> -tetramethylethylenediamine
<b>TMS</b>	trimethylsilyl or tetramethylsilane (can be used for either as long as no ambiguity results)
<b>TPAP</b>	tetrapropylammonium perruthenate
<b>Tr</b>	trityl
<b>tris</b>	tris(hydroxymethyl)aminomethane
<b>Ts</b>	toluene- <i>p</i> -sulfonyl
<b>TTF</b>	tetrathiafulvalene

# **CHAPTER 1**

## **Introduction**

## 1.1

## INTRODUCTION

Biological systems in general, tend to recognise each member of a pair of enantiomers as different substances, and the two enantiomers will elicit different responses. Thus one enantiomer may act as a very effective therapeutic drug whereas the other enantiomer could be highly toxic. The sad case of thalidomide is a well-known example. It has been shown that only one enantiomer produces the desired activity and the other is totally inactive or toxic. Another example of this is L-DOPA.

The importance and practicality of asymmetric synthesis as a tool to obtain enantiomerically pure or enriched compounds is recognised by chemists in synthetic organic chemistry, medicinal chemistry, agricultural chemistry, natural product chemistry, pharmaceutical industries and agricultural industries. This prominence is due to the explosive development of new and more efficient methods during the last decade.

Among the types of asymmetric reaction, the most desirable and most challenging is catalytic asymmetric synthesis because one chiral molecule can create millions of chiral product molecules, just as enzymes do in biological systems. Among the significant achievements in asymmetric synthesis are asymmetric hydrogenation,<sup>1,2</sup> the Sharpless epoxidation,<sup>3,4</sup> and the transition-metal catalysed allylic displacement reaction.<sup>5-7</sup>

### **Transition metal catalysed allylic displacement**

The transition metal catalysed allylic displacement reaction presents an advantage to form many different types of carbon-carbon and carbon-heteroatom bonds.<sup>8-10</sup>


The transition metal catalysed allylic displacement reactions have emerged as one of the more powerful tools for the controlled introduction of various chemical bonds into organic compounds (Scheme 1).




**Scheme 1**

The reaction involves a ( $\pi$ -allyl) metal complex as a key intermediate, which can be exploited for various transformations with high chemo, regio and stereoselectivities.<sup>11-13</sup>


The mechanism of metal catalysed allylic displacement reaction is generally believed to involve the four fundamental steps shown in Scheme 2.




The key feature of the catalytic cycle is the intermediate ( $\pi$ -allyl) metal complex . Its generation and subsequent reaction represent the bond-breaking and making event in which the source of chiral induction can be derived. The  $\pi$ -allyl complex may undergo many useful transformations; for example the conjugated 4,6-dienone shown in Scheme 2.1 may be prepared from a steroidal enone *via* a  $\pi$ -allyl palladium complex.<sup>14</sup> Catalytic oxidative dehydrogenation of ethyl 3-butenedicarboxylate, Scheme 2.2, proceeds via a  $\pi$ -allyl palladium complex.<sup>15</sup> Oxidation of a  $\pi$ -allyl palladium complex with peracids produces allylic alcohols regioselectively with retention of configuration (Scheme 2.3).<sup>16</sup>



**Scheme 2.1**




Scheme 2.2



Scheme 2.3


Isomerization of the double bond in vitamin D analogues such as calciferol occurs by an oxidation reduction sequence via  $\pi$ -allyl palladium complex (Scheme 2.4).<sup>17</sup>



**Scheme 2.4**


Depending on the structure of substrate every step may provide an opportunity for enantioselection with the exception of the decomplexation, which occurs after bond formation.

Although ionization proceeds with inversion of stereochemistry, regardless of the nucleophile, the addition of a nucleophile can involve two pathways in which the nature of the nucleophile leads to a different stereochemical outcome (Scheme 3).<sup>18-20</sup>



Scheme 3

A unique feature of the transition metal catalysed allylic displacement is its ability to convert starting materials of various types, such as racemic, meso and achiral compounds, into optically pure material. Strategies which effect such transformations are derived from recognition of the stereochemical courses in each step of the catalytic cycle (Scheme 2), and analysis of symmetry elements in the substrate or intermediate. Scheme 4 summarizes potential sources of enantio-discrimination in transition-metal catalysed allylic displacement.



Scheme 4

The first strategy (A) involves discrimination between enantiotopic leaving groups, the transition metal catalyst can be made to choose deftly between the two leaving groups.<sup>21</sup>

The second strategy (B), involves two enantiomers of a racemic substrate converge into a meso  $\pi$ -allyl complex wherein preferential attack of the nucleophile at one of either allylic terminal leads to asymmetric induction.<sup>22a</sup>

If the  $\eta^3$ -allylic intermediate is not symmetrically 1,3-disubstituted, enantioselection will be dictated by which face of the allylic fragment is presented to the nucleophile. This is represented by (C).<sup>22b</sup>

The fourth strategy shown in (D), involves the transition metal testing both faces of the olefin, but only one face will lead to metal ionization,<sup>23</sup> this strategy only works if the olefin is not symmetrically distributed.

Enantioselectivity can also be achieved if the nucleophile is prochiral or is an equilibrating mixture of racemic nucleophiles. Here, enantioselectivity is achieved by enantioface discrimination of the nucleophile moiety (E).

Examples of chiral ligands used in allylic displacement are shown in Fig 1






Fig 1

### Zinc mediated conjugate addition


Another very important reaction is the Michael addition to  $\alpha, \beta$ -unsaturated systems which is a fundamental process in organic chemistry, because it forms carbon-carbon (Scheme 5.1),<sup>24</sup> carbon-nitrogen (Scheme 5.2),<sup>25</sup> carbon-sulfur (Scheme 5.3)<sup>26</sup> or carbon-oxygen bonds at the  $\beta$  position of the carbonyl group and results in highly functionalized compounds with one or two asymmetric centres.



Scheme 5.1




Scheme 5.2



Scheme 5.3

The golden age of organocopper chemistry began in 1941 when Kharasch and Tawney<sup>27</sup> reported the 1,4 addition reaction of Grignard reagent to an  $\alpha, \beta$ -unsaturated ketone in the presence of a small amount of  $\text{Cu}^{\text{I}}$  salt.<sup>28</sup> Gilman reported in 1952 that the addition of one equivalent of  $\text{MeLi}$  to a  $\text{Cu}(\text{I})$  salt resulted in the formation of a yellow precipitate, which then afforded a colourless solution upon addition of another equivalent of  $\text{MeLi}$  (Scheme 5.4).<sup>29</sup>



Scheme 5.4

In 1966, Costa isolated a complex between phenylcopper (I) and magnesium, as well as crystals of lithium diphenyl cuprate (I) complex.<sup>30</sup>

The organic chemistry of organocuprates started its rapid development in 1966, when House showed that the reactive species of conjugate addition is the lithium diorganocuprate (I) called a Gilman reagent.<sup>31</sup> Foundations for subsequent vigorous synthetic development were laid by Corey and Posner.<sup>32</sup>

Shortly after the opening of the Lewis acid era with the discovery of the Mukaiyama aldol reaction in 1973,<sup>33</sup> Yamamoto and Maruyama's reports on the  $\text{RCuBF}_3$  reagents<sup>34</sup> introduced the new concept of Lewis acid assistance in organocopper chemistry.<sup>35</sup> Although the identity of the  $\text{RCuBF}_3$  moiety is still elusive,<sup>36</sup>  $\text{BF}_3$  activation was applied to numerous synthetic work, as illustrated by a diastereoselective addition of a homoenolate species in the total synthesis of cortisone.<sup>37</sup>


Since the initial discovery by Nakamura and Kuwajima in 1984,<sup>38</sup>  $\text{Me}_3\text{SiCl}$  has become a standard reagent for acceleration of conjugate addition. This effect was reported first for the copper-catalysed conjugate addition of the zinc homoenolate of propanoic acid ester, then for Grignard-based catalytic reagents<sup>39,40</sup> and stoichiometric lithium diorganocuprates.<sup>41-45</sup>

The zinc homoenolate started the chemistry of metal homoenolates. (Scheme 5.5).<sup>46-48</sup> The synthetic scope of such a nucleophile-bearing electrophile was further exploited by Tamaru, Yoshida, and co-workers first,<sup>49</sup> and then extensively by Knochel and others.<sup>50-51</sup> Dominated by lithium and magnesium-based systems until the mid 1980s, organocopper chemistry now routinely utilizes much milder organometallic nucleophiles such as organozinc, titanium,<sup>52</sup> zirconium,<sup>53,54</sup> and

aluminium reagents.<sup>55</sup> With the aid of proper activators, these mildly reactive reagents show selectivities unavailable with the conventional reagents, as illustrated in Scheme 5.6 for Me<sub>3</sub>SiCl-dependent chemoselectivity.<sup>56</sup>




Scheme 5.5



Scheme 5.6


Organocopper chemistry is still rapidly expanding its synthetic scope. The scope of carbocupration, previously limited to acetylenes, has been extended to olefins.<sup>57-59</sup> Enantioselective conjugate addition<sup>60</sup> has become truly useful through the use of dialkylzinc, a cationic copper catalyst and a chiral ligand (Scheme 6).

In 1997 Feringa reported one of the most enantioselective conjugate addition to date of dialkylzinc reagents, catalysed by chiral copper phosphoramidite complexes.<sup>61</sup> The reaction is mediated by 2 mol % Cu(OTf)<sub>2</sub> and 4 mol % chiral ligand (Scheme 6), and he reported an ee of 92% when the reaction was carried at -35 °C.




Scheme 6

A possible pathway for the 1,4-addition is shown in Scheme 7. This involves the transfer of an alkyl fragment from  $\text{R}_2\text{Zn}$  to the copper complex,<sup>62</sup> followed by  $\pi$ -complexation of the resulting copper alkyl species to the double bond of the enone and of the alkylzinc ion to the enone-carbonyl. Next, alkyl transfer to the  $\beta$ -position of the enone generates a zinc enolate, which upon protonation provides the 1,4-adduct.




Scheme 7


The presence of a ligand is fundamental, as without it the reaction is very slow and with many side products found. Some examples of chiral ligands used in this type of reaction are shown in Fig 2.




Feringa 1997  
(2a)



Kobayashi 1994  
(2b)



Alexakis 1997  
(2c)




Ming 1999 X=binaphthyl  
(2d)

Fig 2


The asymmetric 1,4-addition of diethylzinc to cyclohex-2-enone (Scheme 6) has been well developed and is well understood, and using this methodology it has been possible to incorporate stereoselectively functionalized cyclohexane and larger rings in natural product synthesis.<sup>63</sup> In contrast, the asymmetric 1,4-conjugate addition to cyclopent-2-enone (Scheme 8) presents more of a challenge. The importance of the cyclopentane skeleton stems for example from its presence in prostaglandins,<sup>64</sup> and alkaloids including dendrobine.<sup>65</sup> Recent reports by Chan,<sup>66</sup>

Pfaltz<sup>67</sup> and Hoveyda<sup>68</sup> give ees of up to 97% for the asymmetric 1,4-addition of diethylzinc to cyclopent-2-enone at  $-35\text{ }^{\circ}\text{C}$ .



**Scheme 8**


Attempts by Pfaltz<sup>67</sup> to apply the asymmetric 1,4-addition to acyclic enones (Scheme 9) showed little success with ees achieved of only up to 50% when the reactions were carried out at  $-35\text{ }^{\circ}\text{C}$ .




**Scheme 9**

### Heck reaction

In recent years there has been considerable interest in the use of the Heck-type arylation and alkenylation of olefins for constructing carbon skeletons of biologically important organic compounds.<sup>69,70</sup> Development of the asymmetric Heck reaction, where the carbon-carbon bond formation proceeds with high enantioselectivity, would provide new efficient routes to the optically active compounds. Pioneering work in this area by Shibasaki<sup>71</sup> and Overman<sup>72</sup> has opened the way for the asymmetric Heck reaction. The catalytic cycle, shown below in Scheme 10, is termed the neutral pathway, which is followed for unsaturated halide substrate.




However, there is also a cationic pathway, which is shown below in Scheme 11. This pathway is followed by unsaturated triflate substrates, or for unsaturated halides in the presence of halide scavengers such as Ag(I) or Tl(I) salts.<sup>73,74</sup> The individual steps are similar to the neutral pathway, but the difference between the two mechanisms lies in the nature of palladium(II) intermediate, which is now cationic, and this difference has a marked effect on the reactivity and stereoselectivity.




Scheme 11

The salient features of the cationic pathway shown in Scheme 11 were deduced by Cabri<sup>75</sup> and Hayashi.<sup>76</sup> Subsequent to oxidative addition, a vacant co-ordination site is generated by triflate dissociation or by halide abstraction by  $Ag(I)$ ; this vacant site facilitates double bond coordination to form the cationic intermediate which ultimately results in the formation of the Heck product.


The key feature of this reaction is the removal of an anionic ligand from the palladium coordination sphere, which allows the coordination of the alkene while the integrity of the chiral ligand is maintained. Ozawa<sup>77</sup> reported the application of the ligand (3) in the asymmetric Heck reaction (Scheme12), and published ees exceed 96%.




A common feature of the ligands involved in the palladium catalysed allylic displacement reaction, 1,4-addition and the Heck reaction is that in most cases nitrogen appears as one of the ligating atoms. The need to balance the steric and electronic conditions in order to achieve the best results in the catalytic reaction<sup>78</sup> (Scheme 13) may dictate the need to introduce substituents at nitrogen. The usual method is the application of the Eschweiler-Clarke approach.<sup>79</sup> The reaction involves heating a primary or secondary amine with formaldehyde and formic acid to yield the tertiary amine. The hydride acceptor is the iminium ion, from condensation of the amine with formaldehyde. The resulting *N*-methyl amino alcohol core structures are found in several bioactive molecules.<sup>80,81</sup>



$\alpha$ -Amino alcohols are important intermediates in many organic synthesis, but there are few methods to generate these compounds. One method is to construct an oxazolidine ring, which is obtained by condensation of an amino alcohol and aldehyde or ketones, and then reductively cleave the ring to give the  $\alpha$ -amino alcohol.<sup>82</sup> An alternative strategy for the synthesis of  $\alpha$ -amino alcohols is through the synthesis of cyanohydrins derived from aldehydes or ketones followed by reduction using lithium aluminium hydride,<sup>83</sup> or by reduction of  $\alpha$ -amino acids.




The oxazolidine ring system **4** has been studied extensively by Knorr and coworkers<sup>83</sup> who ascribed the cyclic structure to products which were obtained, e.g. from ethanolamine and aldehyde or ketones with loss of a molecule of water (Scheme 14), without taking into account the possibility that the condensation product might simply be the Schiff base (Scheme 15).



ha

Scheme 14




Scheme 15

In 1951 McCasland and Horswill<sup>86</sup>, who were investigating some complex cases arrived at the conclusion that the product structures were not well-established. The formation of oxazolidines from aldehydes and simple aliphatic ketones has since been studied extensively.<sup>87-90</sup> However their formation from aromatic ketones has not been reported prior to our work.<sup>91</sup>

Since the first reports of  $\pi$ -allyl palladium catalysed allylic substitution,<sup>92</sup> and the subsequent reviews by Trost,<sup>93</sup> many examples of chiral ligands have been synthesized for application in this reaction, and here we report novel approaches to the synthesis of some ligands to be applied in this reaction. However, few attempts have been made to use thioethers. Thioethers are powerful ligands for transition metals, and the affinity of thioether groups for transition metals has been used to direct the outcome to stoichiometric reactions mediated by transition metals.<sup>94</sup>


Recently, Krafft and co-workers have shown that the Pauson-Khand reaction gives excellent regiocontrol when a thioether is present in the substrate.<sup>95</sup> The most dramatic advance has been the demonstration that a sulfur atom in the substrate can exert an influence in the palladium-catalysed alkylation, dictating the regiochemistry of nucleophilic attack.<sup>96</sup> More recently, thioethers have proved to modify reactivity in the platinum-catalysed hydrosilylation of olefins (Scheme 16).<sup>97</sup> The importance of the thioether moiety was demonstrated in a competition experiment, in which homoallyl sulfide was mixed with octene and subjected to hydrosilylation at room temperature; only the thioether substrate underwent hydrosilylation. Hydrosilylation of octene fails under similar conditions, demonstrating that homoallylic thioether was accelerating an otherwise sluggish reaction.<sup>97</sup>



**Scheme 16**

The final class of ligand which we consider here are those containing sulfoxides. The synthesis of sulfoxides has been widely reviewed. The efficiency of the sulfoxide group in asymmetric synthesis is thought to be due to the presence of three different groups at the asymmetric centre; while nitrogen tends to be hindered by pyramidal inversion this is not the case with sulfur, where pyramidal inversion is not observed below 150 °C.<sup>99</sup>

Since the first reports of sulfoxidation by Belenovic<sup>100</sup> and Montanovi,<sup>101</sup> many authors have attempted asymmetric sulfoxidation with the modified Sharpless asymmetric epoxidation,<sup>102</sup> adopted independently by Kagan<sup>103</sup> and Modena<sup>104</sup> (Scheme 17).



**Scheme 17**

Kagan modified the Sharpless procedure by adding one equivalent of water, a poison in the epoxidation reaction, and increased the ratio of (+) DET to titanium isopropoxide to 2:1 (Scheme 18). Under these conditions, double bonds, amines, alcohols and phenols are not oxidized.

Davis reported the use of *N*-sulfonyl oxaziridines as sulfoxidation reagents.<sup>105,106</sup> Later, Davis reported the use of cyclic oxaziridines based upon camphor sulfonic acid,<sup>107</sup> and this was later modified by Page.<sup>108</sup> In oxaziridines the nitrogen is a stable chiral centre, and electron-releasing groups tend to stabilize oxaziridines, reducing their electrophilicity. In the camphor system, the sulfonyl group is electron-withdrawing, rendering these compounds highly electrophilic, a property exploited in their use for sulfur oxidation.<sup>109</sup>

To summarize, the aim of the work described in this thesis is to design ligands which show multipurpose usage in asymmetric synthesis. The main focus of our design is to synthesize ligands for the asymmetric palladium catalysed allylic substitution reaction, the asymmetric 1,4-addition of dialkylzinc to enones, and

the asymmetric Heck reaction. Additionally it was intended to investigate the efficiency of the Page sulfoxidation reagent in the presence of electron withdrawing groups (EWG) and electron donating groups (EDG) in the substrate molecules.

## 1.2 REFERENCES

1. Knowles, W.S.; Sabacky, M.J.; Vineyard, B.D. *J. Chem. Soc. Chem. Commun.* **1972**, 10.
2. Vineyard, B.D.; Knowles, W.S.; Sabacky, M.J.; Bachman, G.L.; Weinkauff, D.F. *J. Am. Chem. Soc.* **1977**, 99, 5946.
3. Katsuki, T.; Sharpless, K.B. *J. Am. Chem. Soc.* **1980**, 102, 5974.
4. Hanson, R.M.; Sharpless, K.B. *J. Org. Chem.* **1986**, 51, 1922.
5. Trost, B.M.; Strege, P.E. *J. Am. Chem. Soc.* **1977**, 99, 1610.
6. Tsuji, J. *Tetrahedron* **1986**, 42, 4361.
7. Trost, B.M.; Paterson, D.E. *J. Org. Chem.* **1998**, 63, 1339.
8. Consiglio, G.; Waymouth, R.M. *Chem. Rev.* **1989**, 89, 257.
9. Heumann, A.; Reglier, M. *Tetrahedron* **1995**, 51, 975,
10. Reiser, O. *Angew. Chem. Int. Ed.* **1993**, 32, 547.
11. Noyori, R.J.; Kitamura, M. *Angew. Chem. Int. Ed.* **1991**, 30, 49.
12. Oguni, N.; Matsuda, Y.; Kaneko, T. *J. Am. Chem. Soc.* **1988**, 110, 7877.
13. Kitamura, M.; Okada, S.; Suga, S.; Noyori, R. *J. Am. Chem. Soc.* **1989**, 111, 4028.
14. Haynes, R.K.; Jackson, W.R.; Stragalinou, A. *Aust. J. Chem.* **1980**, 33, 1537.
15. Suzuki, T.; Tsuji, J. *Bull. Chem. Soc. Jpn.* **1973**, 46, 655.
16. Jones, D.N.; Knox, S.D. *J. Chem. Soc. Chem. Commun.* **1975**, 166.
17. Barton, D.H.R.; Patin, H. *J. Chem. Soc. Chem. Commun.* **1977**, 799.

18. Trost, B.M.; Strege, P.E. *J. Am. Chem. Soc.* **1975**, *97*, 2534.
19. Trost, B.M.; Verhoeven, T.R. *J. Org. Chem.* **1976**, *41*, 3215.
20. Hayashi, T.; Hagihara, T.; Konishi, M.; Kumada, M. *J. Am. Chem. Soc.* **1983**, *105*, 7767.
21. Trost, B.M.; van Vranken, D.L. *Angew. Chem. Int. Ed.* **1992**, *31*, 1992.
- 22a. Trost, B.M.; Murphy, D.J.; Konishi, M.; Kumada, M. *J. Am. Chem. Soc.* **1983**, *105*, 7767.      22b. Trost, B.M.; Sterge, P.E. *J. Am. Chem. Soc.* **1977**, *99*, 1650.
23. Murahashi, S.I.; Taniguchi, Y.; Imada, Y.; Tanigawa, Y. *J. Org. Chem.* **1989**, *54*, 3292.
24. Shimizu, S.; Ohori, K.; Arai, T.; Sasai, H.; Shibasaki, M. *J. Org. Chem.* **1998**, *63*, 7547.
25. Sibi, M.; Shay, J.J.; Liu, M.; Jasperse, C.P. *J. Am. Chem. Soc.* **1998**, *120*, 6615.
26. Nishimura, K.; Ono, M.; Nagaoka, Y.; Tomioka, K. *J. Am. Chem. Soc.* **1997**, *119*, 8959.
27. Kharash, M.S.; Tawney, P.O. *J. Am. Chem. Soc.* **1941**, *63*, 2309.
28. Paterson, J.M. *J. Org. Chem.* **1957**, *22*, 170.
29. Gilman, H.; Jones, R.G.; Woods, L.A. *J. Org. Chem.* **1952**, *17*, 1630.
30. Costa, G.; Camus, A.; Gatti, L.; Marsich, N. *J. Organomet. Chem.* **1966**, *5*, 568.
31. House, H.O.; Respess, W.L.; Whitesides, G.M. *J. Org. Chem.* **1966**, *31*, 3128.
32. Corey, E.J.; Posner, G.H. *J. Am. Chem. Soc.* **1967**, *89*, 3911.
33. Mukaiyama, T.; Banno, K.; Naraska, K. *J. Am. Chem. Soc.* **1974**, *96*, 7503.
34. Yamamoto, J.; Maruyama, K.J. *J. Am. Chem. Soc.* **1977**, *99*, 8068.

35. Smith, A.B.; Jerris, P.J. *J. Am. Chem. Soc.* **1981**, *103*, 194.
36. Lipshutz, B.H.; Ellsworth, E.L.; Dimock, S.H. *J. Am. Chem. Soc.* **1990**, *112*, 5869.
37. Horiguchi, Y.; Nakamura, E.; Kuwajima, I. *J. Am. Chem. Soc.* **1989**, *111*, 6257.
38. Nakamura, E.; Kuwajima, I. *J. Am. Chem. Soc.* **1984**, *106*, 3368.
39. Kuwajima, I.; Enda, J. *J. Am. Chem. Soc.* **1985**, *107*, 5493.
40. Nakamura, E.; Matsuzawa, S.; Horiguchi, Y.; Kuwajima, I. *Tetrahedron Lett.* **1986**, *27*, 4029.
41. Corey, E.J.; Boaz, N.W. *Tetrahedron Lett.* **1985**, *26*, 6015.
42. Alexakis, A.; Berlan, J.; Besace, Y. *Tetrahedron Lett.* **1986**, *27*, 1047.
43. Horiguchi, Y.; Matsuzawa, S.; Nakamura, E.; Kuwajima, I. *Tetrahedron Lett.* **1986**, *27*, 4025.
44. Linderman, A.; Godfrey, A. *Tetrahedron Lett.* **1986**, *27*, 4853.
45. Alexakis, A.; Frutos, J.C.; Mangeney, P. *Tetrahedron : Asymmetry* **1993**, *4*, 2427.
46. Nakamura, E.; Auki, S.; Oshino, H.; Kuwajima, I. *J. Am. Chem. Soc.* **1985**, *109*, 8056.
47. Nakamura, E.; Isaka, M.; Matsuzawa, S. *J. Am. Chem. Soc.* **1988**, *110*, 1297.
48. Nakamura, E.; Kuwajima, I. *J. Am. Chem. Soc.* **1977**, *99*, 7360.
49. Tamaru, Y.; Ochiai, H.; Nakamura, T.; Yoshida, Z. *Angew Chem. Int. Ed.* **1987**, *99*, 1193.
50. Knochel, P.; Yeh, M.C.P.; Berk, S.C.; Talbot, J. *J. Org. Chem.* **1988**, *53*, 2390.
51. Singer, R.D.; Knochel, P. *Chem. Rev.* **1993**, *93*, 2117.

52. Nakamura, E.; Arai, M.; Lipshultz, H. *J.Org. Chem.* **1991**, *56*, 5489.
53. Lipshultz, H.; Ellsworth, E. *J. Am. Chem. Soc.* **1990**, *112*, 7440.
54. Lipshultz, H.; Ellsworth, E.; Dimock, S. *J. Am. Chem. Soc.* **1990**, *112*, 5869.
55. Ibuka, T.; Minakata, H. *Synth. Commun.* **1980**, *10*, 125.
56. Arai, M.; Lipshultz, H.; Nakamura, E. *Tetrahedron* **1992**, *48*, 5709.
57. Stoll, A.; Negishi, E. *Tetrahedron Lett.* **1985**, *26*, 5671.
58. Berlan, J.; Beface, J.; Stephan, E.; Cresson, P. *Tetrahedron Lett.* **1985**, *26*, 5768.
59. Nakamura, E.; Isaka, M.; Matsuzawa, S. *J. Am. Chem. Soc.* **1988**, *110*, 1297.
60. Alexakis, A. in *Transition Metal Catalysed Reactions*, Blackwell Science, Oxford, UK. **1999**, 303.
61. Feringa, B.L.; Pineschi, M.; Arnold, L.A.; Imbos, R.; de vries, A.H.M. *Angew. Chem. Int. Ed.* **1997**, *36*, 2620.
62. Knochel, P.; Singer, R.D. *Chem. Rev.* **1993**, *93*, 2117.
63. Naaz, R.; Arnold, L.A.; Pineschi, M.; Keller, E.; Feringa, B.L. *J. Am. Chem. Soc.* **1999**, *121*, 1104.
64. Noyori, R. in *Asymmetric Catalysis in Organic Synthesis*, J. Wiley and Sons. New York, **1994**.
65. Roush, W.R. *J. Am. Chem. Soc.* **1980**, *102*, 1390.
66. Yan, M.; Chan, A.S.C. *Tetrahedron Lett.* **1989**, *40*, 6648.
67. Escher, I.H.; Pfaltz A. *Tetrahedron* **2000**, *56*, 2879.
68. Degrado, S.J.; Mizutani, H.; Hoveyda, A.H. *J. Am. Chem. Soc.* **2001**, *123*, 755.

69. Ashimori, A.; Bachand, B.; Overman, L.; Poon, D. *J. Am. Chem. Soc.* **1998**, *120*, 6477.
70. Sato, Y.; Sodeoka, M.; Shibasaki, M. *J. Org. Chem.* **1989**, *54*, 4738.
71. Honzawa, S.; Mizutani, T.; Shibasaki, M. *Tetrahedron Lett.* **1999**, *40*, 311.
72. Carpenter, N.E.; Kuceta, D.J.; Overman, L.E. *J. Org. Chem.* **1989**, *54*, 5845.
73. de Meijere, A.; Meyer, F.E.. *Angew. Chem. Int Ed.* **1994**, *33*, 2379.
74. Shibasaki, M.; Boden, C.D.J.; Kojima, A. *Tetrahedron* **1997**, *53*, 7371.
75. Gabri, W.; Candiani, J.; de Deymadis, S.; Francalanci, F.; Penco, S. *J. Org. Chem.* **1991**, *56*, 5796.
76. Ozawa, F.; Kubo, A.; Hayashi, T. *J. Am. Chem. Soc.* **1991**, *113*, 1417.
77. Ozawa, F.; Kubo, A.; Hayashi, T. *Tetrahedron Lett.* **1992**, *33*, 1485.
78. Van Haaren, R.J.; Goubitz, K.; Fraanje, J.; van Strijdonck, G.P.F.; Oeeving, Henk; Coussens, B.; Reek, J.N.H.; Kamer, P.C.J.; Van Leeuwen, P.W.N.M. *Inorg. Chem.* **2001**, *40*, 3363.
79. Pine, S.H.; Sanchez, B.L. *J. Org. Chem.* **1971**, *36*, 829.
80. Strattmann, K.; Burgoyne, D.L.; Moore, R.E.; Patterson, G.M.L.; Smith, C.D. *J. Org. Chem.* **1994**, *59*, 7219.
81. Petti, G.R.; Singh, S.B.; Herald, D.L.; Lloyd, P.; Kantosci, D.; Brukett, D.; Barkoczy, J.; Hogan, F.; Wardlaw, T.R. *J. Org. Chem.* **1994**, *59*, 6287.
82. Bergman, E.D. *Chem. Rev.* **1953**, *53*, 309.
83. Coates, R.M.; Vettel, P.R. *J. Org. Chem.* **1980**, *54*, 5430.
84. Knorr, L.; Mathess, H. *Ber.* **1901**, *34*, 3484.
85. Knorr, L.; Roessler, P. *Ber.* **1903**, *36*, 1278.
86. McCasland, G.E.; Horswell, E.C. *J. Am. Chem. Soc.* **1951**, *73*, 3933.

87. Kelly, R.; van Rheenen, V. *Tetrahedron Lett.* **1973**, 1709.
88. Contreras, R.; Santiesteban, F.; Paz-Sandoval, M.A. *Tetrahedron* **1984**, *40*, 3838.
89. Lambert, J.B.; Majchrzak, M.W. *J. Am. Chem. Soc.* **1980**, *102*, 3588.
90. Alva Astudillo, M.E.; Chokotho, N.C.J.; Jarvis, T.C.; Johnson, D.; Lewis, C.C. *Tetrahedron* **1985**, *41*, 5919.
91. Meyers, A.I.; Shimano, M. *Tetrahedron Lett.* **1993**, 4893.
92. Shaw, B.L.; Sheppard, N. *Chem. Ind. (London)* **1961**, 517.
93. Trost, B.M.; van Vrankev, D.L. *Chem. Rev.* **1996**, *96*, 365.
94. Holton, R.A.; Kjonaas, R.A. *J. Am. Chem. Soc.* **1997**, *99*, 4177.
95. Krafft, M.E.; Juliano, C.A.; Scott, I.L.; Wright, C.; McEachin, M.D. *J. Am. Chem. Soc.* **1991**, *113*, 1693.
96. Krafft, M.E.; Wilson, A.M.; Fu, Z.; Proctor, M.J.; Dasse, O.A. *J. Org. Chem.* **1998**, *63*, 1748.
97. Stratmann, K.; Burgoyne, D.L.; Moore, R.E.; Patterson, G.M.L.; Smith, C.D. *J. Org. Chem.* **1994**, *59*, 7219.
98. Carreno, A.J. *Chem. Rev.* **1995**, 1717.
99. Rayner, D.R.; Gordon, A.J.; Mislow, K.J. *J. Am. Chem. Soc.* **1968**, *90*, 4954.
100. Balenovic, K.; Bregnant, N.; Francetic, D. *Tetrahedron Lett.* **1960**, 20.
101. Mayer, A.; Montanari, F.; Tramotini, M. *Tetrahedron Lett.* **1961**, 607.
102. Pitchen, P.; Kagan, H.B.; Deshmukh, M.N.; Dunach, E. *J. Am. Chem. Soc.* **1984**, *106*, 8188.
103. Pitchen, P.; Kagan, H.B. *Tetrahedron Lett.* **1984**, *25*, 1049.
104. Di Furia, S.H.; Modena, G.; Seragilia, G. *Synthesis* **1984**, 325.

105. Davis, F.A.; Nadir, U.K.; Kluger, E.W. *Chem. Commun.* **1977**, 25.
106. Davis, F.A.; Jenkins, R.H.; Award, S.B.; Stringer, O.; Watson, W.H.; Galloy, J. *J. Am. Chem. Soc.* **1982**, *104*, 5412.
107. Davis, F.A.; Towson, J.C.; Carroll, P.J.; Weismiller, M.C.; Lal, S. *J. Am. Chem. Soc.* **1988**, *110*, 8477.
108. Page, P.C.B.; Heer, J.P.; Bethell, D.; Collington, E.W.; Andrews, D.M. *Tetrahedron Lett.* **1994**, *35*, 9629.
109. Katsuki, T.; Sharpless, K.B. *J. Am. Chem. Soc.* **1980**, *102*, 5974.

# **CHAPTER 2**


## **Ligand Design**

## LIGAND DESIGN


In attempting to design ligands, there are certain constraints which must be considered, these constraints often limit the scope or degree to which the ligands can function. Thus it is up to the designer to try and overcome those constraints. In designing ligands for complexing with transition metal the following constraints must be considered.

- a) The metal: The metal to which the ligand is complexing has a great influence on the outcome. Some metals complex to the ligand and never release it, others do not complex at all. Ligands can also form inert and labile complexes.
- b) The nature of the ligand heteroatoms: The binding nature of the heteroatom again has fundamental importance, the ability or otherwise of a heteroatom to act as a Lewis base is very important if complexation is to occur. The Lewis basicity has a great influence on the stereoselectivity; for example, as has been well documented for the palladium catalysed alkylation, the nucleophile approaches *trans* to the better  $\pi$  accepting heteroatom.<sup>1</sup>
- c) Type of reaction: Some ligands which are designed for alkylation might not be useful for epoxidation or other reactions and *vice versa*.

For the case of the allylic displacement reaction shown below in Scheme 18, the general catalytic cycle shown in Scheme 19,<sup>1</sup> must be considered in order to enable us to design a good ligand. The basic cycle consists of metal-olefin complexation, ionization, alkylation and decomplexation.



**Scheme18**




**Scheme 19**


To achieve asymmetric induction, the asymmetric environment of the ligands must be felt on the opposite face of the  $\pi$ -allyl unit, where bond making and breaking are occurring. For complexes containing bidentate ligands, it has been suggested that

regioselectivity is determined by the bonding of the allyl moiety.<sup>2</sup> When the allyl group is substituted at one of its terminal positions, the symmetry of its bond to palladium is distorted. QSAR studies<sup>1</sup> have shown that the palladium-allyl bond is distorted from  $\eta^3$  to  $\eta^1$ - $\eta^2$  (Scheme 20). The Pd-C<sub>1</sub> bond is shorter than the Pd-C<sub>3</sub>, this distortion occurs if the allyl group is substituted at one terminus.


It appears that the malonate nucleophile (Scheme 18) attacks preferentially at the allylic carbon atom with the largest palladium-C distance.<sup>3,4</sup> To avoid this complication we opted for a symmetrically substituted olefin (Scheme 21), this allows the designed ligand to provide the required environment distortion; for example, this could be achieved by altering the bite angle. This is the angle which the ligand makes with the palladium (Figure 5).



**Scheme 20**




**Scheme 21**



**Fig 5**

It has been shown that increasing the size of the bite angle enhances the regioselectivity towards the branched product (Scheme 22).




**Scheme 22**

Considering the Heck reaction (Schemes 10,11) the catalytic cycle reveals many differences from the allylic displacement reaction, here the key step which determines enantioselectivity is the association and insertion into the palladium- $R^1$  bond.

Moreover, here the two pathways shown result in different enantioselectivity. For example, the cationic pathway is regarded as being more stereoselective than the neutral pathway. This is due to the former involving a vacant site on the metal which gives a 16 electron species (Figure 6A below). In this case the integrity of the whole system is preserved, thus maximising the asymmetric induction, which results in better stereoselectivity, while in the neutral pathway dissociation of chiral ligand


(Figure 6B) in order to accommodate alkene insertion, tends to reduce enantioselectivity.



In practice it has been possible to influence which pathway will be followed in a given Heck process either by adding silver salts to the reaction<sup>5</sup> or by adding excess halide anions to reactions using triflate, which results in nucleophilic displacement of triflate anion from the product of oxidative addition.<sup>6</sup>

Finally, the 1,4-conjugate addition catalytic cycle is as shown in Scheme 7 in the introduction (page 14).


Here again the asymmetric induction is achieved in a different manner from both of the above reactions. The assumption here is that the chiral ligands are located at opposite corners of the planar array of the metal atom to minimize unfavourable electronic and steric interactions (Figure 7).<sup>7</sup>



**Fig 7**

For chiral lithium organocuprates, one model of the conjugate addition reaction assumes that the cuprate reagent is a dimer with a planar array of metal atoms,<sup>8</sup> and that the chiral ligands are located at the diagonal corners of the planar array of metal atoms to minimize any unfavourable interaction. The observed enantioselectivities then depend upon a number of factors, including substrate structure, ligand stereochemistry, solvent, and the cuprate composition. For a model of the zinc-mediated reaction, see Scheme 51 page 97.

Our design strategy is shown below in Scheme 23.




**Scheme 23**

Armed with this information we proceeded to design a number of ligands. Each class of ligands is considered in terms of their ligating atoms S,N, P,N, etc. Within each class the synthesis is considered and any new novel approaches are explained.

## 2.1

## S,N LIGAND SYNTHESIS

The first attempt in this class of ligands started from the enantiomerically pure commercially available isoquinoline shown in Scheme 24.



Scheme 24

We were able to synthesize the ligand 8e by first reducing the acid using lithium aluminium hydride to give the alcohol. The alcohol was then refluxed with benzaldehyde or pivalaldehyde to give the oxazolidines 8a and 8b, the oxazolidines were then subjected to reductive cleavage to afford the corresponding amino alcohols 8c and 8d, which were then subjected to nucleophilic displacement by the isopropyl thiol after activation of the hydroxy group as the mesylate, prepared using methanesulfonyl chloride.


Only poor enantioselectivity was achieved when the ligands were tested in the palladium catalysed allylic displacement reaction of 1,3-diphenyl propyl-2-enyl acetate using dimethyl malonate as the nucleophile, as described in Chapter 3. The

**Table 1**  
**Palladium Allylic Alkylation Reaction**

R	Yield %	Ligand	ee %
Ph	78	8e	12
t-butyl	92	8d	23

Ephedrine and pseudoephedrine have been used as chiral auxiliaries in many asymmetric syntheses,<sup>9, 10</sup> and we decided to proceed with the design of our ligand using both as our chiral materials. It is essential when planning a design sequence to minimize the number of steps, in order to increase simplicity and to maximise efficiency.

Thioethers (sulfides) have been widely used as chiral ligands in many catalytic asymmetric reactions, for example for the palladium catalysed allylic displacement,<sup>11</sup> and we decided to attempt the synthesis of this type of ligand. The synthetic route is outlined in Scheme 25.



**Scheme 25**

This route involved formation of oxazolidines (9a-d) by condensing ephedrine or pseudoephedrine with the aldehyde in the presence of PTSA under Dean-Stark conditions to give the oxazolidines in 67-90% yield. Table 2 below shows the data for the oxazolidines.

**Table 2**  
**Oxazolidines from Aldehyde**


Compound Number	$R^1$	$R^2$	$R$	Yield %
9a	Ph	H	Me	86
9b	Ph	H	Ph	65
9c	H	Ph	Me	89
9d	H	Ph	Ph	90

We then proceeded to ring open the oxazolidines (9a-d) to obtain the *N*-substituted amino alcohols (10a-d) (Scheme 26). These are usually synthesized by utilizing the Eschweiler-Clarke approach, whereby primary or secondary amines are condensed with formaldehyde followed by hydride transfer from formic acid to yield the *N*-substituted product. Our approach was to use NaBH<sub>3</sub>CN/TMSCl in acetonitrile and to our delight we were able to ring open a variety of substrates.<sup>12</sup> The resulting ring opened structures (10a-d) are shown in Table 3 below. The detailed mechanism of this ring opening procedure is shown below in Scheme 26.

**Table 3**  
**Ring Opening of Oxazolidines Using TMSCl, NaBH<sub>3</sub>CN**

number	<i>R</i> <sup>1</sup>	<i>R</i> <sup>2</sup>	<i>R</i>	Yield %
10a	Ph	H	Me	98
10b	Ph	H	Ph	96
10c	H	Ph	Me	83
10d	H	Ph	Ph	93

**Ring Opening Using  
TMSCl, NaBH<sub>3</sub>CN**




**Scheme 26**

The ring opening procedure presumably occurs by coordination of the oxygen atom of the oxazolidine to the TMSCl, resulting in iminium ion formation. Hydride ion is then transferred from NaBH<sub>3</sub>CN to yield the intermediate (A). Hydrolysis during work up of intermediate (A) yields the required amino alcohol (10a-d).

Having ring-opened the oxazolidines (9a-d) to obtain the *N*-substituted chiral amino alcohols (10a-d), we then proceeded to synthesize the thioethers (11a-d), by substituting the hydroxy groups with the mesylate and then displacing the mesylate with the thiol as outlined in Scheme 25. The mechanism of formation of the thioethers is outlined in Scheme 27 below. The mechanism involves a double inversion sequence at the reacting carbon atom.<sup>13</sup> This overall result has been proved by our group, by obtaining a single crystal X-ray structure which confirms the retention of configuration of the original amino alcohol.<sup>76b</sup> Table 4 shows the thioethers (11a-d) obtained. When tested in the palladium catalysed allylic

displacement (Chapter 3, page 80-81, tables 16 and 17), using 10 mol % of the ligands and 10 mol % of palladium, the outcome was an improvement in ee as compared to (8d-e).



Scheme 27


**Table 4**  
Chiral Sulfide Ligands from Aldehydes

Ligand	$R^1$	$R^2$	$R$	Yield %
11a	Ph	H	Me	65
11b	Ph	H	Ph	63
11c	H	Ph	Me	62
11d	H	Ph	Ph	69

We considered the possibility of using the same sequence of reactions, but instead of an aldehyde, to condense the chiral amino alcohol with aromatic and bulky aliphatic ketones. In doing so we are entering the realm of the unknown because many groups have tried to form oxazolidines from aromatic and bulky aliphatic ketones, but were not successful.<sup>14</sup>

Our original attempt to form the oxazolidines from aromatic ketones used the procedure outlined in Scheme 10, but instead of PTSA, we used  $\text{MgBr}_2$ ,  $\text{AlCl}_3$  and subsequently even camphor sulfonic acid, but in all cases we did not isolate the required oxazolidines.

We noted that scandium triflate has been used as a mild Lewis acid agent in radical cyclisation reactions,<sup>15</sup> in studies of the intramolecular reactions of *N*-acyliminium ions to synthesize the neuvamine skeleton,<sup>16</sup> and in a high yielding cascade sequence.<sup>17</sup> We decided to use this reagent as outlined in Scheme 28. Due to the importance of the ring closure reaction using ketones, a description of the experimental procedure is given here for clarity. 10 mol % of scandium (iii) triflate was added to a solution of 1.2 eq of the ketone in dry dichloromethane, then 1 eq of the ephedrine or pseudoephedrine was added, then 10 g of 4Å molecular sieves was added and the container was sealed and left standing for a period of 7-10 days. Fresh sieves were added four times during the course of the reaction; this was essential for the success of the reaction. The reaction was followed by carbon-13 NMR spectroscopy for the appearance of a peak at 97-110 ppm. When the required peak had been observed, the molecular sieves were removed by filtration and washed with dry dichloromethane (20 ml), the filtrate was stirred with 10-12g of sodium hydrogen carbonate for 2 hours and then removed by filtration. The filtrate was washed with water, dried over magnesium sulfate, and removal of the solvent yielded the oxazolidines (12a-n).




**Scheme 28**

Table 5 below shows the oxazolidines obtained together with yields and diastereoisomeric ratios. The diastereoisomeric ratios were measured using NMR spectroscopy. The proposed mechanism is shown below in Scheme 29.

**Table 5**  
**Formation of Oxazolidines from Ketones using Sc(III) Triflate**

Compound Number	$R^1$	$R^2$	$R^3$	$R^4$	Ratio of Major/ Min	Yield %
12a	Ph	H	Ph	Me	13:1	76
12b	Ph	H	Ph	Et	5 : 3	65
12c	Ph	H	Furyl	2-Me	2 : 1	62
12d	Ph	H	Pyridyl	2-Me	3 : 2	61
12e	Ph	H	p-OMe Ph	Me	6 : 5	63
12f	Ph	H	Et	Me	7 : 1	77
12g	Ph	H	$P_r^i$	Me	16:1	81
12h	H	Ph	Ph	Me	15:1	74
12i	H	Ph	Ph	Et	7:1	73
12j	H	Ph	Furyl	2-Me	4 : 1	72
12k	H	Ph	Pyridyl	2-Me	7 : 1	74
12l	H	Ph	p-OMe Ph	Me	9 : 1	71
12m	H	Ph	Et	Me	12:1	88
12n	H	Ph	$P_r^i$	Me	only one observed	83



Scheme 29

We believe the reaction to be a thermodynamically controlled one. The diastereoselectivity of a thermodynamic controlled reaction is generally dictated by the difference in energy between the products. One will therefore expect an influence of external conditions on the equilibrium states. Indeed it has been documented that the equilibrium shows a dependency on concentration. Furthermore, Cope and Hancock<sup>18</sup> have shown that the product from ethanolamine and dipropyl ketone changes its refractive index upon standing for two months, the phenomenon is reversed by distillation of the product. Metzger<sup>19</sup> showed that when 2-ethyl-2-hexenal and 2-amino-3-methyl-3-butanol were condensed at elevated temperature,

there is a tendency for the Schiff base to be formed and no oxazolidines were observed.

The presence of a Lewis acid such as scandium triflate increases the electrophilicity of the functionality to which it is complexed. In our system we believe that the Lewis acid complexes to the oxygen in the ketone. The diastereoselectivity observed relates to the structure of the substrate. The formation of one diastereoisomer is disfavoured because of the orientation of the large proximal group which raises the energy. The presence of scandium (iii) triflate seems to influence the process by favouring the equilibration between isomers.

Having synthesized the oxazolidines, we needed to know the stereochemistry at C2, and decided to carry out nuclear Overhauser enhancement (nOe) studies on oxazolidines derived from each series of ephedrine and pseudoephedrine. These are shown in Appendix 1. On the basis of the nOe data we were able to confirm the C2 configuration. These are shown in Figure 13.





Fig 13


These results are of considerable interest. The pseudoephedrine configuration at C2 is as expected, that is, the large group is on the same side as the smaller ring methyl group. The result from the ephedrine series, where the large group (L) and the ring

phenyl are on the same side, is surprising. However, if we draw the envelope conformation as shown in Figure 14, in this case the large group (L) and the phenyl are both in pseudoequatorial positions and in doing so they minimize their interaction and result in the lowest energy conformation .



**Fig 14**

Having finally obtained the oxazolidines, we proceeded to investigate their ring opening using our method with  $\text{NaBH}_3\text{CN/TMSCl}$ .<sup>12</sup> However, to our disappointment the method did not work. This may be because the reductive cleavage using this sequence must proceed through the iminium ion intermediate, Figure 15.




**Fig 15**

When  $\text{R}^1$  is a methyl and  $\text{R}^2$  is a hydrogen, or  $\text{R}^1$  is a methyl and  $\text{R}^2$  is a methyl, the reaction is successful, as is also the case when  $\text{R}^1$  is a phenyl and  $\text{R}^2$  is hydrogen. However, when  $\text{R}^1$  is a phenyl and  $\text{R}^2$  is a methyl, the reaction does not work. The explanation may be that in the three former cases the iminium ions are less stable


than in the latter case, because in the latter case where phenyl and methyl substituents are present, Figure 15B, these will stabilize the iminium ion to such an extent that the sodium cyanoborohydride is not able to induce reduction of the iminium intermediate. There is no precedent for reductive ring opening of this type of compound derived from ketones.

An alternative approach was thus needed to ring open this new class of oxazolidines.

We first investigated pyridine borane complex, however, this was unsuccessful. A recent report<sup>20</sup> has described the use of dibutylchlorotin - HMPA complexes, but again with the present substrates the reaction did not give the desired product. Polymethyl hydrosiloxane<sup>21</sup> has been used to reduce ketones but it also did not work in our case. More recently the reductive cleavage of C-N bond in cyclic amidines using DIBAL has been reported,<sup>22</sup> we therefore decided to use this reagent in hexanes or toluene solutions, and to our delight we were able to reduce a range of chiral oxazolidines to obtain a wide range of  $\alpha$ -amino alcohols with high selectivities shown in Table 6 (Scheme 31). For the case of oxazolidine (12n) the reductive ring opening resulted in the amino alcohol (16n) Scheme 30



**Scheme 30**



Scheme 31


Table 6

Formation of aminoalcohols from Oxazolidines using DiBAL


Compound Number	$R^1$	$R^2$	$R^3$	$R^4$	Ratio of Major /Min	Yield %
16a	Ph	H	Ph	Me	13: 1	85
16b	Ph	H	Ph	Et	5: 3	83
16c	Ph	H	Furyl	2-Me	7: 1	87
16d	Ph	H	Pyridyl	2-Me	3: 2	88
16e	Ph	H	p-OMe	Me	6: 5	71
16f	Ph	H	Et	Me	7: 1	79
16g	Ph	H	iPr	Me	6: 1	87
16h	H	Ph	Ph	Me	5: 1	82
16i	H	Ph	Ph	Et	7: 1	86
16j	H	Ph	Pyridyl	2-Me	4: 1	87
16k	H	Ph	Furyl	2-Me	7: 1	83
16l	H	Ph	p-OMe	Me	9: 1	81
16m	H	Ph	Et	Me	12: 1	89
16n	H	Ph	iPr	Me	only one	98

Since the application of DIBAL to the oxazolidines resulted in the required amino alcohols (6a-n), then the pathway may well be different from that using

$\text{NaBH}_3\text{CN/TMSCl}$ . Also, the reaction in DIBAL is successful when the solvent is either hexanes or toluene, but not THF. This may indicate that the oxygen atom of the oxazolidine is coordinating with the aluminium atom, when hexane or toluene is used as solvent. However, when THF is the solvent, the THF coordinates to the aluminium, and the oxazolidine oxygen atom is presumably not sufficiently basic to coordinate with the aluminium and hence the reduction does not occur. This suggests that the coordination to aluminium is an important step in the ring opening process. A possible mechanism is suggested in Scheme 32.




Scheme 32



Scheme 33


Two cases must be considered. In route A Scheme 32, the iminium ion is formed first, then the hydride is delivered. This will result in the formation of two iminium species which will give two different stereoisomeric alcohols. For the case where  $R^3$  and  $R^4$  are methyl and isopropyl the two possible stereoisomers are shown in Scheme 33 above, but we only obtained one. In route B, reductive cleavage occurs by attack of hydride at C2 without formation of iminium ion. This mechanism would be expected to lead to higher stereoselectivities. The two pathways shown in scheme 32 above are likely to take place but we are not sure which is correct. The stereoselectivities obtained using DIBAL are shown in Table 6. These were determined by NMR spectroscopy. These results show that the pseudoephedrine derivatives give higher stereoselectivity than the ephedrine derivatives.

In order to make certain that the amino alcohol had been formed, we decided to synthesize a 3,5-dinitrobenzoate derivative, as shown in Scheme 34 below. We were hoping to obtain an X-ray crystal structure, however to our disappointment, the compound (17) was an oil, Appendix 5 shows the relevant NMR and COSY 45 spectra of this compound.




Scheme 34

Having formed the chiral amino alcohol, we decided to compare our method with the traditional method of ring opening using Grignard reagents (Scheme 35).<sup>26</sup>




The results obtained from these reactions are shown in Table 7 below.

**Table 7**  
**Ring Opening of Oxazolidines**  
**using Grignard Reagent**



**18a-b**



**19a-b**

Entry	Compound Number	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Grignard Reagent	Ratio major/min	Yield (%)
1	18a	Ph	H	Ph	Me	13: 1	76
2	19a	Ph	H	Me	Ph	8: 1	62
3	18b	H	Ph	Ph	Me	3: 1	59
4	19b	H	Ph	Me	Ph	2: 1	69

As mentioned earlier, when using DIBAL to ring open the oxazolidines, the pseudoephedrine series were reduced more selectively. However, as the results in Table 7 show, when Grignard reagents are used to ring open the oxazolidines, the ephedrine derivatives gave the better stereoselectivities. This could be due to the possibility that when oxazolidines are opened with Grignard reagent, they do so through iminium ion formation,<sup>26</sup> while in our proposed mechanism the ring opening using DIBAL proceeds in a concerted fashion without going through the iminium ion. Moreover, when using methylmagnesium iodide as the Grignard reagent, we obtained better selectivity than with phenylmagnesium bromide agents. This is unexpected.

For the case where the oxazolidines were formed from benzaldehyde or acetaldehyde and ephedrine, the oxazolidines formed would be expected to adopt an envelope

conformation, since this results in the lowest energy, with the larger groups pseudoequatorial. The Grignard reagent can then attack either the *si* or *re* faces of the iminium unit (fig 20).




Fig 20

In the case of oxazolidines formed from benzaldehyde, acetaldehyde and pseudoephedrine, attack by the Grignard reagents occurs via iminium ions shown in Figure 21.





Fig 21

Attack by Grignard reagent could again occur at either the *si* or *re* faces of the iminium unit (Table 7).


Having formed the chiral amino alcohols, the next step in the synthesis was to proceed and synthesize the thioethers (Scheme 36).



**Scheme 36**

This was achieved by displacing the hydroxyl group with the mesylate followed by nucleophilic attack by the thiol substrate to yield thioethers (22a-l), Table 8 below. Thioethers obtained through this procedure retain the configuration of the alcohol from which they were derived by a process of double inversion, (Scheme 27).<sup>23</sup>

**Table 8**  
**Chiral Sulfide Ligands derived from Ketones**




Compound Number	<i>R</i> <sup>1</sup>	<i>R</i> <sup>2</sup>	<i>R</i> <sup>3</sup>	<i>R</i> <sup>4</sup>	Yield %
22a	Ph	H	Ph	Me	65
22b	Ph	H	Ph	Et	61
22c	Ph	H	Furyl	2-Me	55
22d	Ph	H	Pyridyl	2-Me	45
22e	Ph	H	iPr	Me	76
22f	Ph	H	Et	Me	78
22g	H	Ph	Ph	Me	71
22h	H	Ph	Ph	Et	74
22i	H	Ph	Furyl	2-Me	61
22j	H	Ph	Pyridyl	2-Me	55
22k	H	Ph	iPr	Me	88
22l	H	Ph	Et	Me	91

On testing these ligands in the palladium catalysed allylic reaction of 1,3 diphenyl propyl-2-enyl acetate, see Chapter 3, an improvement in ee was observed over ligands (8a-b) and (11a-d).

Anderson used imine-derived ligands to achieve some respectable results in the palladium catalysed allylic displacement reaction,<sup>24</sup> and Fiore used pyridine-derived ligands, also to give respectable ees in the same reaction.<sup>25</sup> Therefore we


decided to investigate this approach and to synthesize a number of imine ligands (Scheme 37).



**Scheme 37**


The imines were prepared by refluxing the nitrile, typically 1 eq, with the required Grignard reagent (2.5 eq) in dry ether for 2 h, or until the nitrile had fully reacted. The reaction mixture was then allowed to cool and placed in a cool bath at  $0-15^\circ C$  and 20 eq of dry methanol was added. The solid precipitate was filtered off and washed with ( $3 \times 30$  ml) dry methanol. The methanol was evaporated to dryness to yield the imine ( $23a-h$ ) (Table 9).

**Table 9**  
**Imine from Nitrile**




Compound Number	R	R'	Yield %
23a	Me	Ph	82
23b	Propyl	Ph	85
23c	Et	Ph	89
23d	Bu	Ph	83
23e	Ph	Ph	95
23f	P <sub>i</sub>	Ph	88
23g	1-Naph	1-Naph	72
23h	2-Naph	2-Naph	65


The required imine ligands were obtained by imine exchange. A recent report<sup>27</sup> describes transalkylation using the homogeneous catalysts  $\text{Ru}_3(\text{CO})_{12}$ ,  $\text{Rh}_6(\text{CO})_{16}$  to afford the tertiary amines (Scheme 38). Imine exchange,<sup>28</sup> where Pd black is used as catalyst in the presence of an amine to afford the exchange (Scheme 39) has also been reported. However we opted to use the O'Donnell protocol (Scheme 40),<sup>29</sup> but instead of the Lewis acid ( $\text{TiCl}_4$ ) we used the protic acid camphor sulfonic acid (CSA). A suggested mechanism is shown in scheme 41. The imine exchange was carried out by stirring a mixture of 1:1 of the imine (23a-h) and the amine (Scheme 37) in DCM in the presence of 10 mol % of CSA overnight, and then the crude product was purified to yield the imine ligand (24a-g) in 81-95% yields, (Table 10).




Scheme 38



Scheme 39




Scheme 40




Scheme 41

**Table 10**  
**Sulfur Imine Ligands**



Compound Number	<i>R</i>	<i>R</i> <sup>1</sup>	Yield %
24a	Me	Ph	95
24b	Propyl	Ph	92
24c	Pr <sup>i</sup>	Ph	96
24d	Ph	Ph	96
24f	1-Naph	1-Naph	81
24g	2-Naph	2-Naph	81


On testing these ligands in the Pd catalysed allylic displacement of 1,3-diphenyl propyl-2-enyl acetate, a dramatic improvement of ees was observed of up to 96%. An X-ray structure of one of the imine ligands is shown in Appendix 2 and is represented in Figure 25.



**Fig 25**

For the cases where the imine was not symmetrical, we needed to know the stereochemistry of the ligand and we resorted to the use of nOe spectroscopy.


Appendix 3 shows the results for the case of isopropyl, phenyl, and indicates that for this unsymmetrical imine, the isopropyl group is on the same side as the carbon-nitrogen bond (Figure 26).



**Fig 26**

This stereochemistry may affect the conformations of the palladium complexes with the ligands, and so help to determine enantioselectivity in the palladium catalysed allylic displacement reactions.

From the pattern of ees observed in the palladium catalysed allylic displacement, we noted that substitution on nitrogen improved enantioselectivity, an observation also noted by Feringa,<sup>30</sup> and we decided to explore this effect by choosing ligands with the only structural variation being at nitrogen. Scheme 42 shows the ligands (29a-b) derived from L-phenylalanine.



Scheme 42


The nitrogen group is either Cbz or BOC substituted. Typically, a solution of the Cbz, or BOC substituted L-phenylalanine was stirred in THF, 1 eq of N-methyl morpholine and 1.2 eq of N, O-dimethyl hydroxylamine hydrochloride were added and the solution was stirred for 30 minutes. 1 eq of EDCI was added portion-wise over a 1 hour period, the THF was evaporated and the crude product was purified by chromatography to yield the Weinreb amides (27a-b) in ca 95% yield. The Weinreb amides (27a-b) were reduced to the aldehydes with lithium aluminium hydride at  $0^\circ\text{C}$ . The mixture was quenched with potassium hydrogen sulfate, worked up, and purified by chromatography to give the aldehydes in 96-97% yield.

The aldehydes (28a-b) were treated with propane-1,3-dithiol in the presence


of  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  and the resulting products purified by chromatography yield the ligands (29a-b) in 86-88% yield.

On testing these ligands in the palladium catalysed allylic displacement reaction of 1,3-diphenyl propyl-2-enyl acetate using dimethyl malonate as the nucleophile, a dramatic improvement in enantioselectivity occurred from changing the substituent at nitrogen from Cbz (17% ee) to Boc (44% ee).


So far, we have concentrated on the design of S,N ligands. Figure 30 shows the S,N bidentate ligands which we designed, the uses of which in asymmetric reactions are discussed below,




**11 a-d**  
**R=Me,Ph**




**R<sup>3</sup>=Me**  
**R<sup>4</sup>=Ph,iPr,Furyl,Pyridyl,Et**



**8 d-e**  
**R=Ph,tBut**



**24 a\_f**  
**R<sup>1</sup>=Ph,1-Naph,2-Naph**  
**R=Ph,Me,Et,Prop,iPr**  
**1-Naph,2-Naph**




**29 a-b**  
**R=Cbz,Boc**

**Fig 30**


## P-N LIGANDS

Phosphorus and nitrogen have different  $\pi$ -accepting properties, with phosphorus being the better  $\pi$ -acceptor than nitrogen,<sup>30</sup> thus in the palladium catalysed allylic displacement reaction, nucleophile will attack from the direction *trans* to the better  $\pi$ -acceptor phosphorus atom in the transition state. Also, Morimoto<sup>31</sup> has reported some excellent results in the application of P-N imine ligands in the 1,4 addition of diethylzinc to cyclic and acyclic enones. We decided to design a series of P-N ligating ligands. The starting point was the P-N compound (31c) derived from the phosphine benzaldehyde (31b) and pseudoephedrine (31a) (Scheme 43).



When tested in the palladium catalysed allylic displacement reaction, this ligand gave a respectable enantiomeric excess of 78%. An X-ray crystal structure of this ligand is shown in Appendix 4.

Based on our previous results obtained from the sulfur imine ligands, and from results obtained by Morimoto,<sup>31</sup> we decided to proceed and design phosphorus imine ligands and prepared the three ligands (32a-c) shown in Scheme 44 (Table 11), by condensation of the three aminoesters with 2-(diphenylphosphiny) benzaldehyde.

**Scheme 44**

**Table 11**  
**Phosphorus Imine Ligands**

Compound Number	R	Yield %
32a	iPr	87
32b	Ph	79
32c	(C <sub>6</sub> H <sub>4</sub> OH)CH <sub>2</sub>	65

These ligands did not perform well in the palladium catalysed allylic displacement reaction. However, in the copper-catalysed 1,4 addition of diethylzinc to cyclohexenone (Scheme 45) the ligands gave ees of up to 96%. It is perhaps no surprise that ligands which do very well in one reaction may do badly in others. Thus once a ligand is prepared it pays to test it in more than one field of catalysis.





Figure 33 below shows the P-N ligands which we designed and prepared. The testing

of these ligands is described in the subsequent chapters.



**Fig 33**

The ligands in Figs. 30 and 33 were tested in three types of catalytic asymmetric reactions:

the palladium catalysed allylic displacement using 1,3-diphenylprop-2-enyl acetate and dimethyl malonate as a nucleophile;

the asymmetric copper- catalysed 1,4-addition of diethylzinc to cyclic and acyclic enones;

the asymmetric Heck cross coupling reaction.

## **CHAPTER 3**


# **Testing the Ligands; “The Moment of Truth”**

## TESTING THE LIGANDS; “The MOMENT OF TRUTH”

In this chapter is reported the tests on our ligands which were synthesized as described in the previous chapter. This chapter is divided into three independent sections. Section one deals with the palladium catalysed allylic displacement reaction, section two deals with the 1,4-addition of diethylzinc, and finally section three deals with the asymmetric Heck reaction.

### 3.1 PALLADIUM CATALYSED ALLYLIC DISPLACEMENT


The palladium catalysed displacement reaction represented in Scheme 46 is a classic test of new ligands for asymmetric catalysis.



**Scheme 46**

The reaction is influenced by many factors, such as the solvent, bases, nucleophile, substrate, temperature and time. Thus there is a need to find the best or optimum conditions for each ligand type for a fixed substrate and nucleophile.

To do this we opted to use the ligand (34c) shown in Scheme 47 below, and opted to use dimethylmalonate as the nucleophile and 1,3- diphenylprop-2-enyl acetate as the substrate.



Scheme 47

The ligand was synthesized by reacting 1.2 eq of propane-1,3 -dithiol with 1 eq of the pyridine aldehyde (34a) in the presence of 10 mol % of PTSA under Dean-Stark conditions, to yield the dithiane (34b), this was then oxidized using the Davis oxaziridine to yield the sulfoxide in 88% yield and 98% ee (Scheme 47). The ee was determined using chiral HPLC.

Having prepared this ligand, we decided to use it in the palladium catalysed displacement reaction shown in Scheme 46.

To start with, we varied the solvent and kept the other variables constant. The results are shown in Table 12, and indicate that the use of  $\text{CH}_2\text{Cl}_2$  (DCM) as solvent gave the best enantiomeric excess. This was not surprising since  $\text{CH}_2\text{Cl}_2$  has no coordinating affinity to palladium while the other solvents do, thus those solvents which have affinity to coordinate to palladium will compete with the catalyst, and

since the solvent is used in excess, they may saturate the palladium sites and thus lower ees will result.<sup>32a</sup>

**Table 12**  
**Effect of Solvent in the**  
**Palladium Catalysed Allylic Displacement Reaction**

Solvent	Reaction Temperature	Reaction Time	Yield %	ee %
THF	RT	24 hr	80	35
Et <sub>2</sub> O	RT	24 hr	76	23
DCM	RT	24 hr	95	41
CH <sub>3</sub> CN	RT	24 hr	93	35
DMF	RT	24 hr	72	21

Next we decided to look at the effect of different bases using CH<sub>2</sub>Cl<sub>2</sub> as the solvent and keeping temperature and time constant. The results are shown in Table 13 and indicate that the use of NaH or Cs<sub>2</sub>CO<sub>3</sub> as bases results in the best ees. Although there was not much difference between NaH and Cs<sub>2</sub>CO<sub>3</sub>, we decided on Cs<sub>2</sub>CO<sub>3</sub> due to the handling difficulties associated with the use of NaH.

**Table 13**  
**Effect of Base Palladium Catalysed Allylic Displacement Reaction**

Solvent	Reaction Temperature	Reaction Time	Base	Yield %	ee %
DCM	RT	24 hr	NaH	87	63
DCM	RT	24 hr	BSA	68	32
DCM	RT	24 hr	DBU	75	35
DCM	RT	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	92	56

The effect of temperature on the ees is well known,<sup>32b</sup> so we wished also to investigate the effect of this parameter on the ees in our case. The results are

shown in Table 14 and indicate that in agreement with published work, the effect of temperature on ee is small.<sup>32b</sup>

**Table 14**  
**Effect of Temperatures on Palladium Catalysed Allylic Displacement**

Solvent	Reaction Temperature	Reaction Time	Base	Yield %	ee %
DCM	RT	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	87	46
DCM	0	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	83	55
DCM	-10	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	80	55
DCM	-20	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	74	59
DCM	10	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	79	53
DCM	Reflux	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	65	44

The ees only varied by less than 2% in going from -20 °C to 0 °C.

Finally, we investigated the influence of time on the ee. As shown in Table 15, time has no influence on the ee, but influences the chemical yield.

**Table 15**  
**Effect of Time on the Palladium Catalysed Allylic Displacement**

Solvent	Reaction Temperature	Reaction Time	Base	Yield %	ee %
DCM	RT	6 hr	Cs <sub>2</sub> CO <sub>3</sub>	42	55
DCM	RT	12 hr	Cs <sub>2</sub> CO <sub>3</sub>	61	52
DCM	RT	24 hr	Cs <sub>2</sub> CO <sub>3</sub>	85	53
DCM	RT	36 hr	Cs <sub>2</sub> CO <sub>3</sub>	93	54

Thus from these experiments we were able to fix our conditions listed below.


Temperature	0 °C
Base	Cs <sub>2</sub> CO <sub>3</sub>
Solvent	CH <sub>2</sub> Cl <sub>2</sub> (dichloromethane)
Time	24 hours (complete conversion)
Pd source	allyl palladium chloride dimer
Nucleophile	dimethylmalonate
Substrate	1,3-diphenyl prop-1-enyl acetate

It is important to discuss the stereochemical outcomes of the palladium catalysed allylic displacement reaction. The  $\pi$ -allyl metal complexes exist in dynamic equilibrium under typical displacement reaction conditions (Scheme 48).<sup>33</sup> The choice of ligand (L) may result in a preference for one configuration, due to steric and/or electronic considerations.<sup>34</sup>



**Scheme 48**


The isomeric forms are able to equilibrate by  $\pi$ - $\sigma$ - $\pi$  mechanism (Scheme 49).<sup>1</sup>



**Scheme 49**

Akermark, Vitagliano and co-workers<sup>35,36</sup> demonstrated that the presence of a  $\pi$ -acceptor in the ligand is of fundamental importance for controlling selectivity, and enhancing reactivity. The  $\pi$ -acceptor properties may be thought of as withdrawing electron density from the metal, which in turn increases the positive charge character of the allyl unit, rendering it more susceptible to nucleophilic attack. Furthermore, it has been observed that  $\pi$ -acceptor properties of the ligand are relayed predominantly in a *trans* manner across the complex.<sup>33</sup>

Once we had established those conditions, we proceeded to test our ligands under these conditions using racemic 1,3-diphenyl-1-en-3-yl acetate and dimethyl malonate as nucleophile (Scheme 50).



Scheme 50

Our first ligand to be tested (8e) gave the *S* product in 55% yield with an ee of 23%. The attacking nucleophile is expected to approach the complex *trans* to the sulfur atom, which is the better  $\pi$ -acceptor,<sup>37,38</sup> but may be hindered in its approach by the large group on the nitrogen. Figure 35 shows the two possible conformations of the  $\pi$ -allyl complex obtained. Attack by the nucleophile *trans* to the better  $\pi$ -acceptor in conformation A will give *S* configuration. This conformation presumably allows the minimization of interactions between the phenyl groups of the substrate and the isopropyl and *t*-butyl groups of the complex.<sup>34</sup>





Fig 35

As indicated in our synthetic strategy (Scheme 24) in the previous chapter, our next set of ligands were derived from commercial chiral ephedrine and pseudoephedrine by condensation with aldehydes. The first set of ephedrine derived ligands (11a-b) is shown in Figure 36 below and the corresponding results of the palladium catalysed allylic displacement are shown in Table 16.



R=Me, 11a

R=Ph, 11b

Fig 36

Table 16

## Palladium Catalysed Displacement Reaction

Entry	Compound Number	R	Yield %	ee %	Configuration
1	11a	Me	78	47	S
2	11b	Ph	83	54	S

The results show some improvement in ee compared with the previous ligand (8a). This may be because approach of the nucleophile is not as congested as that for ligand 8a. Again the absolute configuration of the product may be explained by attack of the nucleophile trans to the better p-acceptor, the sulfur atom, at conformation A shown in Figure 37. Conformation A is preferred because Ph<sub>B</sub> avoids the phenyl group of the ligand.





Fig 37

For the pseudoephedrine derivatives (11c-d), Figure 38, results of the palladium catalysed displacement reaction are shown in Table 17 below.



**Table 17**  
**Palladium Catalysed Allylic Displacement**

Entry	Compound Number	R	Yield %	ee %	Configuration
1	11c	Me	85	56	<i>R</i>
2	11d	Ph	88	64	<i>R</i>

As the results indicate, there is some improvement, and the configuration of the major product is now *R*. This change is presumably due to the pseudoephedrine skeleton which has opposite relative configuration to ephedrine at one of the asymmetric centres (the phenyl group). Figure 39 shows the two possible conformations of the allyl moiety. Transition state A may be preferred as it minimizes interaction between Ph<sub>A</sub> and the phenyl group of the pseudoephedrine backbone.





Fig 39

Presumably the higher selectivity obtained by the ligands where R=Ph than the case R=Me is due to the detailed conformation of the reactive transition state.

Our next series of ligands was derived from ephedrine and pseudoephedrine condensed with several ketones.

The ephedrine derivatives (22a-f) (Figure 40) were tested in the palladium catalysed allylic displacement reaction, and the results are shown in Table 18 below.



22 a-f


Fig 40

Table 18

## Palladium Catalysed Allylic Displacement Reaction


Entry	Compound Number	R <sup>1</sup>	R <sup>2</sup>	Yield %	ee %	Configuration
1	22a	Ph	Me	82	52	S
2	22b	Ph	Et	86	44	S
3	22c	Furyl	2-Me	85	25	S
4	22d	pyridyl	2-Me	70	38	S
5	22e	Pri	Me	88	56	S
6	22f	Et	Me	83	54	S

As indicated in Table 18, the product absolute configuration obtained is S, as obtained from ligand (11a-b) also derived from ephedrine, and this confirmed that substituent changes had no effect on the overall transition state conformation. Attack by the nucleophile *trans* to sulfur atom would result in the required S configuration through transition state conformation A shown in Figure 41. Again conformation A may be preferred to B because interaction is avoided between Ph<sub>B</sub> and the ephedrine phenyl group.



**Fig 41**

The corresponding series of ligands derived from pseudoephedrine (22g-l) (Figure 42) were next used in the palladium catalysed allylic displacement, and the results are shown in Table 19



22g-1

Fig 42

Table 19

## Palladium Catalysed Allylic Displacement Reaction

Entry	Compound Number	$R^1$	$R^2$	Yield %	ee %	Configuration
1	10g	Ph	Me	92	68	<i>R</i>
2	10h	Ph	Et	94	61	<i>R</i>
3	10i	Furyl	2-Me	76	45	<i>R</i>
4	10j	pyridyl	2-Me	78	51	<i>R</i>
5	10k	$Pr^i$	Me	93	78	<i>R</i>
6	10l	Et	Me	92	64	<i>R</i>

Table 19 shows that the configuration obtained is *R*, which is again opposite to that of the ephedrine case, the explanation for this outcome is by considering the proposed transition state conformations shown in Figure 43. Again Figure A may be the preferred conformation because  $Ph_A$  avoids the phenyl group of the ligand, while in Figure 43B  $Ph_A$  may interact with the ligand phenyl group.




Fig 43

Although the backbone substituents of the ephedrine and pseudoephedrine skeletons are far away from the Pd atom, they may also exert indirect influence by forcing the substituent on the binding hetroatom to be organized in such away as to minimize steric interactions. In the ephedrine case the backbone substituents are both pointing up as drawn in figure 44A, thus this will force the substituents on sulfur and nitrogen to point down in order to minimize steric interactions. However, for pseudoephedrine the backbone substituents are opposite, the phenyl pointing down while the methyl points up. This may have the effect of forcing the substituent on sulfur to point up while the larger substituent on nitrogen points down (figure 4B); the situation is however less clear.





Fig 44

We next investigated the sulfur imine ligands (24a-f) (Figure 45). These ligands were derived from L-methonine and the corresponding imine by the application of the O'Donnell protocol.<sup>29</sup> The results of the use of these ligands in the palladium catalysed allylic displacement reaction are shown in Table 20 below.



24 a - f

Fig 45

Table 20

## Palladium Catalysed Allylic Displacement Reaction

Entry	Compound Number	$R^1$	$R^2$	Yield %	ee %	Configuration
1	24a	Me	Ph	91	56	<i>R</i>
2	24b	Propyl	Ph	90	53	<i>R</i>
3	24c	$Pr^i$	Ph	93	60	<i>R</i>
4	24d	Ph	Ph	94	96	<i>R</i>
5	24e	1-Naph	1Naph	78	82	<i>R</i>
6	24f	2-Naph	2Naph	78	83	<i>R</i>

One of these ligands gave exceptionally good ees, and the absolute configuration obtained in the product was (*R*) in each case.

The reacting conformation is presumed to be that shown in Figure 46A, because again  $Ph_A$  avoids the ester group; the second conformation (Figure 46B) is disfavoured because  $Ph_A$  would interact with the ester and perhaps with substituents  $R^1$  and  $R^2$ . The imines, as mentioned in Chapter 2, have the *Z* geometry where the groups are different. Attack *trans* to imine in the conformation shown (Figure 46A) would result in the (*R*) configuration of the product.




Fig 46

In this ligand system we believe that the imine group is the better  $\pi$ -acceptor as thioethers are considered to be poorer  $\pi$ -acceptors. Also we believe the unsaturated nitrogen atom may offer greater stabilization to low oxidation state electron-rich metals, such as palladium.<sup>43</sup>

Very good ees, up to 96%, resulted from use of the aromatic imine, ie. R<sup>1</sup>, R<sup>2</sup> = Ph (Figure 47) (entries (4-6), Table 20). However, if one of R<sup>1</sup> or R<sup>2</sup> is aliphatic this resulted in lower ees (entries (1, 2), Table 20), and indeed the longer the linear chain, the lower the ee, because presumably as the linear chain becomes large, steric interactions increase, and binding to palladium weakens. Low ee results.





Fig 47

Naphthyl derivatives (entries (5, 6) Table 20) gave poorer ees than the diphenyl ligand 24d (entry 4), perhaps due to poorer binding to Pd.

So far in the palladium catalysed allylic displacement reaction, we have used N-S ligating ligands (11a-d), (22a-l) and (24a-g). Before starting to consider the phosphorus-nitrogen ligands (31c) and (32a-c) it is worthwhile taking a look at the ligands (29a-b) (Figure 48, Table 21) and to try to offer an explanation for the observed result that when the nitrogen is BOC substituted a better ee results than when it is Cbz substituted.



29a-b R=BOC,Cbz

Fig 48

Table 21  
Palladium Catalysed Allylic Displacement Reaction

Ligand	R	Yield %	ee %	Configuration
29a	BOC	88	44	S
29b	Cbz	52	17	S

The possible conformers are shown in Figure 49.




Fig49

For both cases where R=BOC, Cbz the conformation A is preferred because in this conformation Ph<sub>A</sub> does not interact with the BOC group or Cbz group. Attack of the nucleophile *trans* to sulfur will then result in the observed S configuration of the

product, while in conformation B there is interaction between  $\text{Ph}_A$  and the BOC or Cbz groups.

Also this conformation explains why in the case of BOC substituted derivative (entry 1) a better ee was observed than in the case of the Cbz substituted derivative (entry 2). In the former a larger steric interaction between  $\text{Ph}_A$  and the BOC group than occurs in the latter between  $\text{Ph}_A$  and the Cbz group reinforces the preference for conformation A.

Next is considered the phosphorus-nitrogen ligand (Figure 50, ligand 31c). In the test palladium catalysed allylic displacement reaction this ligand provided a 92% yield and an ee of 78%. The configuration obtained in the product is the R; the possible transition state conformations are indicated in Figure 50. Conformation A must be preferred because it gives rise to the observed enantiomer of the product. It is however difficult to predict the conformation of the ligand-metal complex, particularly around the five-membered ring.





Fig 50

A similar phosphorus oxazolidine ligand has been reported by Svensson .<sup>39</sup>

The next set of phosphorus-nitrogen ligands (32a-c) which were tested are shown in Figure 51, and the results of the palladium catalysed allylic displacement are shown in Table 22.



32 a - c

Fig 51

Table 22

## Palladium Catalysed Allylic Displacement

Compound Number	Entry	R	ee %	Yield %	Configuration
32a	1	Ph	61	82	S
32b	2	Pr <sup>i</sup>	55	75	S
32c	3	p-(OH) C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>	67	82	S

The *S* configuration is obtained by attack of the nucleophile *trans* to phosphorus which is the better  $\pi$ -acceptor.<sup>40</sup> Figure 52 below shows the two possible square planar conformations. Figure 52B must represent the favoured arrangement as it is this transition state which gives rise to the *S* enantiomer of the product. The preferred orientation of the nitrogen substituent cannot easily be predicted. The diagrams shown are not intended to imply any particular conformation of this substituent.




Fig 52

Table 22 shows that the ees are very similar for all of the ligands, perhaps due to the similar sizes of the R group. However, an interesting outcome, entry 3, showed an improved ee when a hydroxyl group (OH) is present. This group is electron-donating and thus may help to transfer electron density via the aromatic ring to the N-atom thus enhancing the binding to palladium.<sup>40,41</sup> Moreover, steric influence of the large  $-\text{CH}_2\text{Ph}(\text{OH})$  unit, may play a part, presumably  $\text{Ph}_A$  avoids this unit, dictating the exact preferred conformation.

The ability of an element to act as good  $\pi$ -acceptor depends on whether the element has a vacant site to accommodate an electron. Second row elements have available to them the d orbital, this is not the case for first row elements. However, first row elements like nitrogen can be made to be good  $\pi$ -acceptors if the hybridization is changed from  $\text{sp}^3$  to  $\text{sp}^2$ , because then the donated electron can be accommodated in  $\pi^*$  orbital. Thus the general accepted hierarchy is  $\text{sp}^2 (\text{N}) > \text{S} > \text{sp}^3 (\text{N})$ .


Having screened our ligands using the palladium -catalysed displacement reaction, we decided to screen some of our ligands in the copper-catalysed conjugate addition of diethylzinc to enones, discussed in section 3.2 below.


For this our first choice was to investigate the phosphorus-nitrogen ligands shown above in Figure 51, since hybrid ligands are often used in this type of reaction.<sup>42</sup>

## 3.2


## 1,4- Conjugate Addition to Enones with Diethyl Zinc

The 1,4- addition to enones is a fundamental process in organic chemistry as mentioned in the introduction. The accepted mechanism for the copper-catalysed 1,4-addition of diethylzinc to 2-cyclohexenone is shown in Scheme 51. The mechanism shows a transition state represented by (A). Recent investigations of this reaction were reported by Feringa<sup>44</sup>, Alexakis,<sup>45</sup> and Schinner.<sup>46</sup> For cyclic enones the reaction is represented in Scheme 52, and for acyclic enones is represented in Scheme 53.






Scheme 52



Scheme 53

As mentioned at the end of Section 3.1 we decided to investigate our P,N phosphorus imine ligands (32a-c) (Figure 53) because others have used P,N hybrid ligands for this reaction.<sup>47</sup> A typical procedure for the catalytic addition reaction is as follows. A solution of  $\text{Cu}(\text{OTf})_2$  and ligand (32a-c) in toluene was stirred under a  $\text{N}_2$  atmosphere at room temperature for 1 hour. The flask was placed in an ice-bath. Diethylzinc and a solution of enone were added sequentially. After stirring for 3 hours the mixture was poured into 1M HCl solution and was extracted with ethyl acetate.



(32a-c)

Fig 53

The organic layer was washed with brine and dried over  $\text{MgSO}_4$ . After the solvent was evaporated the colourless oil obtained was purified by column chromatography (hexane/ethylacetate 9:1). The ees and yields obtained are shown in Tables 23-26 inclusive. The highest ee obtained for cyclopentenone (table 24, entry 3) was a pleasing 88% and this compares well with the published results by Knoble.<sup>48b</sup>

Table 23  
Asymmetric Conjugate Addition of  $\text{Et}_2\text{Zn}$




Entry	R	Enone	Temperature (°C)	Yield %	ee %	$[\alpha]_D$	Configuration
1	Ph		0	79	78	+44	R
2	$\text{Pr}^i$		0	81	58	+33	R
3	$\text{p(OH)C}_6\text{H}_4\text{CH}_2$		0	75	> 99	+58	R

Table 24  
Asymmetric Conjugate Addition of Et<sub>2</sub>Zn




Entry	R	Enone	Temperature (°C)	Yield %	ee %	[α] <sub>D</sub>	Configuration
1	Ph		0	80	66	+51	R
2	Pr <sup>i</sup>		0	65	35	+28	R
3	p-(OH)C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>		0	55	88	+88	R

Table 25  
Asymmetric Conjugate Addition of Et<sub>2</sub>Zn







Entry	R	Enone	Temperature (°C)	Yield %	ee %	[α] <sub>D</sub>	Configuration
1	Ph		0	60	36	+27	R
2	Pr <sup>i</sup>		0	51	17	+13	R
3	p-(OH)C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>		0	78	65	+44	R

Table 26  
Asymmetric Conjugate Addition of Et<sub>2</sub>Zn

Entry	R	Enone	Temperature (°C)	Yield %	ee %	[α] <sub>D</sub>	Configuration
1	Ph		0	55	31	+33	<i>R</i>
2	P <sup>r</sup> i		0	32	9	+11	<i>R</i>
3	p-(OH) C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>		0	82	44	+46	<i>R</i>

In every case the absolute configuration in the product obtained was *R*, suggesting a common transition state stereochemistry. Schinner<sup>46</sup> and Morimoto<sup>48</sup> suggested that a possible intermediate is that shown below in Figure 54A. Using this and the observation of Dieter,<sup>49</sup> Figure 54B (*S*) and (*R*) configurations could be explained. In our case the presence for the (*R*) configuration is due to enone zinc complex avoiding the bulky *R* group and the ester group on the ligand (Figure 54A).





Fig 54

A proposed mechanism for the transfer of alkyl group from  $R_2Zn$  to the  $Cu(II)$  ligand complex is given in Scheme 54. The mechanism is based on observation and kinetic studies by Kraus<sup>54</sup> and Nakamura<sup>55</sup> suggesting that a  $Cu(III)$  species is present. This species could be obtained by electron transfer from the ligand to form a  $Cu(I)$  ligand complex, this then attacks the enone which results in formation

form a Cu(I) ligand complex, this then attacks the enone which results in formation of a Cu(III) ligand enone complex (Scheme 54); this intermediate then attacks the  $R_2Zn$  to result in the transfer of an R group and results in reduction of the Cu(III) ligand complex, to a RCu(I) ligand complex, which on work up yields either the (*R*) or (*S*) enantiomer of the product depending on which face is coordinated, according to Dieter.<sup>49a</sup> The geometry of the copper dimer is believed to be distorted-square-planar.<sup>49b</sup>



**Scheme 54**

In each case, entry 3 (ligand (32c)) gave the best ee in this series. This may be because the presence of (OH) in the para position donates electron density and enhances the binding of the ligand to the metal, moreover the increase in electron density may have a direct effect on the concentration of the enone-cuprate complex.<sup>50</sup>

A lack of this effect may be the cause of the isopropyl derivative ligand (32b) giving low ees in this reaction.

Another effect which requires explanation is the difference in stereoselectivity seen between the cyclohexen-1-one and the cyclopentenone substrates. The much higher ees obtained for cyclohexenone presumably results from its much greater rigidity compared with cyclopentenone.

Having obtained such impressive results, we decided to investigate the effect of temperature on our best ligand (32c) to see if we could improve the yield, while keeping the ee value high. Table 27 shows the results for the conjugate addition to cyclohexen-1-one using our ligand 32c.

**Table 27**  
**Conjugate Addition to Cyclohexen-1-one**  
**with Ligand 32c**

Entry	Temperature ( °C)	Yield %	ee %
1	-10	75	> 99
2	-20	70	> 99
3	-35	65	> 99
4	10	86	81
5	30	88	44

We then repeated the same sequence of reactions to cyclopenten-1-one; the results are shown in Table 28.


**Table 28**  
**Conjugate Addition to Cyclopenten- 1- one**  
**with Ligand 32c**

Entry	Temperature ( °C)	Yield %	ee %
1	-10	50	91
2	-20	48	92
3	-35	47	92
4	10	66	69
5	30	70	48


These results compare very well with those of Degrado<sup>51</sup> who obtained impressive ees of 98% for both the cyclohexen-1-one and cyclopenten-1-one substrates.

These results in Tables (27) and (28) indicate that the yield and ees are divergent quantities, thus for this type of reaction one needs to sacrifice the least important of either ee or yield and in this case the ee is the important entity.

A recent report<sup>52</sup> on the use of thiol ligands in the addition of diethylzinc to aryl aldehydes (Scheme 55) describes impressive ees, and we decided to employ our thioether ligands (22g-l), Figure 55, in the 1,4 addition to cyclohexen-1-one, Scheme 53. Table 29 shows the results obtained.



Scheme 55



22g-1


Fig 55

Table 29  
Asymmetric Conjugate Addition to Cyclohexen-1-one  
using Ligands (22g-1)

Entry	R <sup>1</sup>	R <sup>2</sup>	Yield %	ee %	[α] <sub>D</sub>	Configuration
1	Pr <sup>i</sup>	Me	75	> 99	+65	<i>R</i>
2	furyl	Me	42	66	+49	<i>R</i>
3	pyridyl	Me	62	62	+43	<i>R</i>
4	Ph	Me	66	> 99	+63	<i>R</i>


These results are again very impressive given the fact that thioethers have never been employed as ligands in this type of reaction. A possible transition state model

is shown in Figure 56. The very good ees provided by ligands, 22a, 22k (entries (1) and (4), Table 29), are perhaps due to bulk at nitrogen.<sup>53</sup>



**Fig 56**

In the case of the pyridyl derivative ligand (22i) (entry 3, 62% ee), we were disappointed with the outcome. However, Morimoto reported a similar ee of 52% when his ligand (Figure 57) was used in the asymmetric conjugate addition to cyclohexen-1-one (Scheme 52).<sup>53</sup> The explanation could be that in this case we have two possible transition states shown in Figure 58 below.



**Fig 57**




Fig 58

In the case of transition state model A, the face to face approach of the enone to the copper ligand complex is unhindered and the Cu binds to all three atoms, similarly to Morimoto.<sup>53</sup> Thus this results in stable binding and consequently good ee. The second possibility (Figure 58B), where binding of the sulfur atom is hindered by the t-Bu group, could result in an unfavourable ee. In our case it may be that the complex adopts conformation B (Figure 58).

Further, for the case of the furyl derivatives ligand (22j) (entry 2, 66% ee), again two possible transition state models are possible (Figure 59), each resulting in a different product configuration, similar to the pyridyl derivatives. Here, in the case of 59A, the approach of the enone is unhindered the Cu again binds to all three atoms similar to the Morimoto case and this should result in a product with good ees. However, in Figure 59B the sulfur atom is prevented from binding to the Cu; this may result in an unfavourable ees.




Fig59

Once we were able to screen the thioether ligands, we decided to use the methyl isopropyl ligand (22k) (Figure 60) with other enones, as shown in Table 30.







Fig 60

Table 30  
Conjugate Addition to Enones at 0 °C

Entry	Enone	Configuration	Yield %	ee %
1	Cyclopenten-1-one 	<i>R</i>	72	88
2	Chalcone 	<i>R</i>	46	> 99
3	Benzalacetone 	<i>R</i>	42	> 99

The results were very encouraging because to date no one has yet utilized these types of ligands in this kind of reaction. The absolute configuration of the product obtained was the (*R*) configuration. Again in order to explain the absolute configuration the possible transition state models may be examined. Figure 61A is presumably the

preferred conformation as this will result in the least steric interaction between the enone and the substituent on nitrogen, while Figure 61B will result in interaction between the enone and the substituent on nitrogen. The (*R*) configuration would then result from facial approach as reported by Dieter<sup>49</sup> through transition state model A.




Fig 61

Having obtained these encouraging results, we wondered what effect the catalyst loading would have on the ee, in the conjugate addition (Scheme 53). We varied the ratio of ligand to copper triflate, and we repeated the conjugate addition to cyclohexen-1-one (Table 31), again using our ligand (22k), the methyl isopropyl derivative.

Table 31

**Asymmetric Conjugate Addition to Cyclohexen-1-one  
using Ligand (22k) and Varying the Loading**


Entry	Loading	Time (hours)	Solvent	Temperature (°C)	Yield %	ee %
1	2.5 : 1	4	Toluene	0	82	> 99
2	3 : 1	4	Toluene	0	86	> 99
3	4 : 1	4	Toluene	0	88	> 99
4	4.5 : 1	4	Toluene	0	88	> 99
5	5 : 1	4	Toluene	0	88	> 99
6	5.5 : 1	4	Toluene	0	88	> 99

It can be seen from Table 31 that increasing the loading (ratio of ligand to Cu(OTf)<sub>2</sub>) has no effect on the ee, but does increase the yield. This suggests that there is an optimum ratio of ligand/Cu(OTf)<sub>2</sub> needed, which is around 2.5/1 in this case.

## 3.3


## The Asymmetric Heck Reaction

As is mentioned earlier in the Introduction, there has been considerable interest in the use of the Heck type arylation of olefins for constructing carbon skeletons of biological important organic compounds.<sup>56,57</sup> The asymmetric version, where the carbon-carbon bond formation proceeds with high enantioselectivity, has been demonstrated,<sup>58</sup> where BINAP is used as the chiral ligand. This is represented in Scheme 55 for the reaction of dihydrofuran with phenyl triflate.



Scheme 55

Due to the conflicting reports on the use of base,<sup>58,59</sup> we decided to carry out our own optimization experiment using our phosphorus imine ligand (32b)(Figure 64). The ligand was chosen due to its ready availability.

32b  
Fig 64

Because of time constraints we decided to use the most popular bases as shown in Table 32, and use the procedure shown in Scheme 55 as the test reaction.

**Table 32**  
**Optimization of Asymmetric Heck Reaction**  
**with Ligand 32b**

Entry	Base	Time (hours)	Conversion	Ratio (S) 63 / (R) 62	ee % <sup>a</sup>	
					(S) 63	(R) 62
1	(iPr) <sub>2</sub> NEt	48	100	4 : 1	60	42
2	Et <sub>3</sub> N	48	100	6 : 1	66	47
3	pyridine	48	100	100	-	39
4	proton sponge	48	100	4 : 1	60	46


<sup>a</sup> Absolute configurations were determined by comparison of optical rotations with literature values.

The results in Table 32 indicate that the base of choice is triethylamine. However, other reports contradict our findings and indicate that the use of proton sponge gives better results.<sup>58</sup> A recent report on the use of bases in the asymmetric Heck reaction agrees with our findings and supports triethylamine as the preferred base.<sup>59</sup>

The racemate of each of the products had to be prepared in order to confirm the ee analysis. This was carried out as reported by R Larock.<sup>61</sup> Racemic (63) was prepared by heating together bromobenzene, dihydrofuran, Pd(OAc)<sub>2</sub>, Ph<sub>3</sub>P and silver carbonate in acetonitrile at 80 °C. Removal of the solvent and column chromatography in 70/30 petroleum ether/ ethyl acetate gave the product in 53% yield. Racemic (62) was prepared by stirring together at room temperature bromobenzene, dihydrofuran, Pd(OAc)<sub>2</sub>, Ph<sub>3</sub>P, nBu<sub>4</sub>NCl, and Et<sub>3</sub>N, in benzene. Removal of solvent and column chromatography gave the product in 66% yield.

A striking feature of the asymmetric Heck reaction shown in Scheme 55 is that the two


products (62a) and (63a) have not only opposite absolute configuration, but also a different double bond position. The rationale proposed by many authors<sup>60-64</sup> is shown in Scheme 56; it is hypothesized that the addition of the catalytic complex to either face of the substrate can take place, ultimately producing the complexes (*R*) 56 and (*S*) 56, but in the case of the latter, unfavourable steric factors caused by the ligand induce an immediate dissociation of the Pd species producing the minor product (63) (Scheme 56). However for (*R*) 56, another reinsertion of the alkene into Pd-H bond followed by a second  $\beta$ -hydride elimination occurs to produce the more thermodynamically stable product (62). The overall effect is a kinetic resolution of (*R*) and (*S*)-56, effectively enhancing the facial selectivity in the initial steps of pathways A and B (Scheme 56).



Scheme 56

Ozawa reported that when he used (*R*) BINAP and proton sponge as the base then the favoured product was (*R*) 62 (Scheme 55), and the above explanation gives a credible explanation.


However, in our case the preference was for (63). One explanation for the difference in the product distribution is that the olefin-bond complex (Scheme 57) formed after migratory insertion and  $\beta$ -elimination is more prone to dissociate to give the (*S*) isomer (63) in the phosphorus-nitrogen catalyst system than in the phosphorus-phosphorus system, where as mentioned earlier (Scheme 56) a reverse  $\beta$ -elimination followed by  $\beta$ -elimination and dissociation occurs to give (*R*) (62).



Scheme 57

Having obtained reasonable results we decided to investigate the use of all of our phosphorus imine ligands (32a-c) (Figure 33) in this reaction (Scheme 55) using Et<sub>3</sub>N as base. Table 33 below shows the results, which are similar for all three ligands, giving (63) as the major product in reasonable ees.


**Table 33**  
**Results of Asymmetric Heck Reaction**  
 using Ligands 32a -c



32a\_c

Compound Number	R	Time (h)	Conversion	Ratio (S) 63 / (R) 62	ee %	
					(S) 63	(R) 62
32a	Pr <sup>i</sup>	48	100	5 : 1	70	50
32b	Ph	48	100	60 : 1	66	47
32c	p-(OH)Bn	48	100	6 : 1	67	45

Having obtained resonable results using the dihydrofuran as substrate, we then proceeded to investigate the Heck reaction using the alkene shown in scheme 58 below.




Scheme 58

To our delight we were able to achieve ees of up to 97% and chemical yields of up to 80% using our phosphorus imine ligands (32a-c) (Table 34).

**Table 34**  
**Asymmetric Heck Reaction**  
**using Ligands 32a -c, Scheme 58**

Entry	R	Time (h)	Conversion	Configuration	ee %
1	P <sup>i</sup>	48	100	S	87
2	Ph	48	100	S	92
3	p-(OH) C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>	48	100	S	97


The absolute configuration of the major enantiomer obtained was (*S*). The double bond migration could be explained by looking at Scheme 59. The Pd ligand complex after *syn*-addition to the alkene takes up the conformation A shown; (Scheme 59). This then undergoes *syn* β-hydride elimination to yield product (65) (Scheme 58).<sup>65</sup>



**Scheme 59**

The excellent ees produced by this set of ligands, especially entry (3), may throw some light on the possible factors that govern enantioselectivity. The key step is the

association of the alkene and carbopalladation of the alkene. Since we obtained such high ees, the reaction most probably proceeds by an ionic pathway (Scheme 11) below, because according to Shibasaki this route will lead to high ees since the integrity of the ligand-palladium complex is not violated, while the neutral pathway (Scheme 10) will result in dissociation of the ligand-palladium complex in order to accommodate the alkene, and this generally leads to low ee.<sup>62</sup>




**Scheme 11**

Apart from the mechanistic pathway, other factors which play an important role are the nature of the olefins, with electron-rich olefins favouring the ionic pathway.<sup>63</sup>

Solvents have been varied, with ionic solvents having an improved effect on the enantioselectivity and yield.<sup>64</sup>

Thioethers had not been used as ligands in the asymmetric Heck reaction before the

investigation reported here. We therefore decided to apply our first generation thioether ligand (11d) (Figure 66) to the asymmetric Heck reaction (Scheme 58).



**(11d)**

**Fig 66**


To our delight we obtained an ee of 77% and a yield of 72%. The absolute configuration obtained in the product was (*S*).

## **CHAPTER 4**

### **Substituent effect on the enantioselective oxidation of sulfides to sulfoxides**

As mentioned in the Introduction, chiral sulfoxides are an important class of compounds that find increasing use as chiral auxiliaries in asymmetric synthesis.<sup>66</sup>


There are various approaches to enantiomerically pure sulfoxides; they include the modified Sharpless process reported by Kagan<sup>67</sup> and Modena,<sup>68</sup> the use of Davis oxaziridines,<sup>69</sup> and Page's modified procedure.<sup>70</sup> Our aim in the work described in this chapter was to use the Page modified procedure with the Davis oxaziridine (Figure 67) to investigate the effect of substituent groups in the sulfide substrate on the enantiomeric purity of the products of oxidation in certain classes of sulfide.




**(+)-((8,8-dimethoxycamphoryl) sulfonyl) oxaziridine**

**Fig 67**

To do this we decided to proceed as shown in Scheme 61 to form amino- and nitro-phenyl substituted 1,3-dithianes, and then to carry out the sulfoxidation as shown in Scheme 62



Scheme 61



Scheme 62

The results of the asymmetric sulfoxidation are shown in Table 35.

**Table 35**  
**Sulfoxidation of 4- substituted 2- phenyl- 1,3- dithianes**

Entry	Compound	R	Yield %	ee %	Configuration
1	62a	p-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	87	92	<i>R</i>
2	62b	p -NH <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	92	95	<i>R</i>
3	62c	Ph	98	98	<i>R</i>
4	62d	H	95	36	<i>R</i>


We next investigated the effect of changing the position of the substituent group on the ees. We prepared the *meta*-substitued substrate using similar chemistry (Scheme 61), and proceded to oxidize it as shown in (Scheme 62); the results are shown in table 36.

**Table 36**  
**Sulfoxidation of 2- substituted- 2- phenyl 1,3- dithianes**

Entry	Compound	R	Yield %	ee %	Configuration
1	62e	m -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	92	59	<i>R</i>
2	62f	m- NH <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	94	69	<i>R</i>
3	62c	Ph	98	98	<i>R</i>
4	62d	H	95	36	<i>R</i>

These results agree with the published work of Donnoli, that the introduction of a *meta* substituent eg (Br, NO<sub>2</sub>) group to a phenyl will result in a lowering of ee due

to the inductive influence of this group. However, in Donnoli's case they utilized  $\text{Ti}(\text{O}^i\text{Pr})_4$  and t-butyl hydroperoxide as the oxidant (Scheme 63).




**Scheme 63**

When the substituent groups are in the para position the ees are not affected (Table 35).

Although more work is needed in this area, it is apparent that, when there is no substituent on the aryl group of the 2-aryl-1,3-dithiane, the ee is low. The introduction of a benzene ring at 2-position seems to improve the ees considerably. However, the introduction of a *meta*-nitrogen substituent group on the benzene seems to result in lower ees, but if the substituent is *para* then there is no effect on ee.

As shown in Tables 35 and 36, the absolute configuration of product sulfoxide was (*R*) in each case. Davis has examined the extremes of the possible transition states for the 2-sulfamyloxaziridine shown in figure 68 below, with the common feature being the approach of the sulfur atom lone pair attacking in the plane of the oxaziridine ring. In one case, both lone pairs are coplanar with the three-membered ring, the spiro transition state, while in the other extreme, the plane occupied by the lone pairs of the sulfur atom is perpendicular to that of the oxaziridine ring, the planar transition state. Scheme 64 shows the possible conformations which result in the (*R*) sulfoxide configuration. The Newman


projections show the sulfur atom with its reacting lone pair colinear with the oxygen-nitrogen bond.




**Scheme 64, Newman projections  
as reported by F.A.Davis**

Davis<sup>72</sup> suggested that the most representative of the true transition states are those that show the fewest non-bonded interactions between the Ar' and ZSO<sub>2</sub> group of the oxaziridine and the sulfide (Ar-S-R), thus transition state (a) and (c) (Scheme 64) are favoured while (b) and (d) are disfavoured. Davis also looked at the possible transition states which result in the (S) configuration. These are shown in Scheme 65. He again argued that a presence of the (S) configuration arises from the transition state which shows the least non-bonded interactions.

The favoured transition states are the (a) and (c), while the disfavoured transition states are (b) and (d) (Scheme 65).<sup>72,73</sup> These transition states are again for the oxaziridine shown in figure 68 below.




**Scheme 65, Newman projections as reported by F. A. Davis.**



**Fig 68**

Both the spiro and planar transition states are modelled by the lone pair of sulfur attacking the oxygen of the oxaziridine ring. It is apparent from Schemes 64 and 65 that planar transition states (a, c) have fewer nonbonded interactions than do the spiro transition states (b, d), therefore Davis argued that the planar transition state is favoured. Therefore the transition states (a, c) result in product formation. The preferred diastereomeric transition state for oxidation by (*R, R*) and (*S, S*) oxaziridines is one where the sulfur atom attacks the electrophilic oxygen atom in such a way that Ar' and *R* groups of the substrate Ar'-*S-R* face the small and large regions of the oxaziridine three membered ring. In this case ZSO<sub>2</sub> is considered to be bulkier than C-Ar group.


Using similar argument to the above case the Davis oxaziridine (Figure 67) could be expected to adopt the spiro or planar transition state conformations shown in figure 69 below, again with the planar conformation resulting in the least number of interactions.




The results in Tables 35 and 36 show that there is a considerable improvement of ee when an unsubstituted phenyl group is introduced, compared to when there is no substituent on the 1,3-dithiane moiety; this perhaps is due to the steric influences in the transition state of oxygen transfer, but could also be partly due to electronic effects. However, the presence of a substituent group more electronegative than carbon at the *meta*-position of the ring has the effect of making the sulfur atom less nucleophilic and thus has the effect of reducing the ability of sulfur to donate electrons to the oxaziridine. This effect may contribute towards reducing the enantioselectivity (Table 36).

The introduction of a similar substituent group at the *para* position however, seems to have very little effect on the enantioselectivity, as the results in Tables 35, 36 show. Davis in his work had explained that although electronic effects do play some part in determining the enantioselectivity of the sulfoxidation reaction, the steric influence is the more important.<sup>74</sup>

The possible catalytic cycle for the sulfoxidation by oxaziridine and catalysed by the corresponding imine can be represented as in Scheme 66 below, and the mechanism of sulfoxidation using oxaziridine and  $\text{H}_2\text{O}_2$  is shown in Scheme 67.



**Scheme 66**



Scheme 67

Having considered the effect of some phenylsubstituent groups on the sulfoxidation of 2-phenyl-1,3-dithiane, we wished to investigate different substituents on the oxaziridine. We decided to synthesize the diethoxy derivative shown below in figure 70. The synthesis of this compound is shown in Scheme 68. The synthesis proceeded from camphor sulfonyl chloride and reaction with ammonium hydroxide to yield the sulfonylimine, which was oxidized *in situ* using selenium dioxide to obtain the ketoimine. This was then reacted with triethylorthoformate to form the diethoxy imine which was converted to the oxaziridine derivative with hydrogen peroxide.






Fig70



Scheme 68

We next decided to use the diethoxy oxaziridine (Figure 70) and dimethoxy derivative Figure 67 in the sulfoxidation of sulfide 71 as shown in Scheme 69. The sulfide was synthesized from the 2-chloromethyl pyridine as shown. Using the

diethoxy oxaziridine we obtained an ee of 58% while for the case of dimethoxy oxaziridine an ee of 52% was obtained, thus an improvement of nearly by 10% was achieved using the diethoxy derivative over the dimethoxy derivative. The major product (72), a new compound is believed to have the (R) configuration on the basis of the known results from oxidation of other sulfides with these reagents.<sup>77</sup>



Scheme 69

# **CHAPTER 5**

## **Conclusion**

## CONCLUSION

The aim of the current work was to synthesize chiral hybrid ligands with various ligating heteroatoms including S-N and P-N types, and to examine the possibility of applying these to asymmetric catalysis. Special attention was paid to palladium catalysed allylic displacement reaction, conjugate addition of diethylzinc to enones, and the asymmetric Heck reaction.

Through our journey towards our goal of the synthesis of these ligands we were able to devise methods of synthesis of the various important intermediates, including the oxazolidines derived from aryl ketones, and the very important  $\alpha$ -amino alcohols, achieved by diastereoselective ring-opening using DIBAL in hexanes. From there we were able to synthesize various chiral thioethers which gave respectable ees of up to 78% in the palladium catalysed allylic displacement reaction, and > 99% in the conjugate addition of diethylzinc to cyclic and acyclic enones.

In our quest to improve the enantioselectivity we were able to synthesize a series of sulfur imine ligands which performed extremely well in the palladium catalysed allylic displacement reaction, providing ees of up to 96%.

We then proceeded to the synthesis of phosphorus imine ligands, which were very poor performers in the palladium catalysed allylic displacement reaction, but proved to be very impressive in the conjugate addition of diethylzinc to cyclic

and acyclic enones, with ees up to 99%. Moreover, these ligands showed impressive results in the asymmetric Heck reaction with ees of up to 97%. Overall, we have managed to design and optimize the conditions for all three types of reaction mentioned above, and in all cases we managed to achieve stereoselectivity comparable with the best results in the literature. Further, our ligands are all structurally very simple, and are based on readily available inexpensive materials, and are prepared through extremely simple synthetic routes.

In sulfoxidation, we examined the effects of substituents on the phenyl ring of a 1,3-dithiane system. We were able to show that the presence of *para* nitrogen substituents has no effect on the absolute configuration obtained, and also that the presence of a *meta* nitrogen substituents on the benzene ring has the effect of reducing the enantioselectivity.

We have also utilized our first generation thioether ligands (4a-d) in the asymmetric Heck reaction and obtained an ee of 76%. It is hoped that further work should be carried out in this field since this type of ligand has never been used in this reaction.

It is to be hoped that the present work may contribute in a small way to the understanding of how chiral ligands are incorporated into the transition states of the reactions investigated and how stereoselectivity is achieved thereby.

## EXPERIMENTAL PROCEDURES

Commercially available solvents and reagents were used without further purification. However, low grade solvents were distilled before use. Petroleum ether was distilled from calcium chloride, ethyl acetate and dichloromethane were distilled from calcium hydride. Tetrahydrofuran was distilled from the Na/benzophenone ketyl radical before use.

Air and moisture sensitive reactions were carried out under nitrogen, and all glassware was dried for 24 hours at 150 °C before use.

Column chromatography was carried out using either silica or alumina. Thin layer chromatography was carried out on silica plates. Compounds were visualized using a UV lamp or by permanganate dip.

Melting points were performed on Reichert hot stage apparatus.

Microanalyses were performed on a Perkin-Elmer 2400 Analyser at Loughborough University.

Infrared spectra were recorded in the range 4000-600  $\text{cm}^{-1}$  using a Perkin-Elmer 88 model. Liquids were run neat, solids were run as nujol mulls.

NMR spectra were run on either a Bruker 250 MHz or 400 MHz spectrometer. Tetramethylsilane in deuteriochloroform was used as internal standard. Where needed the chiral shift reagent used to determine ees was (+) europium tris [3, (heptafluoropropyl)hydroxymethylene camphorate], 10 mol % equivalent.

All other enantiomeric excesses were determined by HPLC using a Chiralcel OD column, For Pd catalysed allylic displacement reactions, the eluent used was


99:1 hexane:isopropanol; flow rate 0.5 ml/min, retention times were 30.5 min (*R*) and 33.7 min (*S*). For the asymmetric Heck reaction using 2-phenyl cyclohex-1-ene as alkene, the eluent used was 60:40 hexane:isopropanol; retention times were 7.2min (*S*), 9.4 min (*R*).

For the conjugate addition of diethylzinc, a Perkin Elmer gas chromatograph 8700 was used, using chiral Chrompack CP7502 CP-Chirasil Dex CB 25mm 0.25 mm 1D column. For cyclohexen-1-one, the retention times were 10.5 min (*R*) and 10.6 min (*S*) at isothermal 200 °C. For cyclopenten-1-one, the retention times were 14.5 min (*R*) and 15.2 min (*S*).

Optical rotations were measured on an Optical Activity polAAr-2001 polarimeter at  $\lambda = 589$  nm.

## **EXPERIMENTAL PROCEDURES**


## Oxazolidine Synthesis from Ketones and Pseudoephedrine



To an excess (1.5 eq) of the ketone in DCM (40 ml), scandium (III) triflate (1.34-1.4 g, 27 mmol, 15 mol %) was added, then pseudoephedrine (3.0 g, 18 mmol) was added, followed by 4Å molecular sieves (13 g). The resulting mixture was allowed to stand for two weeks. The mixture was stirred with solid sodium hydrogen carbonate for two hours, then filtered and washed with water (2 × 50 ml). The organic solutions was dried over magnesium sulfate and removal of the solvent yielded the required oxazolidine.

**(4S,5S)-2,3,4-Trimethyl-2,5-diphenyl 1,3-oxazolidine** (1.8 g 74% yield)

[pseudoephedrine(1.5 g 9.0 mmd), acetophenone(1.63 g 13.63 mmd)]




$\nu_{\max}$  (neat) 4379, 3488, 3698, 2309, 1483, 1437, 1360, 1352, 1278, 1230, 1187, 1121  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  (CDCl<sub>3</sub>, 250 MHz) 1.1 (3 H, d,  $J=4.0$  Hz, Me a), 1.6 (3 H, s, Me c), 2.3 (3 H, s, Me b), 2.6-2.8 (1 H, m, Hb), 4.51 (1 H, d,  $J=4.4$  Hz, Ha), 7.2-7.4 (8 H, m), 7.5-7.6 (2 H, m).

$\delta_{\text{C}}$  (CDCl<sub>3</sub>, 100 MHz) 14.26, 20.94, 37.35, 83.96, 96.48, 124.57, 126.14, 127.16, 127.58, 128.03, 128.43, 139.77, 143.86, 145.57.

$m/z$  FAB (Found: 268.1702 ( $M^++1$ ) C<sub>18</sub>H<sub>22</sub>NO requires: 268.1623).  $[\alpha]_{\text{D}} = +67$  (CCl<sub>4</sub>, 10 mg/ml).

**(4S,5S)-2-Pyridine-2-yl-2,3,4-trimethyl-phenyl 1,3-oxazolidine** (1.79 g 74% yield) [(pseudophedrine(1.5 g 9 mmd), 2-acetylpyridine(1.63 g 13.5 mmd)]



**(12k)**

$\nu_{\text{max}}$  (neat) 4350, 3482, 3679, 1304, 1491, 1437, 1356, 1200, 1135, 1126 cm<sup>-1</sup>.


$\delta_{\text{H}}$  (CDCl<sub>3</sub>, 250 MHz) 1.15 (3 H, d,  $J=4.4$  Hz, Me a), 1.83 (3 H, s, Me c), 2.30 (3 H, s, Me b), 2.57-2.60 (1 H, m, Hb), 4.41 (1 H, d,  $J=6.7$  Hz, Ha), 7.1-7.2 (2 H, m), 7.2-7.4 (5 H, m), 7.62-7.80 (2 H, m), 8.52-8.65 (1 H, m).

$\delta_C$  (CDCl<sub>3</sub>, 250 MHz) 16.31, 18.91, 25.32, 35.76, 65.35, 86.21, 99.76, 119.85, 121.76, 125.37, 125.82, 126.76, 127.02, 135.32, 139.46, 149.26, 165.72.

$m/z$  FAB (Found: 269.1656 ( $M^{++1}$ ) C<sub>17</sub>H<sub>21</sub>N<sub>2</sub>O requires: 269.1575).  $[\alpha]_D^{25} = +65$  (CCl<sub>4</sub>, 10 mg/ml).

**(4S,5S)-2-Ethyl-3,4-dimethyl-3,5-diphenyl-1,3-oxazolidine** (1.85 g 79% yield)

[pseudophedrine(1.5 g 9 mmol), propiophenone(1.82 g 13.5 mmol)]



**(12i)**


$\nu_{\max}$  (neat) 4382, 3496, 3703, 2306, 1489, 1436, 1359, 1297, 1198, 1135, 1146  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.9 (3 H, d,  $J=4.4$  Hz, Me a), 1.15 (3 H, t,  $J=5.0$  Hz, Me c), 2.2 (3 H, s, Me b), 2.51-2.53 (1 H, m, Hb), 2.9 (2 H, q,  $J=5.0$  Hz, Hc, Hd), 4.1 (1 H, d,  $J=4.2$  Hz, Ha), 7.15-7.25 (2 H, m), 7.35-7.44 (5H, m), 7.62-7.85 (2 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 8.23, 15.46, 31.75, 37.43, 61.26, 101.23, 126.21, 127.03, 127.96, 128.35, 128.54, 132.85, 136.92, 142.63.

$m/z$  FAB (Found: 282.1878( $\text{M}^++1$ )  $\text{C}_{19}\text{H}_{24}\text{NO}$  requires: 282.1799).  $[\alpha]_{\text{D}} = +45.6$  ( $\text{CCl}_4$ , 10 mg/ml).

(4S,5S) 2,3,4- Trimethyl- 2- (1 methyl ethyl) 5- phenyl]- 1,3- oxazolidine (35 g 83% yield) [pseudophedrine(3.0 g 18.18 mmd), methyl isopropyl ketone(2.3 g 27.27 mmd)]



(12 n)


$\nu_{\max}$  (neat) 43289, 3734, 3130, 2924, 2761, 1459, 1373, 1326, 1189, 1135  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.9 (3 H, d,  $J=5.0$  Hz, Me a), 1.07 (6 H, d,  $J=4.2$  Hz, Me d Me e), 1.12 (3 H, s, Me c), 1.8-1.9 (1 H, m, Hc), 2.1 (3 H, s, Me b), 2.21-2.23 (1 H, m, Hb), 4.21 (1 H, Ha, d,  $J = 5.0$  Hz, Ha), 7.2-7.3 (5 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 7.71, 14.38, 14.56, 33.66, 36.43, 65.14, 85.39, 98.63, 126.22, 126.67, 127.00, 127.74, 140.43.

$m/z$  FAB (Found: 234.1801, ( $\text{M}^++1$ )  $\text{C}_{15}\text{H}_{24}\text{NO}$  requires: 234.1799).  $[\alpha]_{\text{D}} = +39$  ( $\text{CCl}_4$ , 10 mg/ml).

(4S,5S)-2-Furan-2-yl-2,3,4-trimethyl-5-phenyl-1,3-oxazolidine (1.66 g 72% yield) [pseudophedrine (1.5 g 9 mmol), 2-acetyl furan (1.27 g 13.5 mmol)]



$\nu_{\max}$  (neat) 4371, 3454, 2986, 2201, 1481, 1354, 1200, 1175, 1121  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.90 (3 H, d,  $J=4.4$  Hz, Me a), 1.71 (3H, s, Me c), 2.2 (3H, s, Me b) 2.58-2.61 (1H, m, Hb), 4.12 (1 H, d,  $J=4.3$  Hz, Ha), 6.29-6.31 (1 H, m), 7.11-7.21 (1 H, m), 7.31-7.45 (5 H, m), 7.61-7.63 (1 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 250 MHz) 14.41, 25.99, 33.45, 61.29, 77.50, 100.12, 108.50, 112.26, 126.83, 127.83, 127.32, 127.80, 142.01, 146.43, 152.89.

$m/z$  FAB (Found: 258.1499,  $(\text{M}^++1)$   $\text{C}_{16}\text{H}_{20}\text{NO}_2$  requires: 268.1415).  $[\alpha]_{\text{D}} = +37.4$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5S)-2,3,4-Trimethyl-2-(ethyl)-5-phenyl 1,3-oxazolidine (3.8 g 88% yield).**

[pseudophedrine(3.0 g 18.18 mmd), methyl ethyl ketone(1.96 g 27.27 mmd)]



**(12 m)**


$\nu_{\max}$  3827, 3812, 2870, 2787, 1493, 1454, 1370, 1166  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.9 (3 H, d,  $J=7.4$  Hz, Me d), 1.1 (3 H, d,  $J=4.4$  Hz, Me a), 1.3 (3 H, s, Me c), 1.6-1.71 (2 H, m, Hc, Hd.), 2.2 (3 H, s, Me b), 2.6-2.7 (1 H, m, Hb), 4.6 (1 d,  $J=7.6$  Hz, Ha), 7.2-7.3 (5 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 7.70, 114.54, 23.86, 32.18, 32.48, 64.71, 85.71, 96.83, 126.67, 127.76, 127.85, 140.03.

$m/z$  FAB (Found: 220.1625, ( $\text{M}^++1$ )  $\text{C}_{14}\text{H}_{22}\text{NO}$  requires: 220.1623).  $[\alpha]_{\text{D}} = +43$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5S)-2,3,4-Trimethyl-2-(methylphenyl)-5-phenyl 1,3-oxazolidine** (1.9 g 71% yield). [pseudophedrine(1.5 g 9 mmd), para-methoxyacetophenone(2.02 g 13.5 mmd)]




$\nu_{\max}$  4327, 3472, 2206, 1489, 1359, 1198, 1142, 1125  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.9 (3 H, d,  $J=4.4$  Hz, Me a), 1.8 (3 H, s, Me c), 2.2 (3 H, s, Me b), 3.59-3.61 (1 H, m, Hb), 3.9 (3 H, s, Me d), 4.4 (1 H, d,  $J=4.3$  Hz, Ha), 6.9-7.1 (2 H, m), 7.2-7.4 (5 H, m), 8.1-8.2 (2 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 15.12, 26.31, 33.56, 62.39, 77.6, 101.12, 125.31, 26.01, 126.82, 127.31, 127.82, 133.42, 147.35, 153.92.

$m/z$  FAB (Found: 298.2012, ( $\text{M}^++1$ )  $\text{C}_{19}\text{H}_{24}\text{NO}_2$  requires: 298.1728).  $[\alpha]_{\text{D}} = +41.3$  ( $\text{CCl}_4$ , 10 mg/ml).


## Oxazolidine Synthesis from Ketones and Ephedrine



To an excess (1.5 eq) of the ketone in DCM (40 ml), scandium (III) triflate (1.34-1.4 g, 27 mmol, 15 mol %) was added, then ephedrine (3.0 g, 18 mmol) was added, followed by 4Å molecular sieves (13 g). The resulting mixture was allowed to stand for two weeks. The mixture was stirred with solid sodium hydrogen carbonate for two hours, then filtered and washed with water (2 × 50 ml). The organic solutions were dried over magnesium sulfate and removal of the solvent yielded the required oxazolidine.

(4S,5R)-2-3,4,- Trimethyl- 2,5- diphenyl- 1,3- oxazolidine (1.8 g 76% yidd).

[ephedrine(1.0 g 6.0 mmd), acetophenone(1.08 g 9 mmd)]



$\nu_{\max}$  (neat) 4347, 3028, 2314, 1454, 1377, 1265, 1195, 1152  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$ (CDCl<sub>3</sub>, 250 MHz) 0.83 (3 H, d,  $J=4.7$  Hz, Me a), 1.82 (3 H, s, Me c), 2.23 (3 H, s, Me b), 2.59-2.61 (1 H, m, Hb), 4.52 (1 H, d,  $J=4.6$  Hz, Ha), 7.1-7.5 (8 H, m), 8.15-8.21 (2 H, m).

$\delta_{\text{C}}$ (CDCl<sub>3</sub>, 100 MHz) 14.27, 18.96, 33.86, 37.78, 64.21, 81.89, 98.88, 126.15, 126.92, 127.49, , 128.28, , 137.08, 137.11, 141.89.

$m/z$  FAB (Found: 268.1705, ( $M^{+}+1$ ) C<sub>19</sub>H<sub>24</sub>NO<sub>3</sub> requires: 268.1623).  $[\alpha]_{\text{D}} = -4.5$  (CCl<sub>4</sub>, 10 mg/ml).

**(4S,5R)-2-Ethyl-3,4-dimethyl-2,5-diphenyl-1,3-oxazolidine** (3.3 g 65% yield).

[ephedrine(3.0 g 18.18 mmd), propiophenone(3.68 g 27.27 mmd)]



**(12b)**

$\nu_{\text{max}}$  (neat) 4390, 3352, 2977, 1450, 1377, 1350, 1218, 1145, 1137 cm<sup>-1</sup>.


$\delta_{\text{H}}$ (CDCl<sub>3</sub>, 250 MHz) 0.8 (3 H, d,  $J=4.4$  Hz, Me a), 1.03 (3H, t,  $J=4.5$  Hz, Me c), 2.2 (3 H, s, Me b), 2.59-2.61 (1 H, m, Hb), 2.7 (2 H, q,  $J=4.5$  Hz Hc, Hd), 4.5 (1H, d,  $J=4.3$  Hz, Ha), 7.25-7.32 (8 H, m), 7.81-7.83 (2 H, m).

$\delta_{\text{C}}$ (CDCl<sub>3</sub>, 100 MHz) 8.26, 14.19, 31.80, 60.45, 72.92, 96.88, 126.94, 127.07, 128.11, 128.56, 132.87, 136.98, 141.43.

$m/z$  FAB (Found: 282.1793,  $(M^++1)$   $C_{20}H_{24}NO$  requires: 282.1796).  $[\alpha]_D = -7.5$  ( $CCl_4$ , 10 mg/ml).

(4S,5R)-2-Furan-2-yl-2,3,4-trimethyl-5-phenyl 1,3-oxazolidine (2.9 g 62% yield).

[(3.0 g 18.18 mmol), 2-acetylfuran (2.0 g 27.27 mmol)]



$\nu_{max}$  4373, 3393, 2201, 1969, 1392, 1358, 1199, 1137  $cm^{-1}$ .


$\delta_H$  ( $CDCl_3$ , 250 MHz) 0.93 (3 H, d,  $J=4.4$  Hz, Me a), 1.75 (3 H, s, Me c), 2.35 (3 H, s, Me b), 2.59-2.61 (1 H, m, Hb), 4.6 (1 H, d,  $J=4.3$  Hz, Ha), 6.2-6.3 (1 H, m, furan), 7.1-7.2 (1 H, m, Furan), 7.3-7.4 (5 H, m), 7.6-7.7 (1 H, m, Furan).

$\delta_C$  ( $CDCl_3$ , 100 MHz) 14.56, 26.32, 33.67, 62.14, 78.13, 99.96, 107.30, 1121.76, 126.62, 127.93, 127.92, 142.3, 146.53, 153.12.

$m/z$  FAB (Found: 258.1501 ( $M^++1$ )  $C_{17}H_{21}NO_2$  requires: 258.1415).  $[\alpha]_D = -6.9$  ( $CCl_4$ , 10 mg/ml).

**(4S,5S)-2-Pyridine-2-yl-2,3,4-trimethyl-phenyl 1,3-oxazolidine** (2.98 g 61% yield).

[ephedrine(3.0 g 18.18 mmd), 2-acetylpyridine(3.3 g 27.27 mmd)]



**(12d)**

$\nu_{\max}$  3294, 3085, 1585, 1493, 1454, 1358, 1327, 1238, 1196, 1136  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.98 (3 H, d,  $J=4.89$  Hz, Me a), 1.32 (3 H, s, Me c), 2.63 (3 H, s, Me b), 2.65-2.66 (1 H, Hb, m, Hb), 5.22 (1 H, d,  $J=4.8$  Hz, Ha), 7.25-7.91 (8 H, m), 8.52-8.53 (1 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 4.92, 24.96, 33.90, 60.19, 76.89, 82.12, 97.28, 120.65, 121.20, 127.92, 128.09, 128.39, 136.21, 137.01, 140.72, 148.75, 162.97.

$m/z$  FAB (Found: 269.1651( $\text{M}^++1$ )  $\text{C}_{17}\text{H}_{21}\text{N}_2\text{O}$  requires: 269.1575).  $[\alpha]_{\text{D}} = -3.9$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5R)-2,3,4-Trimethyl-2-(methoxyphenyl)-5-phenyl-1,3-oxazolidine** (3.4 g, 63% yield).

[ephedrine(3.0 g, 18.18 mmol), methoxyacetophenone(2.73 g, 36.36 mmol)]



$\nu_{\max}$  4333, 3444, 2970, 1454, 1419, 1358, 1258, 1198, 1145  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.83 (3 H, d,  $J=4.5$  Hz, Me a), 1.92 (3H, s, Me c), 2.33 (3 H, s, Me b), 2.52-2.53 (1 H, m, Hb), 4.64 (1 H, d,  $J=4.7$  Hz, Ha), 7.1-7.2 (2 H, m), 7.3-7.4 (5 H, m), 7.5-7.6 (2 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 14.25, 26.33, 33.70, 55.38, 63.42, 73.37, 88.12, 103.68, 126.18, 126.94, 127.31, 128.06, 128.39, 130.28, 130.56, 142.17.

$m/z$  FAB (Found: 298.1808( $\text{M}^++1$ )  $\text{C}_{19}\text{H}_{24}\text{NO}_2$  requires: 298.1728).  $[\alpha]_{\text{D}} = -8.6$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5R)-2,3,4-Trimethyl-2-(ethyl)-5-phenyl 1,3-oxazolidine** (3.4 g 81% yield).

[ephedrine(3.0 g 18.8 mmd), methyl isopropyl ketone(2.35g 27.27 mmd)]



**(12f)**

$\nu_{\max}$  4390, 3352, 2977, 1450, 1377, 1350, 1218, 1145, 1137  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.8 (3 H, d,  $J=4.4$  Hz, Me a), 1.03 (3 H, t,  $J=4.5$  Hz, Me d), 1.07 (3 H, t,  $J=4.3$  Hz, Me c), 1.72 (2 H, m, Hc, Hd), 2.2 (3 H, s, Me b), 4.92 (1 H, d,  $J=4.3$  Hz, Ha), 7.2-7.3 (5 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 7.68, 14.55, 32.10, 32.44, 60.20, 65.11, 77.55, 96.45, 126.68, 126.74, 127.57, 127.75, 140.48.

$m/z$  FAB (Found: 219.1698( $\text{M}^++1$ )  $\text{C}_{19}\text{H}_{21}\text{NO}$  requires: 219.1623).  $[\alpha]_{\text{D}} = -10.7$ . ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5R) - 2,3,4- Trimethyl- 2- (1- methylethyl) 5- phenyl- 1,3- oxazolidine** (3.43 g 81% yield).

[ephedrine(3.0 g 18.18 mmol), methyl isopropyl ketone(1.34 g 27.27 mmol)]



**(12 g)**


$\nu_{\max}$  4299, 3736, 3261, 2929, 2362, 1460, 1375, 1336, 1198, 1146  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.8 (3 H, d,  $J=4.8$  Hz, Me a), 1.07 (6 H, d,  $J=4.5$  Hz, Me d, Me e), 1.16 (3 H, s, Me c), 2.1-2.2 (1 H, m, Hc), 2.3 (3 H, s, Me b), 2.4-2.5 (1 H, m, Hb), 4.5 (1 H, d,  $J=4.6$  Hz, Ha), 7.2-7.4 (5 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 8.70, 14.55, 16.30, 17.03, 18.22, 32.54, 34.00, 60.12, 80.69, 96.95, 126.47, 127.48, 128.02, 129.57, 140.64.

$m/z$  FAB (Found: 234.1877 ( $\text{M}^++1$ )  $\text{C}_{25}\text{H}_{24}\text{NO}$  requires: 234.1799).  $[\alpha]_{\text{D}} = -6.1$  ( $\text{CCl}_4$ , 10 mg/ml).

## Ring Opening of Oxazolidines using DIBAL/ Hexanes




R<sup>1</sup>=H, R<sup>2</sup>=Ph ephedrine

R<sup>1</sup>=Ph, R<sup>2</sup>=H pseudoephedrine

DiBAL in hexanes (1.5eq) was added slowly to a stirred solution of the oxazolidines (**12a-n**) in hexanes at 0 °C under nitrogen. The resulting solution was stirred at 0 °C overnight. Ethyl acetate (30 ml), methanol (5 ml) and sodium potassium tartarate (10 ml) were added and the resulting mixture was left to stir for two hours. The solid precipitate was filtered off and washed with ethyl acetate (30 ml). The filtrate was dried over magnesium sulphate; removal of the solvent afforded the required product.

**(1R,2S) - 2- ( Methyl ( phenyl- methyl) amino) - 1- phenyl butan- 1- ol**

(0.86 g 85% yield) [5a(1.0 g 3.7 mmol) DIBAL (5.5 ml, 5.5 mmol)]

**16 a**

$\nu_{\text{max}}$  (neat) 3314, 3061, 3028, 2972, 2877, 2801, 2361, 1950, 1602, 1584, 1492  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.86 (3 H, d,  $J=6.4$  Hz, Me a), 1.49 (3 H, d,  $J=6.4$  Hz, Me c), 2.31 (1 H, br, OH), 2.43 (3 H, s, Me b), 2.74 (1 H, dq,  $J=6.4, 4.0$  Hz, Hb), 4.72 (1 H, d,  $J=4.0$  Hz, Ha), 4.82 (1 H, q,  $J=4$  Hz, Hc), 7.26-7.51 (10 H, aromatics).


$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 14.25, 33.61, 46.26, 70.73, 125.19, 126.21, 127.31, 128.15, 128.25, 128.35, 129.15, 137.32, 141.35, 146.26.

$m/z$  EI (Found: 269.1809  $\text{C}_{18}\text{H}_{23}\text{NO}$  requires: 269.1796). Yielded 89%.

$[\alpha]_{\text{D}} = -23.3$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1R,2S) - 2 - ( Methyl ( 1 - phenyl propyl) amino) - 1 - phenyl propane- 1 - ol**

(1.25 g 83% yield) [5b (1.5 g 5.3 mmd) DIBAL (7.95 ml, 7.95 mmd)]

**16b**

$\nu_{\text{max}}$  (neat) 3313, 3061, 3027, 2969, 2874, 1603, 1492, 1451  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=6.48$  Hz, Me a ), 0.85 (3 H, t,  $J=7.4$  Hz, Me c), 1.72-1.73 (2 H, m, He, Hd), 2.41 (3 H, s, He, Hd), 2.72 (1 H, dq,  $J=5.0, 6.48$  Hz, Hb), 4.52 (1 H, d,  $J=5.0$  Hz, Ha), 4.71 (1 H, t,  $J=2.5$  Hz, Hc), 7.26-7.65 (9 H, m), 7.92-7.961 (1 H, m).


$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 1015, 10.83, 30.25, 31.45, 60.25, 70.65, 125.32, 126.15, 127.45, 127.67, 128.35, 128.15, 141.25, 144.35.

$m/z$  EI (Found: 283.1883  $\text{C}_{19}\text{H}_{24}\text{NO}$  requires: 283.1877).

$[\alpha]_{\text{D}} = -8.3$  ( $\text{CCl}_4$ , 10 mg/ml).

(1R,2S)-2-(1-furan-2-ylethyl)(methylamino)-1-phenylpropane-1-ol

(1.85 g 87% yield) [5c(2.1 g 8.17 mmol) DIBAL (12.2 ml, 12.2 mmol)]



16c


$\nu_{\max}$  (neat) 3315, 2977, 1952, 1814, 1658, 1602  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.85 (3 H, d,  $J=6.6$  Hz, Me a), 1.52 (3 H, d,  $J=6.4$  Hz, Me c), 2.42 (3 H, s, Me c), 2.75 (1 H, dq,  $J=4.0, 6.6$  Hz, Hb), 4.72 (1 H, d,  $J=4.0$  Hz, Ha), 4.82 (1 H, q,  $J=6.6$  Hz, Hc), 6.22 (1 H,  $J=3.5$  Hz, Hd), 6.35 (1 H, dd,  $J=1.85, 3$  Hz, He), 7.26-7.45 (5 H, m), 7.42-7.53 (1 H, Hf, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 9.2, 14.63, 33.26, 42.15, 60.27, 73.26, 104.25, 110.26, 126.15, 126.52, 127.56, 128.26, 140.78, 158.26.

$m/z$  EI (Found: 259.1568  $\text{C}_{16}\text{H}_{21}\text{NO}_2$  requires: 259.1572). Yield 80%.  $[\alpha]_{\text{D}} = -3.6$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1R,2S)-2-(Methyl-(1-pyridin-2-yl-methyl)amino-1-phenylpropane-1-ol)**  
 (1.39 g 88% yield) [5d (1.85 g 6.87 mmol), DIBAL (10.3 ml, 10.3 mmol)]



**16d**

$\nu_{\text{max}}$  (neat) 3313, 3062, 2973, 2879, 1952, 1608, 1595  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=6.5$  Hz, Me a), 1.52 (3 H, d,  $J=6.6$  Hz, Me c), 2.43 (3 H, s, Me b), 2.75 (1 H, dq,  $J=4.0, 6.5$  Hz, Hb), 4.92 (1 H, q,  $J=6.6$  Hz, Hc), 7.26-7.46 (5 H, m), 7.22-7.35 (2 H, m), 7.62-7.72 (1 H, m), 8.42-8.51 (1 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 12.74, 16.31, 25.25, 33.56, 60.25, 69.35, 120.25, 127.11, 127.5, 128.31, 136.21, 141.35, 143.42, 148.22, 163.5.

$m/z$  EI (Found: 270.1703  $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}$  requires: 270.1723).

$[\alpha]_{\text{D}} = -6.4$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1R,2S)-2-(Methyl-(1-(4-methoxyphenyl)-methyl)amino-1-phenylpropan-1-ol) (16e)** (1.25 g 83% yield) [5b (1.5 g 5.3 mmol) DIBAL (7.95 ml, 7.95 mmol)]



$\nu_{\max}$  (thin film) 3065, 3067, 2988, 2978, 1921, 1604, 1573  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=6.4$  Hz, Me a), 1.52 (3 H, d,  $J=6.4$  Hz, Me c), 2.43 (3 H, s, Me b), 2.62 (1 H, br, OH), 2.74 (1 H, dq,  $J=3.2, 6.4$  Hz, Hb), 3.87 (3 H, s, Me d), 4.82 (1 H, d,  $J=3.0$  Hz, Ha), 4.91 (1 H, q,  $J=6.4$  Hz, Hc), 6.91-7.26 (4 H, m), 7.25-7.43 (5H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 13.36, 25.26, 33.16, 55.27, 60.20, 72.15, 113.26, 125.62, 125.92, 126.16, 128.83, , 128.35, , 128.75, 141.26, 158.36.

$m/z$  EI (Found: 299.1803  $\text{C}_{19}\text{H}_{25}\text{NO}_2$  requires: 299.1728).

$[\alpha]_{\text{D}}=8.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1R,2S)-2-(1,2-Dimethyl propyl)-methyl-amino-1-phenyl propane-1-ol**  
 (3.72 g 82% yield) [5f (2.5 g 10.73 mmd) DIBAL (16 ml, 16.01 mmd)]



**16f**

$\nu_{\max}$  (thin film) 3316, 3078, 2965, 2878, 1947, 1612, 1585  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=8.0$  Hz, Me a), 0.91 (3 H, d,  $J=8$  Hz, Me c), 0.98 (6 H, 2d,  $J=12, 8$  Hz, Me d, Me e), 1.6 (1 H, Hd, m, Hd), 2.15 (3 H, s, Me b), 2.45 (1 H, dq,  $J=12, 4$  Hz, Hb), 3.5 (1H, br, OH), 3.7 (1 H, d,  $J=4.0$  Hz, Ha), 7.2-7.4 (5 H, m).


$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 11.04, 11.91, 14.50, 20.50, 21.00, 31.94, 53.43, 60.62, 63.47, 128.74, 125.94, 126.10, 126.94, 128.01, 128.13, 141.38.

$m/z$  FAB (Found: 236.1937 ( $\text{M}^++1$ )  $\text{C}_{15}\text{H}_{26}\text{NO}$  requires: 236.1360).

$[\alpha]_{\text{D}} = -12.5$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1R,2S)-2-(1,2-Methyl ethyl)-methyl-amino-1-phenyl propane-1-ol**

(2.6 g 86% yield) [5g(3.0 g 13.7 mmd) DIBAL (20.5 ml, 20.5 mmd)]



**16g**

$\nu_{\max}$  (thin film) 3095, 2987, 2862, 1602, 1542, 1419, 1360  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.86 (3 H, d,  $J=8.0$  Hz, Me a), 0.91 (3 H, t,  $J=4.0$  Hz, Me d), 0.94-0.96 (2 H, m, Ha Hc), 1.01 (3 H,  $J=12$  Hz, Me c), 2.15 (3 H, s, Me b), 2.65 (1 H, dq,  $J=4.0, 4.7$  Hz, Hb), 4.16 (1 H, d,  $J=4.0$  Hz, Ha), 5.12 (1 H, br, OH), 7.26-7.45 (5 H, m).

$m/z$  FAB (Found: 222.1874 ( $\text{M}^++1$ )  $\text{C}_{14}\text{H}_{24}\text{NO}$  requires: 222.1799).

$[\alpha]_{\text{D}} = -7.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1S,2S) - 2- (Methyl- (phenyl- methyl) amino) - 1- phenyl butan- 1- ol**

(1.3 g 82% yield) [5b (1.6 g 5.9 mmol) DIBAL (9 ml, 8.9 mmol)]



$\nu_{\max}$  (neat) 3402, 3091, 3030, 2982, 2890, 2798, 1960, 1605, 1588, 1497  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=6.5$  Hz, Me a), 1.35 (3 H, s, Me c), 2.43 (3 H, s, Me b), 2.61 (1 H, dq,  $J=6.4, 8.2$  Hz, Hb), 2.91 (1H, br, OH), 4.15 (1 H, d,  $J=8.2$  Hz, Ha), 4.52 (1 H, t,  $J=6.0$  Hz, Hc), 7.25-7.52 (6 H, m), 7.62-7.66 (4 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.26, 33.26, 61.15, 68.26, 77.36, 94.36, 125.35, 126.36, 126.76, 128.15, 128.73, 129.36, 139.67, 142.36.

$m/z$  EI (Found: 269.1802  $\text{C}_{18}\text{H}_{23}\text{NO}$  requires: 269.1796).

$[\alpha]_{\text{D}} = +67$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1S,2S)-2-(Methyl-(1-phenyl-propyl)amino)-1-phenylpropane-1-ol**

(1.38 g 86% yield) [5i (1.6 g 5.69 mmol) DIBAL (8.5 ml, 8.5 mmol)]

**16i**

$\nu_{\text{max}}$  (neat) 3313, 3061, 2970, 2874, 1950, 1603, 1495, 1398  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=6.5$  Hz, Me a), 1.35 (3 H, t,  $J=4.5$  Hz, Me c), 1.78-1.92 (2 H, Hd, He, m, Hd, He), 2.43 (3 H, s, Me b), 2.61 (1 H, dq,  $J=6.5, 8.2$  Hz, Hb), 2.91 (1 H, br, OH), 4.15 (1 H, d,  $J=8.2$  Hz, Ha), 4.52 (1 H, t,  $J=6.0$  Hz, Hc), 7.25-7.51 (6 H, m), 7.62-7.66 (4 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 10.26, 15.26, 31.15, 33.35, 55.67, 61.27, 75.26, 127.66, 127.52, 127.75, 128.36, 128.65, 129.34, 142.61, 148.26.

$m/z$  EI (Found: 283.2008  $\text{C}_{19}\text{H}_{25}\text{NO}$  requires: 283.1936).

$[\alpha]_{\text{D}} = +71$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1S,2S)-2-(1-Furan-2-yl-ethyl)-methylamino-1-phenylpropane-1-ol**

(1.34 g 86% yield) [5j (1.60 g 6.2 mmol) DIBAL (9.5 ml, 9.4 mmol)]



$\nu_{\max}$  (thin film) 3317, 2979, 2285, 1952, 1603, 1584, 1492, 1395, 1319  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, d,  $J=6.5$  Hz, Me a), 1.51 (3 H, d,  $J=6.45$  Hz, Me c), 2.42 (3 H, s, Me b), 2.62 (1 H, dq,  $J=6.5, 8.2$  Hz, Hb), 3.01 (1 H, br, OH), 4.12 (1 H, d,  $J=8.2$  Hz, Ha), 4.91 (1 H, q,  $J=6.45$  Hz, Hc), 6.12 (1 H, d,  $J=3.2$  Hz, Hd), 6.22-6.23 (1 H, m, He), 7.25-7.51 (6 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.25, 20.35, 33.62, 61.50, 63.42, 77.51, 105.25, 110.35, 127.15, 127.8, 18.35, 141.25, 142, 86, 158.65.

$m/z$  EI (Found: 259.1653  $\text{C}_{16}\text{H}_{21}\text{NO}_2$  requires: 259.1572).

$[\alpha]_{\text{D}} = +74$  ( $\text{CCl}_4$ , 10 mg/ml).

(1S,2S)-2-(Methyl-(1-pyridine-2-yl-methyl)amino)-1-phenylpropane-1-ol  
(1.4 g 87% yield) [5k (1.6 g 5.97 mmol) DIBAL (9 ml, 8.95 mmol)]



16k

$\nu_{\text{max}}$  (neat) 3315, 3078, 3288, 1955, 1876, 1610, 1597, 1487, 1391  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.82 (3 H, Me1, d,  $J=4.8$  Hz), 1.22 (3 H, Me3, d,  $J=6.4$  Hz), 2.43 (3 H, Me2, s), 2.73 (1 H, Hb, dq,  $J=4.00, 4.80$  Hz), 3.01 (1 H, br, OH), 4.31 (1 H, Ha, d,  $J=4.9$  Hz), 4.85 (1 H, Hc, q,  $J=6.4$  Hz), 7.26-7.62 (5 H, m), 7.76-8.0 (2 H, m), 8.15-8.32 (2 H, m).

$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 16.75, 17.34, 25.55, 60.37, 69.42, 74.26, 120.42, 127.10, 127.50, 128.00, 128.30, 136.75, 141.45, 148.60, 163.75.

$m/z$  EI (Found: 270.1723  $\text{C}_{17}\text{H}_{22}\text{NO}$  requires: 270.1732).

$[\alpha]_{\text{D}} = +64$  ( $\text{CCl}_4$ , 10 mg/ml).

**v (1S,2S) - 2- ( 1,2- Dimethyl propyl) ( methyl amino) - 1- phenyl propane- 1- ol**  
 (3.15 g 98% yield) [5m (3.2 g 13.79 mmd) DIBAL (21 ml, 20.6 mmd)]



16 m

$\nu_{\max}$  3389, 3256, 2878, 2647, 1645, 1542, 1396, 1327  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.85 (3 H, d,  $J=7.2$  Hz, Me a), 0.92 (3 H, Me d, t,  $J=8$  Hz, Me d), 0.93 (2 H, m, Hd, Hc), 0.96 (3 H, d,  $J=6.5$  Hz, Me c), 1.2-1.21 (1 H, m, Hc), 2.15 (3 H, s, Me b), 2.65 (1 H, dq,  $J=4.7, 4.0$  Hz, Hb), 4.16 (1 H, d,  $J=4.0$  Hz, Ha), 5.12 (1 H, br, OH), 7.28-7.45 (5 H, m).


$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 11.22, 18.47, 27.63, 27.86, 59.80, 61.54, 76.65, 125.91, 126.11, 127.43, 142.67.

$m/z$  FAB (Found: 223.1881(  $\text{M}^++1$ )  $\text{C}_{14}\text{H}_{24}\text{NO}$  requires: 223.1877).

$[\alpha]_{\text{D}} = +58$  ( $\text{CCl}_4$ ,

**(1S,2S) - 2- ( Methyl ( 1- ( 4- methoxyphenyl) amino) 1- phenyl propane- 1- ol**

(1.47 g 81% yield) [5l (1.8 g 6.27 mmd) DIBAL (9.8 ml, 9.4 mmd)]



$\nu_{\max}$  (neat) 3146, 3332, 2879, 1950, 1678, 1600, 1454, 1419, 1359  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$ ( $\text{CDCl}_3$ , 250 MHz) 0.83 (3 H, Me a, d,  $J=6.48$  Hz, Me a), 1.42 (3 H, d,  $J=6.25$  Hz, Me c), 2.43 (3 H, s, Me b), 2.51 (1 H, dq,  $J=6.48$ , 8.3 Hz, Hb), 3.00 (1 H, br, OH), 3.72 (3 H, Me c, s, Me c), 4.12 (1 H, d,  $J=8.3$  Hz, Ha), 4.8 (1 H, Hc, q,  $J=6.25$  Hz), 6.85-7.0 (4 H, m), 7.35-7.56 (5 H, m).


$\delta_{\text{C}}$ ( $\text{CDCl}_3$ , 100 MHz) 16.15, 17.67, 25.25, 35.42, 61.00, 70.25, 115.16, 127.35, 127.70, 127.85, 128.35, 128.56, 139.54, 146.78, 160.35.

$m/z$  EI (Found: 299.1342  $\text{C}_{19}\text{H}_{25}\text{NO}_2$  requires: 299.1455).

$[\alpha]_{\text{D}} = +39$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1S,2S)-2-(1,2-Methyl ethyl) (methyl) amino 1-phenyl propane-1-ol**

(2.7 g, 89% yield) [5n (3.0 g, 13.69 mmol) DIBAL (21 ml, 20.5 mmol)]

**16 n**

$\nu_{\max}$  3335, 3256, 2873, 2470, 1604, 1493, 1453,  $\text{cm}^{-1}$ .


$\delta_{\text{H}}(\text{CDCl}_3, 250 \text{ MHz})$  0.86 (3 H, d,  $J=6.6 \text{ Hz}$ , Me a), 0.92 (3 H, Me c, t,  $J=6.6 \text{ Hz}$ , Me c), 1.02\_1.06 (6 H, 2d,  $J=6.07, 6.6 \text{ Hz}$ , Me d, Me e), 1.72\_1.78 (1H, m, Hd), 2.15 (3 H, s, Me b), 2.34 (1 H, m, Hc), 2.66 (1 H, dq,  $J=6.6, 9.0 \text{ Hz}$ , Hb), 4.15 (1H, d,  $J=9.0 \text{ Hz}$ , Ha), 5.12 (1H, br, OH), 7.28-7.45 (5 H, m).

$\delta_{\text{C}}(\text{CDCl}_3, 100 \text{ MHz})$  11.17, 13.26, 19.16, 21.27, 27.35, 32.62, 65.21, 66.26, 75.35, 127.26, 127.65, 128.15, 142.36.

$m/z$  FAB (Found: 236.1936.1881 ( $\text{M}^++1$ )  $\text{C}_{15}\text{H}_{26}\text{NO}$  requires: 236.1936).

$[\alpha]_{\text{D}} = +86$  ( $\text{CCl}_4$ , 10 mg/ml).


## General Procedure for the Synthesis of Sulfides



To a solution of the amino alcohol 6a-n (1 eq) in DCM (30 ml) at 0 °C, triethylamine (2 eq) was added and the reaction was stirred for 10 minutes, methanesulfonyl chloride (1.2 eq) was added and the resulting mixture was left to stir for 1 hour. Triethylamine (2 eq) was added followed by t-butylthiol (2 eq) and the resulting mixture was stirred overnight. The solvent was evaporated and the residue washed with diethyl ether (3 × 30 ml). The combined ether layers were dried over magnesium sulfate. The solvent was evaporated and the residue chromatographed in 10% ethyl acetate, 90% petrol to yield the title compounds as yellow oils.

**N- {(1R,2S)-2-(1,1-Dimethylethylthio)-1-methyl-2-phenylethyl}-N-(1-methyl-2-phenylethylamine)**

(0.65 g 65% yield) [6a (10.8 g 2.97 mmol), *t*-butylthiol (0.6 ml, 5.94 mmol), triethylamine (1.5 ml, 4 eq)]



$\nu_{\max}$  3597, 2995, 2467, 2312, 1419, 1270, 1216, 1197  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.1 (3 H, d,  $J=7.5$  Hz, Me a), 1.25 (9 H, s, Me d), 2.53 (3 H, d,  $J=8.7$  Hz, Me c), 2.63 (3 H, s, Me b), 3.22 (1 H, dq,  $J=6.5$  Hz, 7.4 Hz, Hb), 4.23 (1 H, d,  $J=7.4$  Hz, Ha), 5.17 (1 H, q,  $J=8.5$  Hz, Hc), 7.16-7.46 (10 H, m, aromatics).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 9.2, 9.89, 15.70, 38.52, 39.87, 46.52, 58.98, 125.67, 127.30, 128.52, 128.79, 129.79, 131.20, 140.36, 142.26.

$m/z$  FAB (Found: 342.2262 ( $\text{M}^++1$ )  $\text{C}_{22}\text{H}_{32}\text{NS}$  requires: 342.2177).  $[\alpha]_{\text{D}} = -33.5$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1R,2S)-2- [(1,1)- Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl -N- methyl N- (1- phenyl propyl amine)}**

( 0.65 g 65% yield) 6b( 0.8 g 2.83 mmd), t- butylthid ( 0.65 ml, 5.66 mmd) , triethylamine (1.50 ml, 4 eq)]



**22 b**

$\nu_{\max}$  3422, 3033, 2935, 2526, 1335, 1227, 1198  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.95 (3 H, d,  $J=6.67$  Hz, Me a), 1.21 (9 H, s, Me d), 1.25 (3 H, t,  $J=7.0$  Hz, Me c), 1.52 (2 H, m, Hd, He), 2.43 (3 H, s, Me b), 2.51 (1 H, t,  $J=7$  Hz, Hc), 2.91-2.92 (1 H, m, Hb), 4.3 (1 H, d,  $J=7.4$  Hz, Ha).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 9.21, 9.81, 15.72, 20.52, 38.64, 39.95, 46.92, 59.22, 125.87, 126.21, 127.22, 127.87, 128.15, 129.06, 129.72, 133.15.

$m/z$  FAB (Found: 356.2423 ( $\text{M}^++1$ )  $\text{C}_{23}\text{H}_{34}\text{NS}$  requires: 356.2333).  $[\alpha]_{\text{D}} = -19.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1R,2S)- 2- [(1,1)- Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl} -N- (1-furyl- 2- ethyl) amine**

(1.0 g 55% yield) [6c (1.5 g 5.8 mmd), t- butylthid (1.3 ml, 11.6 mmd), triethylamine( 13.0 ml, 4 eq)]



$\nu_{\max}$  3379, 3298, 2981, 2507, 1654, 1454, 1346, 1191, 1138  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.21 (3 H, d,  $J=6.4$  Hz, Me c), 1.30 (3 H, d,  $J=6.7$  Hz, Me a), 1.31 (9 H, s, Me d), 2.41 (3 H, s, Me b), 3.92 (1 H, dq,  $J=4.6, 6.4$  Hz, Hb), 4.26 (1 H, q,  $J=6.7$  Hz, Hc), 4.91 (1 H, Ha, d,  $J=4.6$  Hz, Ha), 6.21-6.22 (1H, m, He), 6.23-6.24 (1 H, m, Hd), 7.24-7.26 (6 H, m, 5 ar, 1Hf )


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 7.61, 1435, 18.25, 31.26, 31.71, 39.42, 61.21, 75.26, 105.21, 106.76, 127.35, 127.82, 127.98, 128.62, 142.02, 158.10.

$m/z$  FAB (Found: 332.2054 ( $\text{M}^++1$ )  $\text{C}_{20}\text{H}_{30}\text{NO S}$  requires: 332.2048).

$[\alpha]_{\text{D}} = -13.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1R,2S)-2-[(1,1)-Dimethyl ethyl thio] 1-methyl-2-phenylethyl}-N-(1-pyridin-2-ylethyl amine)**

(0.63 g 45% yield) [6d (1.1 g 4.07 mmol), *t*-butylthiol (0.95 ml, 8.14 mmol), triethylamine (2 ml, 4 eq)]



**22 d**

$\nu_{\text{max}}$  3436, 2987, 2941, 1459, 1331, 1197, 1179, 1139  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.00 (3 H, d,  $J=6.4$  Hz, Me a), 1.32 (9 H, s, Me d), 1.45 (3 H, d,  $J=6.6$  Hz, Me c), 2.42 (3 H, s, Me b), 3.22 (1 H, q,  $J=6.8$  Hz, Hc), 3.51 (1 H, dq,  $J=4.6, 6.4$  Hz, Hb), 4.15 (1 H, d,  $J=4.6$  Hz, Ha), 7.16-7.23 (7 H, m), 7.46-7.48 (1 H, m), 8.42-8.43 (1 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.79, 20.01, 31.33, 31.95, 54.06, 60.21, 64.06, 120.21, 121.97, 125.81, 126.82, 128.46, 129.68, 136.25, 136.79, 149.21, 158.15.

$m/z$  FAB (Found: 343.2203 ( $\text{M}^++1$ )  $\text{C}_{21}\text{H}_{30}\text{N}_2\text{S}$  requires: 343.2129).  $[\alpha]_{\text{D}} = -17.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1R,2S) - [(1,1) - Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl}-N- (1,2- dimethyl propyl- N- methyl amine)**

(2.4 g 76% yield) [6f (2.5 g 10.6 mmd), t- butylthid (2.4 ml, 21.2 mmd), triethylamine(6.0 ml, 4 eq)]



$\nu_{\max}$  3752, 2989, 2309, 1435, 1269, 1196  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.95 (6H, 2d,  $J=6.4, 6.7$  Hz, Me e, Me f), 0.96 (6H, 2 d,  $J=6.7, 8.2$  Hz, Me a, Me c), 1.15 (9 H, s, Me c), 1.83-1.84 (1 H, m, Hc), 2.43-2.45 (1 H, m, Hd), 3.00 (3 H, s, Me b), 3.22 (1 H, dq,  $J=6.4, 6.7$  Hz, Hb), 7.15-7.26 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.75, 29.31, 30.26, 31.35, 41.75, 54.72, 64.75, 125.48, 125.92, 126.95, 127.98, 136.12, 142.31, 165.82.

$m/z$  FAB (Found: 308.2411 ( $\text{M}^++1$ )  $\text{C}_{19}\text{H}_{34}\text{NS}$  requires: 308.2333).  $[\alpha]_{\text{D}} = -24.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {( (1R,2S) - [ (1,1) - Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl)-N- (1-methylethyl) - N- methyl amine**

( 258 g 78% yield) [6g(2.5 g 11.31 mmd), t- butylthid (2.5 ml, 22.32 mmd), triethylamine(6 ml, 4 eq)]



**22 f**

$\nu_{\max}$  3672, 2898, 2401, 1426, 1269, 1201, 1198  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.95 (3 H, d,  $J=7.5$  Hz, Me e), 0.98 (3 H, d,  $J=6.7$  Hz, Me a), 1.0 (3 H, d,  $J=6.4$  Hz, Me a), 1.42 (9 H, s, Me d), 1.52-1.54 (2 H, m, Hd, He), 2.46 (3 H, s, Me b), 3.0 (1 H, dq,  $J=6.7, 7.2$  Hz, Hb), 3.91 (1 H, d,  $J=6.7$  Hz, Ha), 7.15-7.35 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.82, 13.91, 27.35, 30.54, 31.54, 31.62, 45.89, 60.82, 62.46, 126.35, 120.49, 127.38, 127.85, 134.65.

$m/z$  FAB (Found: 294.2253 ( $\text{M}^++1$ )  $\text{C}_{18}\text{H}_{32}\text{NS}$  requires: 294.2255).  $[\alpha]_{\text{D}} = -13.6$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- (1S,2S) - 2- [(1,1) - Dimethyl ethyl thio] 1- methyl- 2- phenylethyl}-N- methyl- N- (1 phenylethyl amine)**

( 0.9 g 71% yield) [6h (1.0 g 3.72 mmd), t- butylthid (1.7 ml, 7.44 mmd)]



**22 g**

$\nu_{\max}$  3651, 2898, 2471, 2309, 1418, 1269, 1211, 1198  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.20 (3 H, d,  $J=7.4$  Hz, Me a), 1.21 (9 H, Me d, s, Me d), 2.53 (3 H, d,  $J=9.3$  Hz, Me c), 2.61 (3 H, s, Me b). 3.22 (1 H, dq,  $J=6.7$ , 7.4 Hz, Hb), 4.51 (1 H, d,  $J=7.4$  Hz, Ha), 4.8 (1 H, q,  $J=9.3$  Hz, Hc), 7.15-7.52 (10 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.93, 9.69, 15.50, 16.69, 31.42, 38.42, 39.74, 46.47, 59.07, 125.72, 127.40, 128.68, 128.77, 129.73, 131.06.

$m/z$  FAB (Found: 342.2258 ( $\text{M}^++1$ )  $\text{C}_{22}\text{H}_{32}\text{NS}$  requires: 342.2177).

$[\alpha]_{\text{D}} = +72$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1S,2S)-2- [(1,1)- Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl}-N-methyl- N- (1 phenyl propyl amine)**

(0.65 g 65% yield) [6i (1.0 g 3.5 mmd), t- butylthid (0.8 ml, 7 mmd), triethylamine(2.0 ml, 4 eq)]



**22 h**

$\nu_{\max}$  3414, 3028, 2898, 2511, 1323, 1219, 1195  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.91 (3 H, d,  $J=6.4$  Hz, Me a), 1.0 (9 H, s, Me d), 1.22 (3 H, d,  $J=7.0$  Hz, Me c), 1.53-1.54 (2 H, m, Hd, He), 2.3 (3 H, s, Me b), 2.52 (1 H, t,  $J=7.0$  Hz, Hc), 3.0-3.05 (1 H, m, Hb), 4.51 (1 H, d,  $J=6.4$  Hz, Ha), 7.0-7.25 (10 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.98, 9.79, 15.60, 20.15, 38.42, 39.84, 46.97, 59.18, 125.92, 126.25, 127.31, 127.92, 128.98, 129.01, 129.62, 133.12.

$m/z$  FAB (Found: 356.2407 ( $\text{M}^++1$ )  $\text{C}_{23}\text{H}_{34}\text{NS}$  requires: 356.2333).

$[\alpha]_{\text{D}} = +66$  ( $\text{CCl}_4$ , 10 mg/ml).

**N-[(1S,2S)-2-[(1,1)-Dimethyl ethyl thio] 1-methyl-2-phenyl ethyl]-N-(1-furyl-2-ethyl) amine**

(0.86 g 61% yield) [6j (1.1 g 4.25 mmol), t-butylthiol (1.00 ml, 8.5 mmol), triethylamine (2.5 ml, 4 eq)]



$\nu_{\max}$  3379, 3050, 2938, 2928, 1350, 1220, 1198  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.20 (3 H, d,  $J=6.4$  Hz, Me a), 1.21 (9 H, s, Me d), 1.92 (3 H, d,  $J=8.0$  Hz, Me c), 2.4 (3 H, s, Me b), 3.21 (1 H, dq,  $J=6.4, 8.0$  Hz, Hb), 4.23 (1 H, q,  $J=8.0$  Hz, Hc), 4.51 (1 H, d,  $J=8.0$  Hz, Ha), 6.22 (1 H, d,  $J=4.0$  Hz, Hd), 6.21-6.22 (1 H, m, He), 7.15 (1 H, d,  $J=4.0$  Hz, Hf), 7.25-7.56 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 7.66, 14.27, 28.21, 31.12, 31.17, 39.39, 60.39, 75.14, 105.01, 106.98, 127.32, 127.76, 127.88, 128.57, 141.78, 157.92.

$m/z$  FAB (Found: 332.2054 ( $\text{M}^++1$ )  $\text{C}_{20}\text{H}_{30}\text{NO S}$  requires: 332.2048).  
 $[\alpha]_{\text{D}} = +82$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1S,2S)-2- [(1,1)- Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl}-N- (1- pyridin- 2- yethyl amine)**

( 0.77 g 55% yield) [6k ( 10.1g 4.07 mmd), t- butylthid ( 1.0 ml, 8.15 mmd), triethylamine (2.1ml, 4 eq)]



**22 j**

$\nu_{\max}$  3425, 2981, 2939, 1454, 1327, 1191, 1176, 1138  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.91 (3 H, d,  $J=6.6$  Hz, Me a), 1.22 (9 H, s, Me d), 1.45 (3 H, d,  $J=6.6$  Hz, Me c), 2.16 (3 H, s, Me b), 3.35 (1 H, q,  $J=6.5$  Hz, Hc), 3.45 (1 H, dq,  $J=4.6, 6.6$  Hz, Hb), 3.91 (1 H, d,  $J=4.6$  Hz, Ha), 7.16-7.28 (5H, m), 7.32-7.33 (2 H, m), 8.35-8.36 (2 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.41, 19.09, 31.24, 31.80, 53.08, 59.35, 63.97, 119.72, 121.81, 125.79, 126.79, 128.36, 136.50, 144.86, 148.37, 148.46, 165.10.

$m/z$  FAB (Found: 343.2213 ( $\text{M}^++1$ )  $\text{C}_{21}\text{H}_{31}\text{N}_2\text{S}$  requires: 343.2129).  
 $[\alpha]_{\text{D}} = +69$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1S,2S) - [(1,1) - Dimethyl ethyl thio]- 1- methyl 2- phenyl ethyl}-N- (1,2 dimethyl, propyl) N- methyl amine**

(02.87 g 88% yield) [6m (2.5 g 10.64 mmd), t- butylthid (2.4 ml, 21.28 mmd), triethylamine (6 ml, 4 eq)]



**22 k**

$\nu_{\max}$  3748, 2984, 2305, 1420, 1265, 1194  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.10 (6 H, 2 d,  $J=6.7$ , 6.4 Hz, Me e, Me f), 1.12 (6H,,2d, $J=7.0,7.6$  Hz, Me a, Me c), 1.12 (9 H, s, Me d), 1.83-1.85 (1 H, m, Hd), 2.42-2.43 (1 H, m, Hc), 3.0 (3 H, s, Me b), 3.12 (1 H, dq,  $J=6.7$ , 6.4 Hz, Hb), 4.2 (1 H, d,  $J=6.4$  Hz, Ha), 7.15-7.36 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.67, 29.31, 30.21, 31.33, 41.67, 54.64, 64.56, 66.02, 125.48, 125.90, 126.94, 127.84,129.56, 143.45.

$m/z$  FAB (Found: 304.2412 ( $\text{M}^++1$ )  $\text{C}_{19}\text{H}_{34}\text{NS}$  requires: 304.2333).  
 $[\alpha]_{\text{D}} = +92$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1S,2S) - [(1,1) - Dimethyl ethyl thio] 1- methyl 2- phenyl ethyl}-N- (1,methylethyl) N- methyl amine**

(2.4 g 91% yield) [6n (2.0 g 9.0 mmd), t- butylthid (2.0 ml, 18.0 mmd), triethylamine (5 ml, 4 eq)]



**22 I**

$\nu_{\max}$  3650, 2897, 2406, 1421, 1267, 1201, 1197  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.9 (3 H, t,  $J=7.15$  Hz, Me e), 0.96 (3 H, d,  $J=6.45$  Hz, Me a), 0.98 (3 H, d,  $J=6.0$  Hz, Me c), 1.42 (9 H, s, Me d), 1.53-1.54 (2 H, m, Hd, He), 2.15 (3 H, s, Me b), 3.0 (1 H, dq, 6.0, 6.7 Hz, Hd), 3.2-3.3 (1 H, d, m, Hc), 3.92 (1 H, d,  $J=6.7$  Hz, Ha), 7.0-7.25 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.72, 13.80, 15.93, 27.35, 31.39, 31.54, 45.86, 60.72, 62.44, 126.27, 126.42, 127.28, 127.72, 134.45.

$m/z$  FAB (Found: 294.2252 ( $\text{M}^++1$ )  $\text{C}_{18}\text{H}_{32}\text{NS}$  requires: 294.2255).  
 $[\alpha]_{\text{D}} = +63$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- (1R,2S)- [(1,1)- Dimethyl ethyl thid] 1- methyl- 2- phenyl ethyl –N- methyl ethyl amine**

(1.3 g 65% yield) [3a( 1.5 g 7.7 mmd), t- butylthid ( 1.74 ml, 15.4 mmd), triethylamine( 4.0 ml, 4 eq)]



**11 a**

$\nu_{\max}$  3698, 2867, 2327, 1445, 1273, 1196  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.99 (3 H, d,  $J=8.0$  Hz Me a), 1.15 (9 H, s, Me d), 1.45 (3 H, t,  $J=7.6$  Hz, Me c), 2.5 (3 H, s, Me b), 3.31 (2 H, q,  $J=8.0$  Hz, Hc , Hd), 3.24-3.26 (1 H, m, Hb), 4.12 (1 H, d,  $J=8.6$  Hz, Ha), 7.15-7.52 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.21, 8.67, 9.21, 29.56, 49.21, 55.26, 67.34, 125.47, 126.42, 127.62, 127.83, 141.29.

$m/z$  FAB (Found: 266.1997 ( $\text{M}^++1$ )  $\text{C}_{16}\text{H}_{28}\text{NS}$  requires: 266.1864).  $[\alpha]_{\text{D}} = -34.7$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- (1R,2S)- (1,1)- Dimethyl ethyl thio 1- methyl- 2- phenyl ethyl -N- 1- phenyl ethyl amine**

(0.74 g 63% yield) [3b (1.2 g 3.6 mmd), t- butylthio (0.8 ml, 7.2 mmd), triethylamine (2 ml, 4 eq)]



$\nu_{\max}$  3666, 2892, 2334, 1440, 1280, 1199  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.19 (9 H, s, Me d), 1.23 (3 H, d,  $J=8.0$  Hz, Me a), 2.2 (3 H, s, Me b), 3.11 (2 H, s, Hc, Hd), 3.49-3.51 (1 H, m, Hb), 3.91 (1 H, d,  $J=8.0$  Hz, Ha), 6.85-7.75 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.40, 30.57, 36.56, 52.65, 58.26, 63.33, 125.98, 126.47, 127.81, 127.86, 128.34, 128.46, 140.01, 145.51.

$m/z$  EI (Found: 328.2017  $\text{C}_{21}\text{H}_{30}\text{NS}$  requires: 328.2014).  $[\alpha]_{\text{D}} = -37$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- {(1S,2S)-[(1,1)-Dimethyl ethyl thio] 1-methyl- 2- phenyl ethyl-N-methyl} ethyl amine**

(0.69 g 62% yield) [3c (1.4 g 4.2 mmd), t-butylthid (0.9 ml, 8.4 mmd), triethylamine (2.5 ml, 4 eq)]



**11 c**

$\nu_{\max}$  3696, 2884, 2331, 1440, 1268, 1194  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.04 (3 H, d,  $J=8.0$  Hz, Me a), 1.24 (9 H, Me d, s, Me d), 1.35 (3 H, t,  $J=8.2$  Hz, Me c), 2.5 (3 H, s, Me b), 3.12 (2 H, q,  $J=8.0$  Hz, Hc, Hd), 3.16-3.18 (1 H, m, Hb), 3.91 (1 H, d,  $J=8.0$  Hz, Ha), 7.15-7.52 (5 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 7.62, 7.67, 8.77, 29.75, 46.51, 51.77, 126.43, 127.49, 128.29, 129.03, 139.29, 143.77.

$m/z$  FAB (Found: 266.1990 ( $M^+ + 1$ )  $\text{C}_{16}\text{H}_{28}\text{NS}$  requires: 266.1864).

$[\alpha]_{\text{D}} = +61$  ( $\text{CCl}_4$ , 10 mg/ml).

**N- (1S,2S) - [(1,1) - Dimethyl ethyl thio] 1- methyl- 2- phenyl ethyl-N- 1- phenyl ethyl amine**

(1.0 g 69% yield) [3d (1.5 g 4.5 mmol), t- butylthiol (1.0 ml, 9 mmol), triethylamine (2.5 ml, 4 eq)]



$\nu_{\max}$  3652, 2897, 2311, 1436, 1270, 1199  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.83 (3 H, d,  $J=6.8$  Hz, Me a), 1.12 (9 H, s, Me d), 2.15 (3 H, s, Me b), 2.43 (2 H, s, Hc, Hd), 3.16 (1 H, dq,  $J=6.8, 9.6$  Hz, Hb), 3.91 (1 H, d,  $J=9.6$  Hz, Ha), 7.15-7.52 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.37, 10.30, 31.20, 32.10, 35.75, 51.68, 63.02, 126.72, 126.82, 127.40, 128.02, 128.08, 128.22, 141.89, 143.77

$m/z$  FAB (Found: 328.2099 ( $\text{M}^++1$ )  $\text{C}_{21}\text{H}_{30}\text{NS}$  requires: 328.2020).

$[\alpha]_{\text{D}} = +63$  ( $\text{CCl}_4$ , 10 mg/ml).


## Synthesis of Oxazolidines from Acetone Ephedrine and Pseudoephedrine



Ephedrine or pseudoephedrine (3.0 g, 18 mmol) was dissolved in acetone (50 ml), 4Å molecular sieve added, and the mixture allowed to stand at room temperature for 3 days. The solvent was evaporated to leave the product.

### (4S,5S) - 2,3,4- Tetramethyl- 5- phenyl- 1,3- oxazolidine

(3.0 g 82% yield) [pseudoephedrine(3.0 g 18.18 mmol)]



$\nu_{\text{max}}$  (neat) 3443, 3030, 2474, 2795, 1645, 1450, 1250  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  (250 MHz  $\text{CDCl}_3$ ) 1.16 (3 H, d,  $J=7.0$  Hz, Me a), 1.33 (6 H, s, Me c, Me d), 2.30 (3 H, s, Me b), 2.60-2.61 (1 H, m, Hb), 4.50 (1 H, d,  $J=7.5$  Hz, Ha).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.25, 21.35, 28.25, 32.15, 62.25, 83.65, 95.16, 126.5, 127.11, 128.15, 140.25.

$m/z$  (EI) (Found: 205.1466  $\text{C}_{13}\text{H}_{19}\text{NO}$  requires: 205.1466).  $[\alpha]_{\text{D}} = +35.6$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4R,2S) - 2,3,4- Tetramethyl- 5- phenyl- 1,3- oxazolidine**

(3.2 g 85% yield) [ephedrine(3.0 g 18.18 mmol)]



(Found: C 75.57, H 9.24, N 6.79,  $C_{13}H_{19}NO$ . requires: C 76.01, H 9.27, N 6.83%)

$\nu_{\max}$  (slurry) 3422, 2976, 2932, 2800, 2796, 1607, 1592, 1492  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  (250 MHz  $\text{CDCl}_3$ ) 0.63 (3 H, d,  $J=5.0$  Hz, Me a), 1.22 (3 H, s, Me c), 2.25 (3 H, Me d, s), 3.60-3.61 (1 H, Hb, m), 5.0 (1 H, Ha, d,  $J=7.0$  Hz), 7.26-7.54 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.25, 19.35, 35.25, 60.35, 81.25, 95.15, 127.35, 127.5, 127.7, 128.3, 140.15.

$m/z$  (EI) (Found: 205.1463  $C_{13}H_{19}NO$  requires: 205.1466).

$[\alpha]_{\text{D}} = +49$  ( $\text{CCl}_4$ , 10 mg/ml).


## Synthesis of Oxazolidines using Aldehydes, Ephedrine or Pseudoephedrine under Dean-Stark Conditions



Ephedrine or pseudoephedrine (3.0 g, 18.0 mmol) was dissolved in toluene (50 ml), then the aldehyde was added (2 eq), followed by CSA (10 mol %). The resulting mixture was heated under Dean-Stark conditions. The solvent was removed, and DCM (30 ml) added, and the solution was washed with aqueous sodium hydrogen carbonate (3 × 20 ml). The organics were dried over magnesium sulfate. Removal of the solvent yielded the title compound.

### (4R,5S)-3,4-Dimethyl-2,5-phenyl-1,3-oxazolidine

(3.5 g 76% yield) [pseudoephedrine(3.0 g 18.18 mmol)]



(Found: C 81.0, H 7.51, N 5.53; C<sub>17</sub>H<sub>19</sub>NO, requires: C 81.17, H 7.65, N 5.45 %)

$\nu_{\max}$  (slurry) 3448, 2968, 2794, 2719, 1605, 1495, 1456, 1191 cm<sup>-1</sup>.

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.81 (3 H, Me1, d,  $J=5.0$  Hz), 2.22 (3 H, Me2, s), 3.0-3.01 (1 H, Hb, m), 4.71 (1 H, Hc, s), 5.22 (1 H, Ha, d,  $J=5.0$  Hz), 7.10-7.62 (10 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.25, 35.15, 65.32, 100.21, 127.75, 128.12, 128.4, 130.0, 130.15, 131.21, 131.45, 132.15, 139.8.

$m/z$  EI (Found: 253.1468  $\text{C}_{17}\text{H}_{19}\text{NO}$  requires: 253.1468).

$[\alpha]_{\text{D}} = -55$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4R,5S)-2,3,4-Trimethyl-5-phenyl-1,3-oxazolidine**

(2.85 g 82% yield) [ephedrine(3.0 g 18.18 mmd)]



**9 a**

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.65 (3 H, d,  $J=7.0$  Hz, Me a), 1.45 (3 H, d,  $J=5.0$  Hz, Me c), 2.25 (3 H, s, Me b), 2.74-2.75 (1 H, m, Hb), 3.91 (1 H, q,  $J=5.0$  Hz, Hc), 5.0 (1 H, d,  $J=6.0$  Hz, Ha), 7.24-7.43 (5 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.19, 36.64, 65.12, 82.25, 94.75, 127.15, 127.40, 127.60, 128.1, 129.35.

$m/z$  EI (Found: 191.1308  $\text{C}_{12}\text{H}_{17}\text{NO}$  requires: 191.1310).

$[\alpha]_{\text{D}} = -21.2$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5S)-2,3,4-Trimethyl-5-phenyl-1,3-oxazolidine**

(1.43 g 62% yield) [pseudophedrine(2.0 g 12.12 mmd) acetaldehyde(1.1 g 24.24 mmd)]

**9 c**

$\nu_{\max}$  3485, 2713, 2803, 1608, 1492, 1362  $\text{cm}^{-1}$ .

$^1\text{H}$  ( $\text{CDCl}_3$ , 250 MHz) 1.15 (3 H, d,  $J=6.4$  Hz, Me a, ), 1.42 (3 H, d,  $J=6.8$  Hz, Me c), 2.24 (3 H, s, Me b), 2.31-2.34 (1 H, m, Hb), 4.22 (1 H, q,  $J=6.9$  Hz, Hc), 4.52 (1 H, d,  $J=6.8$  Hz, Ha), 7.26-7.45 (5 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.25, 19.35, 36.25, 64.51, 82.50, 94.75, 127.15, 128.7, 128.85, 141.25.

$m/z$  EI (Found: 191.1308  $\text{C}_{12}\text{H}_{17}\text{NO}$  requires: 191.1310).

$[\alpha]_{\text{D}} = +22$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5S) - 3,4- Dimethyl- 2,5- di phenyl- 1,3- oxazolidine**

(2.1 g 69% yield) [pseudophedrine(2.0 g 12.12 mmd) benzaldehyde(2.5 g 24.24 mmd)]



(Found: C 80.83, H 7.61, N 5.65. requires: C 80.63, H 7.51, N 5.57 %)

$\nu_{\max}$  3482, 2908, 2792, 1604, 1485, 1395  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.22 (3 H, d,  $J=6.0$  Hz, Me a), 2.23 (3 H, s, Me b), 2.61-2.63 (1 H, m, Hb), 4.75 (1 H, d,  $J=6.5$  Hz, Ha), 4.95 (1 H, s, Hc), 7.26-7.46 (10 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.65, 35.45, 65.82, 84.75, 101.45, 127.85, 128.32, 128.64, 130.05, 130.25, 131.65, 131.72, 141.25.

$m/z$  EI (Found: 253.1469  $\text{C}_{17}\text{H}_{19}\text{NO}$  requires: 253.1468).

$[\alpha]_{\text{D}} = +46$  ( $\text{CCl}_4$ , 10 mg/ml).

**(4S,5S)- 2- [2- ( 1,1- Diphenyl phosphenyl) phenyl] 3,4- dimethyl- 5- phenyl- 1,3- oxazolidine** (4.10 g 78% yield) mp 233\_236 °C

[pseudophedrine(2.0 g 12.12 mmd) dphenyl phosphinebenzaldehyde(3.5 g 24.24 mmd)]



**31 c**

$\nu_{\max}$  (slurry, thin film) 3460, 2970, 1100, 1037, 1026  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.95 (3 H, Me a, d,  $J=6.5$  Hz, Me a), 2.52 (3 H, s, Me b), 2.73-2.79 (1 H, m, Hb), 4.22 (1 H, d,  $J=8.0$  Hz, Ha), 5.72 (1 H, s, Hc), 7.26-7.91 (19 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 14.15, 65.25, 85.22, 95.75, 126.5, 128.3, 129.11, 130.25, 130.46, 130.74, 130.89, 131.25, 131.45, 131.75, 132.55, 133.45, 133.77, 134.77, 140.65.

$m/z$  EI (Found: 437.1902  $\text{C}_{29}\text{H}_{28}\text{NOP}$  requires: 437.1908).

$[\alpha]_{\text{D}} = +82.5$  ( $\text{CCl}_4$ , 10 mg/ml).

## Ring Opening of Oxazolidines using $\text{NaBH}_3\text{CN}$ / TMSCl




1,3-Oxazolidine derived from ephedrine, pseudoephedrine or isoquinolinol (1.0 g), was dissolved in acetonitrile (40 ml); sodium cyanoborohydride (5 eq) was added, and then TMSCl (5 eq) was added to the vigorously stirred mixture at room temperature.

After 15 minutes the solvent was removed, DCM (30 ml) added and the solution was washed with water (2 × 15 ml). The organic solvents were removed and methanol (30 ml) added followed by potassium carbonate (5 eq) and the mixture stirred at room temperature overnight. The solid residue was filtered off, the filtrate evaporated to dryness, DCM (50 ml) added, and the organic solvents were washed with water (3 × 30 ml). The organic layer was dried over magnesium sulphate. Removal of the solvents gave the desired compound.

**(1R,2S)-2-[Methyl (phenyl methyl) amino]-1-phenylpropan-1-ol** (1.4 g 96% yield)

[2b (1.5 g 5.9 mmol),  $\text{NaBH}_3\text{CN}$  (1.86 g 29.52 mmol), TMSO (3.63 g 29.52 mmol)]



**10 b**

$\nu_{\text{max}}$  (neat) 3419, 3084, 3017, 2416, 1602, 1493, 1451, 1386  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.0 (3 H, d,  $J=6.0$  Hz, Me a), 2.21 (3 H, Me b, s, Me b), 2.91 (1 H, dq, Hb), 3.61 (1 H, br, OH), 3.62 (2 H, s, Hc, Hd), 4.8 (1 H, Ha, d,  $J=4.0$  Hz), 7.26-7.42 (10 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 10.12, 39.26, 59.15, 65.26, 74.26, 126.76, 126.92, 127.15, 127.91, 128.26, 128.63, 140.26, 142.35.

$m/z$  (EI) (Found: 255.1619  $\text{C}_{17}\text{H}_{21}\text{NO}$  requires: 255.1623).

$[\alpha]_{\text{D}} = -31$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1R,2S)-2-[Ethyl (methyl) amino]-1-phenylpropan-1-ol** (1.32 g 87% yield)

[2a (1.5 g 3.85 mmol),  $\text{NaBH}_3\text{CN}$  (2.47 g 39.25 mmol), TMSO (4.83 g 39.25 mmol)]



**10 a**

$\nu_{\text{max}}$  (neat) 3399, 3086, 2970, 1685, 1602, 1493, 1450, 1379  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.85 (3 H, d,  $J=3$  Hz, Me a), 1.0 (3 H, Me c, t,  $J=6$  Hz, Me c), 2.25 (3 H, s, Me b), 2.55 (2 H, q,  $J=6.0$  Hz, Hc, Hd, ), 2.83 (1 H, dq,  $J=3.4$  Hz, Hb), 4.80 (1 H, d,  $J=4$ . Ha), 7.20-7.30 (5 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 10.41, 14.25, 37.25, 62.51, 72.86, 125.6, 126.1, 126.81, 127.9, 142.10.

$m/z$  (EI) (Found: 193.1464  $\text{C}_{12}\text{H}_{19}\text{NO}$  requires: 193.1466).

$[\alpha]_{\text{D}} = -32$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1S,2S) - 2- [Ethyl- methyl amino]- 1 phenyl propane 1- ol** (1.1 g 94% yield)

[2c (1.2 g 6.28 mmd),  $\text{NaBH}_4\text{CN}$  (1.98 g 31.41 mmd), TMSO (3.86 g 31.41 mmd)]



**10 c**

$\nu_{\text{max}}$  (neat) 3405, 3112, 2985, 1692, 1610, 1496, 1385  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.75 (3 H, d,  $J=6.5$  Hz, Me a), 1.15 (3 H, t,  $J=7.0$  Hz, Me c), 2.25 (3 H, s, Me b), 2.65 (2 H, m, Hc, Hd), 2.4 (1 H, Hb, m, Hb), 4.2 (1 H, d,  $J=7.0$  Hz, Hb), 7.26-7.42 (5 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 9.53, 14.25, 37.61, 49.21, 62.65, 72.15, 126.15, 126.70, 126.85, 142.15.

$m/z$  (EI) (Found: 193.1463  $\text{C}_{12}\text{H}_{19}\text{NO}$  requires: 193.1466).

$[\alpha]_{\text{D}} = +56$  ( $\text{CCl}_4$ , 10 mg/ml).

**(1S,2S) - 2- [Methyl ( phenyl methyl) amino]- 1 phenyl propane 1- ol** (1.58 g 92% yield)

[2d (1.7 g 6.72 mmd),  $\text{NaBH}_4\text{CN}$  (2.12 g 33.60 mmd), TMSO (4.13 g 33.60 mmd)]



**10 d**

$\nu_{\text{max}}$  (neat) 3425, 3091, 3025, 1615, 1466, 1391  $\text{cm}^{-1}$ .


$\delta_H$  ( $CDCl_3$ , 250 MHz) 0.86 (3 H, d,  $J=6.5$  Hz, Me a), 2.22 (3 H, Me b, s, Me b), 2.83 (1 H, m, Hb), 3.45 (1 H, d,  $J=13$  Hz, Hc), 3.72 (1 H, d,  $J=9.7$  Hz, Hd), 4.33 (1 H, d,  $J=9.7$  Hz, Ha), 4.92 (1 H, br, OH), 7.27-7.63 (10 H, m).

$\delta_C$  ( $CDCl_3$ , 100 MHz) 9.26, 35.36, 54.25, 58.16, 75.26, 120.15, 123.6, 124.15, 124.62, 126.21, 126.83, 131.91, 140.25.

$m/z$  (EI) (Found: 255.1619  $C_{12}H_{21}NO$  requires: 255.1623).  $[\alpha]_D = +42$  ( $CCl_4$ , 10 mg/ml).

**(1S,2S)-2- {Methyl (1- methyl ethyl) amino} 1- phenyl propane- 1- ol** (1.92 g 95% yield)

[2e(2.0 g 9.76 mmd),  $NaBH_3CN$ (3.07 g 48.78 mmd), TMSO (6.0 g 48.78 mmd)]



$\nu_{max}$  (neat) 3299, 3857, 2971, 2616, 2398, 2361, 2251, 1604, 1493, 1457, 1342  $cm^{-1}$ .

$\delta_H$  ( $CDCl_3$ , 250 MHz) 0.83 (3 H, d,  $J=4.0$  Hz, Me a), 1.16 (6 H, dd,  $J=7.0$ , 7.0 Hz, Me c, Me d), 2.38 (6H, d,  $J=6.0$  Hz, Me c, Me d), 2.83 (1 H, m, Hd), 3.0 (1 H, dq,  $J=4.10$ , 9.70 Hz, Hb), 4.16 (1 H, d,  $J=10.0$ , Ha), 5.32 (1 H, br, OH), 7.26-7.42 (5 H, m).


$\delta_C$  ( $CDCl_3$ , 100 MHz) 10.99, 19.91, 20.16, 30.27, 54.25, 62.16, 127.12, 127.28, 127.62, 128.34, 141.26.

$m/z$  (EI) (Found: 207.1678  $C_{13}H_{21}NO$  requires: 207.1673).

$[\alpha]_D = +72.5$  ( $CCl_4$ , 10 mg/ml).

**(1R,2S)- 2- Methyl {1- methyl (1- methyl ethyl) amino} 1- phenyl propane- 1- ol**  
(2.2 g 88% yield)

[2f (2.5 g 12.2 mmd),  $\text{NaBH}_3\text{CN}$  (3.84 g 360.98 mmd), TMSO (75 g 60.98 mmd)]



$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.86 (3 H, d,  $J=7$  Hz, Me a), 1.05 (6 H, 2d,  $J=7.0$ , 7.0 Hz, Me c, Me d), 2.15 (3 H, s, Me b), 2.86-2.92 (1 H, m, Hd), 3.15-3.26 (1 H, m, Hb), 3.66 (1 H, br, OH), 4.82 (1 H, d,  $J=6.0$  Hz, Ha), 7.26-7.42 (5 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.21, 18.61, 18.92, 31.51, 51.25, 61.51, 72.15, 120.15, 127.25, 127.65, 141.52.

$m/z$  (EI) (Found: 207.1621  $\text{C}_{13}\text{H}_{21}\text{NO}$  requires: 207.1623).

$[\alpha]_{\text{D}} = -86$  ( $\text{CCl}_4$ , 10 mg/ml).

**(3S)- 2- {[4- (Methoxy) phenyl] methyl} 1,2,3,4- tetrahydroisoquinolin- 3- yl) methanol** (1.38 g 92% yield)

[2h (1.5 g 5.3 mmd),  $\text{NaBH}_3\text{CN}$  (1.68 g 26.69 mmd), TMSO (3.28 g 26.69 mmd)]



$\nu_{\text{max}}$  (thin film), 3357, 2931, 2834, 1611, 1511, 1249  $\text{cm}^{-1}$ .

$\delta_H$  ( $CDCl_3$ , 250 MHz) 2.4-2.5 (1 H, dd,  $J=5.0$ , 5.0 Hz, Ha), 2.8-2.9 (1 H, dd,  $J=6.0$ , 6.0 Hz, Hb), 3.1-3.2 (1 H, m, Hc), 3.5-3.6 (4 H, m. Hi, Hj, Hf, Hg), 3.7 (1H,OH , s), 3.9 (3 H,s, Me a), 6.9-7.2 (8 H, m).


$\delta_C$  ( $CDCl_3$ , 100 MHz) 26.10, 50.25, 58.15, 59.00, 62.15, 115.16, 126.72, 126.89, 127.15, 129.32, 130.52, 130.61, 131.21, 14.06, 134.70, 160.12.

$m/z$  (EI) (Found: 283.1570  $C_{18}H_{21}NO_2$  requires: 283.1572).

$[\alpha]_D = -39$  ( $CCl_4$ , 10 mg/ml).

(3S)-2-(2,2-Dimethylpropyl)-1,2,3,4-tetrahydroisoquinidin-3-yl]methanol  
(0.76 g 92% yield)

[2g (1.0 g, 3.56 mmol),  $\text{NaBH}_3\text{CN}$  (1.1 g, 13.74 mmol), TMSO (2.2 g, 17.79 mmol)]




8 d

$\nu_{\text{max}}$  (neat) 3364, 3021, 2952, 2417, 2254, 1643, 1495, 1479, 1360, 1188, 1035  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.92 (9 H, s, Me a), 2.26 (2 H, q,  $J=12$  Hz, Hc, Hd), 3.16 (2 H, AB,  $J=12$  Hz, He, Hf,), 3.42 (2 H, q,  $J=4.0$  Hz, Hi, Hj,), 3.65-3.67 (1 H, m, Hc), 4.21 (1 H, d,  $J=12$  Hz, Ha, Hb,), 5.27 (1 H, br, OH), 7.16-7.23 (5 H, m).

$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 25.26, 27.61, 51.26, 60.26, 63.35, 65.36, 126.26, 126.35, 126.90, 129.33, 134.62.  
 $m/z$  (EI) (Found: 233.18041 C<sub>15</sub>H<sub>23</sub>NO requires: 233.17996).  $[\alpha]_D = -41$  (CCl<sub>4</sub>, 10 mg/ml).

**(3S)- 1,2,3,4- Tetrahydro- 3- isoquinoline- 3- met hanol**

Lithium aluminium hydride (4.0 g, 10.54 mmol, 2 eq) was suspended in dry diethyl ether (150 ml). (S)-1,2,3,4-Tetrahydroisoquinoline 3-carboxylic acid (10 g, 57 mmol) was added. The mixture was stirred overnight at room temperature. Water (30 ml) was added carefully, then sodium hydroxide (20%) solution (40 ml) was added slowly. The solid precipitate was filtered off and the organic solutions extracted. The aqueous layer was extracted with diethyl ether (3 × 50 ml). The solid precipitate was dissolved in water (100 ml) and extracted with DCM (3 × 30 ml). The combined organic solutions were dried over magnesium sulphate. The solvent was removed to give an orange solid (8 g, 87%) mp 112-114 °C. <sup>76</sup>


(Found: C, 73.30, H, 7.53, N, 8.10, C<sub>10</sub> H<sub>13</sub> NO. requires: C, 73.57, H, 7.87, N, 8.38%).

δ<sub>H</sub> (CDCl<sub>3</sub>, 250 MHz) 2.5-2.8 (4 H, , m, Hd, He, Hf, Hg), 2.95-3.10 (1 H, m, Hc), 3.45 (1 H, dd, *J*=8.0 Hz, Ha), 3.7 (1 H, dd, *J*=4.0 Hz, Hb), 4.0 (2 H, br, OH), 7.0-7.1 (4 H, m).

δ<sub>C</sub> (CDCl<sub>3</sub>, 100 MHz) 31.16, 48.26, 55.26, 66.18, 120.16, 125.32, 130.16, 137.35, 168.26.

*m/z* EI (Found: 163.0996 C<sub>10</sub>H<sub>13</sub>NO<sub>2</sub> requires: 163.0997). [α]<sub>D</sub> = -96.5 (CCl<sub>4</sub>, 10 mg/ml).

## Synthesis of Oxazolidines derived from Isoquinoline




(3S),1,2,3,4-Tetrahydroisoquinoline-3-methanol was dissolved in toluene (50 ml). Then the required aldehyde (1 eq) was added, followed by PTSA (10 mol %). The mixture was heated under Dean Stark conditions overnight. Sodium hydrogen carbonate (30 ml) was added. The organic solution was evaporated to result in the required oxazolidine.

**R= ( MeOPh)**

**8a- (3R,10aS) - 3- [4- (Methoxy- phenyl)]- 1,5,10 tetrahydro[1,3]-oxazolid[3,4-b] isoquinoline** (1.87 g 88%) 1g (1.2 g 7.3 mmd)

[p- MeO benzaldehyde (0.88 g 7.3 mmd)]



**8 a**

(Found: C, 76.38, H, 6.65, N, 4.92,  $C_{18}H_{19}NO_2$ . requires: C, 76.46, H, 6.76, N, 4.98%).

$\nu_{\max}$  (slurry) 3421, 2937, 1612, 1512, 1242, 1167  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 2.85-3.10 (3 H, m, Hf, Hg, He), 3.4 (1 H, d,  $J=15$ , Ha), 3.7 (1 H, d,  $J=15$ , Hb), 3.8 (3 H, s, Me), 3.85 (1 H, d,  $J=8$  Hz, Hd), 4.25-4.8 (1 H, m, Hc), 4.75 (1 H, s, Hi), 6.8-7.5 (8 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 47.26, 55.26, 71.35, 72.16, 113.28, 114.26, 126.18, 127.35, 128.61, 129.31, 130.21, 137.35, 137.56, 161.26, 164.35.


$m/z$  (EI) (Found: 281.1410  $\text{C}_{18}\text{H}_{19}\text{NO}_2$  requires: 281.1416).

$[\alpha]_{\text{D}} = -92$  ( $\text{CCl}_4$ , 10 mg/ml), mp 186-188°.

**R = (C Me)<sub>3</sub>**

**8b - (3R,10aS) - 3- (1,1- Dimethyl (ethyl) - 1,5- 10,10a tetrahydro[1,3]oxazolo[3,4,6] isoquinoline (2.3 g 82%) (1 g (2.0 g 12.27 mmd)**

**[t- butyl- acetaldehyde (1.1 g 12.27 mmd)]**



**8 b**

(Found: C 77.98, H 8.86, N 6.19,  $\text{C}_{18}\text{H}_{21}\text{NO}_2$ . requires: C 77.92, H 9.09, N 6.06%).

$\nu_{\text{max}}$  (slurry) 3063, 2956, 1642, 1483, 1452, 1329, 1192  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.01 (9 H, Me1, s), 2.6 AB (3 H, He, Hf, Hg, m), 3.52 (1 H, Ha, d,  $J=7.0$  Hz), 3.6 (1 H, Hb, d,  $J=7.0$  Hz), 4.05 (2 H, Hd, Hi, m), 4.16 (1 H, Hi, s), 7.00-7.26 (4 H, m).

$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 25.16, 30.26, 52.15, 57.52, 70.14, 103.25, 105.15, 121.25, 125.15, 126.26, 126.90, 127.1, 127.6, 129.07, 134.14, 135.01, 136.2.

$m/z$  EI (Found: 231.1625 C<sub>15</sub>H<sub>21</sub>NO requires: 231.1623).

$[\alpha]_D = -96$  (CCl<sub>4</sub>, 10 mg/ml).

**(3S)-2-(2,2-Dimethyl propyl)-1,2,3,4  
[tetrahydroisoquinolin-3-yl]methyl (1-methyl  
ethyl)sulfide**



(3S)-2-(2,2-Dimethylpropyl)-1,2,3,4-[tetrahydroisoquinolin-3-yl]-methanol (1.5 g, 6.43 mmol) was dissolved in DCM (10 ml). The solution was cooled to 0 °C, triethylamine (0.11 g, 1.05 mmol, 3.5 eq) was added, then methylsulfonyl chloride (0.04 g, 0.027 mmol, 0.35 ml, 1.2 eq) was added followed by 2-propanthiol (0.068 g, 0.082 ml, 0.9 mmol, 3 eq), and the resulting solution was left to stir overnight. The solvent was removed and the residual oil subjected to flash chromatography using 95% petrol 5% ethyl acetate as eluent to yield the product as an oil (1.29 g, 69%).

$\nu_{\max}$  (neat) 2957, 2867, 1655, 1604, 1582, 1442, 1414, 1364, 1161, 1141 cm<sup>-1</sup>.


$\delta_H$  ( $CDCl_3$ , 100 MHz), 0.82 (9 H, s, Me b), 1.12 (1 H, m, Ha), 1.35 (6 H, d,  $J=7.0$  H, Me az), 1.46 (2 H, d  $J=8.0$  Hz, Hb, Hc), 2.0 (2 H, s, Hl, Hj), 2.31-2.33 (2 H, m, He, Hf), 2.61-2.73 (1 H, m, Hd), 3.16-3.72 (2 H, m, Hi, Hg), 7.52-7.53 (4 H, m,).

$\delta_C$  (100 MHz,  $CDCl_3$ ) 23.16, 25.26, 28.35, 51.26, 60.15, 65.32, 125.16, 126.30, 126.90, 133.60, 134.25.

( $m/z$ ) (EI) (Found: 291.2018  $C_{18}H_{29}NS$  requires: 291.2020).

$[\alpha]_D = +75$  ( $CCl_4$ , 10 mg/ml).

**Phenyl methyl N-(1S)-2-[methyl(methoxy)amino]-2 oxo-1-Me-(phenyl methyl)(ethyl)ethyl)carbamate**




N-tert-butoxycarbonyl-L-phenylalanine (10 g, 40.16 mmol) OR N-(carbobenzyloxy)-L-phenylalanine (10g, 33.4 mmol) was dissolved in DCM (150 ml) and the solution cooled to -20 °C. N-O Dimethyl hydroxylamine hydrochloride (1 eq) was added, followed by N-methyl morpholine (1 eq). EDCI (1 eq) was added in portions over a period of 1 hour at -20 °C. The reaction was allowed to stir for a further 1 hour at -20°C. The reaction was allowed to stir for a further 1 hour. Ice cooled 1M HCl was added (30 ml) and the mixture extracted with DCM, and the aqueous layer was re-extracted with DCM (3 × 10 ml). The combined organic solutions were washed with aqueous sodium hydrogen carbonate (30 ml) and brine (30 ml). The combined aqueous was washed with DCM (3 × 15 ml) and the combined organic solutions dried over magnesium sulphate. Removal of the solvent and purification by chromatography yielded the desired compound

**R=( Ph CH<sub>2</sub>) =27a**

**Phenyl methyl N- (1S)- 2- [methyl( methoxy) amino]- 2- oxo-1- phenyl methyl ethyl car bamate (11 g 80% yield)**

[Cbz- L- phenylalanine(10 g 40.16 mmd), N, O- Dimethyl hydroxylamine(3.9 g 40.1 mmd), N- methylmorpholine(4.1 g 40.16 mmd), EDC (7.8 g 40.16 mmd)]



**27 a**

$\nu_{\text{max}}$  (neat) 3405, 2843, 1660, 1670, 1341, 1134  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 2.7 (1 H, d,  $J=6.0$  Hz, Ha), 2.8 (1H, d,  $J=6.0$ Hz, Hb), 3.1 (3 H, s, Me b), 3.6 (3 H, s, Me a), 3.6 (1 H, Hc, t,  $J=6.0$ Hz, Hc) , 5.0 (2 H, s, He, Hd), 7.2-7.5 (10 H, aromatics).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 32.16, 35.26, 42.52, 52.21, 61.26, 126.46, 127.91, 128.43, 128.72, 128.16, 129.21, 129.40, 129.64, 155.26, 171.21.

$m/z$  EI (Found 342.1749  $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_4$  required 342.1746).

**R=Q(CH<sub>3</sub>)<sub>3</sub>=27b**

**1,1-Dimethyl-N-[(1S)-2-[methyl-(methoxy)amino]-2-oxo-1-(phenyl methyl ethyl) carbamate] (9.5 g 92% yield)**

[BOG L- phenylalanine( 10 g 33.4 mmol), N, O dimethyl hydroxylaminehydrochloride( 3.26 g 33.4 mmol), N- methylmorpholine( 3.38 g 33.4 mmol), EDCI ( 6.4 g 40.16 mmol)]. mp 186- 189 °C



**27 b**

 $\nu_{\text{max}}$  (neat) 3411, 2872, 2695, 1672, 1346, 1145  $\text{cm}^{-1}$ .



$\delta_H$  (CDCl<sub>3</sub>, 250 MHz) 1.43 (9 H, s, Me c), 3.15 (1 H, d,  $J=7.5$  Hz, Ha), 3.16 (1H, d,  $J=7.5$ Hz, Hb ), 2.82-2.83 (1 H, t,  $J=7.7$ Hz, Hc), 3.26 (3 H, s, Me b), 3.72 (3 H, s, Me a), 7.26-7.51 (5 H, m).

$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 28.15, 32.62, 38.16, 51.26, 61.52, 126.25, 126.62, 127.15, 128.2, 137.26, 155.25, 172.23.

*m/z* EI (Found: 308.1739 C<sub>16</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> requires: 308.1736).

$$[\alpha]_D = -32 \text{ (CCl}_4, 10 \text{ mg/ml)}.$$

**Phenyl methyl N-[(1S)-1 formyl-2-phenyl ethyl] carbamate**




N-BOC substituted carbamate (1 eq) or (1b) N-Cbz substituted carbamate (1 eq) was dissolved in dry THF (30 ml). The solution was cooled to 0 °C in an ice bath. Then LiAlH<sub>4</sub> in THF 1.0M (1/2 eq) was added dropwise. The solution was left to stir for 30 minutes, then cooled to -15 °C and saturated aqueous potassium hydrogen sulphate (20 ml) was added carefully, the mixture was diluted with diethyl ether (50 m), and stirred vigorously for further 30 minutes. The mixture was allowed to reach room temperature and the organic layer separated. The aqueous layer was extracted with diethyl ether (3 × 15 ml). The combined organic solutions were dried over magnesium sulphate. Removal of solvent at temperature no greater than 30 °C yielded the required material which was recrystallized from methanol to yield the required material.

$R = \text{C}(\text{CH}_3)_3 = 28b$

1,1- Dimethyl- N- [(1S)- 1- phenyl- 2- phenyl ethyl car bamate (1.15 g 97% yield)

[N-BOC-L-phenyl carbamate (1.5 g 4.8 mmol),  $\text{LiAlH}_4$  (2.4 ml, 2.4 mmol)]



28 b

(Found: C, 66.68, H, 7.74, N, 5.58,  $\text{C}_{14}\text{H}_{19}\text{NO}_3$ . Requires: C, 67.74, H, 7.26, N, 5.56%.  $[\alpha]_{\text{D}} = -34$  ( $\text{CCl}_4$ , 10 mg/ml). mp 174-177 °C

$\nu_{\text{max}}$  ( $\text{CH}_2\text{Cl}_2$  solution) 3425, 2817, 1687, 1472, 1315, 1192  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.42 (9 H, s, Me a ), 3.18 (1 H, d,  $J=7.4$  Hz, Ha), 3.2 (1H, d,  $J=7.6$ Hz, Hb), 3.16-3.17 (1 H, m, Hc.), 7.16-7.26 (5 H, m), 9.82 (1 H, Hd, s).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 28.15, 35.25, 60.15, 127.15, 128.4, 129.25, 130.15 135.15, 192.35, 196.15.

**R=(CH<sub>2</sub>Ph)=28 a**

**Phenyl methyl- N- [(1S)-1-formyl-2-phenylethyl] carbamate (0.94 g 95% yield)**

[N-Cbz-L-phenyl carbamate (1.2 g 3.5 mmd), LiAlH<sub>4</sub> (2.0 ml, 1.75 mmd)]



**28 a**


$\nu_{\text{max}}$  (slurry) 3407, 2843, 1750, 1680, 1198 cm<sup>-1</sup>

$\delta_{\text{H}}$  (CDCl<sub>3</sub>, 250 MHz) 2.63 (2 H, d,  $J=12.0$  Hz, Ha, Hb), 3.16 (1 H, m, Hc), 4.16 (1H, br, NH), 5.52 (2 H, s, He, Hf), 7.16-7.31 (10 H, m), 9.8 (1 H, s, Hh).

$\delta_{\text{C}}$  (CDCl<sub>3</sub>, 100 MHz) 39.15, 39.26, 120.21, 123.6, 124.2, 131.25, 134.21, 136.26, 138.26, 140.35, 168.26, 171.21.

$m/z$  EI (Found: 283.1203 C<sub>17</sub>H<sub>17</sub>NO<sub>3</sub> requires: 283.1208).  $[\alpha]_{\text{D}} = -12$  (CCl<sub>4</sub>, 10 mg/ml).

**Phenyl methyl N- [(1S)- 1- (1,3- dithiane- 2yl)- 2- phenyl]  
carbamate**




Methanol (50 ml) was cooled to 0 °C, and acetyl chloride (5 ml) was added. The solution was stirred for 10 minutes under nitrogen. The carbamate (28a-b) (1 eq) was added as a solution in methanol (10 ml) and DCM (10 ml) to help solubility. The resulting solution was stirred overnight. The solvent was removed and the resulting solid recrystallized from methanol to yield the title compound.

$R = (\text{CH}_2\text{Ph}) = 29\text{a}$

Phenyl methyl N-[(1S)-1-(1,3-dithiane-2-yl)-2-phenyl] carbamate (0.72 g  
92% yield)

[28a (0.6 g 2.12 mmol), propane-1,3-dithiol (0.34 g 3.18 mmol) acetylchloride (5 ml)]



29 a

$\nu_{\text{max}}$  (slurry) 3328, 3028, 1713, 1510, 1453  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.31-1.44 (1 H, m, Hb), 1.82-1.83 (2 H, m, Hc, Hd), 2.85-2.35 (6 H, m, He-j, Hm), 4.22 (1 H, d,  $J=3.0$  Hz, Ha), 5.1 (2 H, s, Hk, Hl), 7.2-7.4 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 23.16, 25.26, 45.32, 47.61, 62.16, 127.12, 127.18, 128.32, 128.43, 128.82, 128.94, 129.21, 129.47, 129.65, 129.74

$m/z$  (EI), (Found: 373.1164  $\text{C}_{20}\text{H}_{27}\text{NO}_2\text{S}_2$  requires: 373.1170).


$[\alpha]_{\text{D}} = -14$  ( $\text{CCl}_4$ , 10 mg/ml). Mp 143-146  $^{\circ}\text{C}$

**R=(CH<sub>3</sub>)<sub>3</sub>=29b**

**1,1-Dimethylethyl-N-[(1S)-1-(1,3-dithiane-2-yl)-2-phenylethyl]**

**carbamate (1.03 g 97% yield)**

[28b- (0.8 g 3.22 mmol), propane-1,3-dithiol (0.51 g 4.82 mmol) acetylchloride(5 ml)]



**29 b**


(Found: C 60.20, H 7.34, N 4.2, C<sub>17</sub>H<sub>15</sub>NO<sub>2</sub>S<sub>2</sub>. requires: C 60.18, H 7.37, N 4.2% ). [α]<sub>D</sub> = -39.5 (CCl<sub>4</sub>, 10 mg/ml). Mp 165-168 °C

ν<sub>max</sub> (slurry, nujol mull) 3428, 3072, 2969, 1723, 1515, 1462 cm<sup>-1</sup>.

δ<sub>H</sub> (CDCl<sub>3</sub>, 250 MHz) 1.42 (9 H, s, Me), 2.10-2.21 (1 H, m, Hb), 2.9-3.4 (6 H, He, m, He-j, Hm), 4.12 (2 H, d, J=9.0 Hz, Hc, Hd), 4.22 (1 H, br, NH), 4.83 (1 H, d, J=7.5 Hz, Ha), 7.26-7.41 (5 H, m).

δ<sub>C</sub> (CDCl<sub>3</sub>, 100 MHz) 5.25, 28.10, 30.26, 38.15, 50.25, 55.15, 80.25, 122.15, 126.5, 127.1, 129.21, 140.15, 158.15.

## Synthesis of Phosphorus Imine Ligands




To a stirred solution of the phosphine aldehyde (1.2 eq) in DCM (30 ml), the L amine ester hydrochloride salt (1 eq) was added, followed by CSA (10 mol %). 4Å Molecular sieve (2.0 g) was added and the resulting mixture allowed to stand for 72 hours. The molecular sieve was removed by filtration and the solution washed with sodium bicarbonate ( $3 \times 20 \text{ ml}$ ). The organic solutions were dried and the crude mixture subjected to column chromatography using 1% triethylamine, 19% ethyl acetate and 80% petrol as eluent to yield the desired product (32a-c) as coloured oils.

$R = CH(CH_3)_2$

**(32a) Methyl- 2- {( E)- 1- [2- (1,1- diphenyl phosphanyl) phenyl] methyldene}**

**-(3S)- methyl butanoate (3.45 g 72% yield)**

[Phosphinealdehyde (4.15 g 14.33 mmol), Lvalinemethyl (2.0 g 11.94 mmol)]



$\nu_{\max}$  3033, 2960, 2872, 2831, 1732, 1633, 1435, 1194  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.62 (3 H, d,  $J=6.7$  Hz, Me a), 0.86 (3 H, d,  $J=6.8$  Hz, Me b), 2.15-2.16 (1H, m, Hc), 3.42 (1 H, d,  $J=7.2$  Hz, Hb), 3.61 (3 H, s, Me c), 7.25-7.54 (14 H, m), 8.7 (1 H, s, Ha).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 17.67, 18.51, 30.09, 52.13, 77.02, 128.36, 128.42, 128.98, 129.09, 129.19, 129.33, 134.62, 134.65, 136.68, 162.13, 172.60.

$m/z$  FAB (Found: 404.1729 ( $\text{M}^++1$ )  $\text{C}_{25}\text{H}_{27}\text{NO}_2\text{P}$  requires: 404.1651).  $[\alpha]_{\text{D}} = +20.2$  ( $\text{CCl}_4$ , 10 mg/ml).

$R = \text{CH}_2 (\text{C}_6\text{H}_4\text{P}(\text{H}))$

(32c) Methyl- 2- {[( E)- 1- [2- (1,1- diphenyl phosphanyl) phenyl] methyldene} amino] - 1-3- (2S- hydroxyphenyl) propanoate (2.65 g 66% yield)

[L- Tyrosinemethyl ester HCl (2.0 g 8.64 mmol), phosphinealdhyde(3.0 g 10.37 mmol)]



$\nu_{\text{max}}$  3055, 1739, 1516, 1437, 1263, 1170  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 2.15 (1 H, dd,  $J=5.9, 10.5$  Hz, Hc), 3.16 (1 H, dd,  $J=5.9, 10.5$  Hz, Hd), 3.52 (3 H, s, Me), 4.16 (1 H, dd,  $J=10.5, 10.5$  Hz, Hb), 5.25 (1 H, br, OH), 6.95-7.8 (18 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 39.4, 52.46, 75.28, 115.72, 128.35, 128.42, 128.92, 128.96, 129.03, 129.18, 129.22, 130.99, 131.12, 137.73, 134.18, 134.28, 155.05, 162.76, 172.92.


$m/z$  FAB (Found: 468.1729 ( $\text{M}^++1$ )  $\text{C}_{29}\text{H}_{27}\text{NO}_2\text{P}$  requires: 468.1651).

$[\alpha]_{\text{D}} = +62$  ( $\text{CCl}_4$ , 10 mg/ml).

R=Ph

(32b) Methyl- 2- {(E)- 1- [2- (1,1- diphenyl phosphanyl) phenyl] methyldene amino} -2S- phenyl ethanoate (4.27 g 70% yield)

[L- Phenylalaninemethyl ester (3.0 g 13.95 mmd), diphenyl phosphinealdehyde (4.86 g 16.74 mmd)]



32 b

$\nu_{\max}$  (Neat) 3070, 2983, 2875, 1739, 1645, 1462, 1438, 1197  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 3.62 (3 H, s, Me), 5.00 (1 H, s, Hb), 7.16-7.83 (20 H, m), 8.2 (1 H, s, Ha).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 52.25, 70.31, 128.21, 128.30, 128.53, 128.06, 129.06, 129.18, 129.25, 131.24, 132.38, 132.46, 133.15, 133.35, 133.62, 133.75, 162.61, 171.60.

$m/z$  FAB (Found: 438.1618 ( $\text{M}^++1$ )  $\text{C}_{28}\text{H}_{25}\text{NO}_2\text{P}$  requires: 438.1545).

$[\alpha]_{\text{D}} = +19.5$  ( $\text{CCl}_4$ , 10 mg/ml).

## Synthesis of Imines from Nitriles




To a stirred solution of the Grignard reagent (1.5 eq) in ether under nitrogen, the nitrile (1 eq) was added slowly as a solution in ether, then heated under reflux and the reaction was followed by TLC. When all the starting nitrile had reacted, the reaction was quenched by the addition of dry methanol (20 eq) carefully into the cooled reaction mixture at  $-15\text{ }^{\circ}\text{C}$  -  $-20\text{ }^{\circ}\text{C}$ . The solid precipitate was filtered off and the crude product used for the next step.

**R=Ph, R<sup>1</sup>=Me**

**(23a) Methylphenyl imine (4.9 g 85% yield)**

[Benzonitrile (5.9 g 48.5 mmol), methyl magnesium bromide (27 ml, 1.5 eq)]



**23 a**

$\nu_{\text{max}}$  (neat) 3560, 2982, 2800, 1600  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 2.42 (3 H, s, Me), 7.25-7.46 (5 H, m), 7.92 (1 H, d,  $J=4.25\text{ Hz}$ , Ha).

$m/z$  EI (Found: 119.0746,  $\text{C}_8\text{H}_9\text{N}$ , requires: 119.0668).

**R=Ph, R<sup>1</sup>=Et**

**(23c) Ethylphenyl imine** (5.5 g 86% yield)

[Benzonitrile (5.0 g 48.5 mmol), methyl magnesium bromide (32 ml, 1.5 eq)]



$\nu_{\text{max}}$  (neat) 3570, 2982, 2811, 1607  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.15 (3 H, t,  $J=7.4$  Hz, Me), 2.43 (2 H, q,  $J=7.8$  Hz, Hb, Hc), 7.26-7.47 (5 H, m), 7.92 (1 H, d,  $J=4$  Hz, Ha,).

$m/z$  EI (Found: 133.0813,  $\text{C}_9\text{H}_{11}\text{N}$ , requires: 133.0891).

**R<sup>1</sup>=propyl, R=Ph**

**(23b) Propylphenyl imine** (6.10 g 85% yield)

[Benzonitrile (5.0 g 48.5 mmol), propyl magnesium chloride (36.5 ml, 1.5 eq)]



$\nu_{\text{max}}$  (neat) 3530, 2982, 2885, 2822, 1610  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.93 (3 H, t,  $J=4.6$  Hz, Me), 1.82 (2 H, dq,  $J=4.57, 4.5$  Hz, Hb, Hc), 2.93 (2 H, t,  $J=4.5$  Hz, Hd, He), 7.26-7.46 (5H, m), 7.92 (1 H, s, Ha).

$m/z$  EI (Found: 147.1126,  $\text{C}_{10}\text{H}_{13}\text{N}$ , requires: 147.1048).

$\text{R}^1 = i\text{Pr}$ ,  $\text{R} = \text{Ph}$

**(23f) Isopropylphenyl imine** (7.2 g 92% yield)

[Benzonitrile (5.0 g 48.5 mmol), isopropyl magnesium chloride (36.4 ml, 1.5 eq)]



**23 f**

$\nu_{\text{max}}$  (neat) 3555, 2982, 2887, 1622  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.10 (6 H, d,  $J=6.8$  Hz, Me a, Me b), 3.0 (1 H, m, Hb), 7.2-7.4 (5 H, m), 7.9 (1 H, d,  $J=6.8$  Hz, Ha).

$m/z$  EI (Found: 161.1126,  $\text{C}_{11}\text{H}_{15}\text{N}$ , requires: 161.1204).

$R^1 = 1\text{-NaPh}$ ,  $R = 1\text{-NaPh}$

(23 g) 1,1- Dinaphthalene- imine (9.1 g 65% yield)

[1- Cyanonaphthalene(7.6 g 50 mmd), 1- bromonaphthalene(20.7 ml, 100 mmd)  
magnesium(2.43 g 100 mmd)]



$\nu_{\max}$  (neat) 3555, 2982, 2887, 1630  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 7.1-8.0 (14 H, m), 8.1 (1 H, s, Ha).

$m/z$  EI (Found: 281.1283,  $\text{C}_{21}\text{H}_{15}\text{N}$ , requires: 281.1204).

**R<sup>1</sup>=2- Naph, R=2- Naph**

**2,2- Dinaphthalene- imine (4.36 g 62% yield)**

(23h) [2- Cyanonaphthalene(2.14 g 25 mmd), 2- bromonaphthalene( 10.83 g 50 mmd), magnesium( 1.215 g 50 mmd)]




$\nu_{\text{max}}$  (neat) 3561, 2979, 2892, 1640  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 7.1-8.0 (14 H, m), 8.1 (1 H, s, Ha ).

$m/z$  EI (Found: 281.1282,  $\text{C}_{21}\text{H}_{15}\text{N}$ , requires: 281.1204).

## Synthesis of Sulfur Imine Ligands




To a stirred solution of the imine (1 eq) in dry DCM, L-methionine methyl ester hydrochloride (1 eq) was added, then CSA 10 mol %, followed by dry 4Å molecular sieves . The reaction mixture was allowed to stand at room temperature for 48 hours. The molecular sieve was then filtered off and the mixture washed with saturated aqueous sodium bicarbonate (3 × 20 ml). The organic solutions were dried over magnesium sulfate and removal of the solvents yielded the crude product which was purified by column chromatography using 1% triethylamine, 15% ethyl acetate, 84% hexane.



R- Ph, R<sup>1</sup>=propyl

(24b) Methyl (2s)-2-[(z)-2-[(z)-1-phenylbutylidene]amino]-4-(methylthio)butanoate (11.34 g 81% yield)

[Propylphenyl imine (7.0 g 47.62 mmol), L-methoninemethyl ester (5.7 g 47.62 mmol)]



24 b

$\nu_{\max}$  3588, 2941, 2871, 2825, 1625  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.9 (3 H, t,  $J=4.5$  Hz, Me a), 1.51 (2 H, d q,  $J=4.6, 4.8$  Hz, Ha, Hb), 2.1 (3 H, s, Me b), 2.2 (2 H, t,  $J=4.8$  Hz, Hc, Hd), 2.3-2.4 (2 H, m, Hg, Hf), 2.5-2.6 (2 H, m, Hi, Hj), 3.6 (3 H, s, Me c), 4.5 (1 H, t,  $J=3.3$  Hz, He). 7.2-7.6 (5H,m)


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 16.25, 20.51, 20.64, 28.37, 32.89, 39.51, 52.10, 63.62, 126.71, 126.83, 127.02, 127.16 140.89, 172.62.

$m/z$  EI (Found: 293.1530  $\text{C}_{16}\text{H}_{23}\text{NO}_2\text{S}$  requires: 293.1528).

**R=Ph, R<sup>1</sup>=iPr**

(24c) Methyl (2 s)- 2- [(z) (methyl- 1- phenyl, pr opylidene amino) - 3- 4- methyl thio] butanoate (7.27 g, 73% yield)

[isopropyl-phenyl-imine (5.0 g 34 mmol), L-methioninemethylest HCl (6.8 g 34 mmol)]

 $\nu_{\max}$  3598, 2961, 2841, 2827, 1635  $\text{cm}^{-1}$ .

$\delta_H$  (CDCl<sub>3</sub>, 250 MHz) 1.1 (3 H, d,  $J=4.5$ , Hz, Me a), 1.1(3H,d,  $J=4.6$ Hz, Me b), 2.1 (3 H, s, Me c), 2.2-2.3 (2 H, m, Hc, Hd,), 2.4-2.41 (2 H, m, He, Hf), 2.6 (1 H, m, Ha), 3.6 (3 H, s, Me d,), 3.91 (1 H, t,  $J=4.7$  Hz, Hb), 7.2-7.4 (5 H, m).


$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 15.56, 20.14, 20.52, 28.27, 32.79, 39.41, 52.05, 63.43, 126.44, 126.61, 126.92, 129.15, 140.72, 172.48.

*m/z* EI (Found: 293.1529 C<sub>16</sub>H<sub>23</sub>NO<sub>2</sub>S requires: 293.1527).

**R=Ph, R<sup>1</sup>=Ph**

**(24d) Methyl (2 s)-2-((diphenyl methyldene) amino 3-4- methyl  
thiol)butanoate (1.18 g 72% yield)**

[Benzophenoneimine (0.91 g, 5 mmol), L-methioninemethyl ester HCl (0.8 g, 5 mmol)]



(Found: C 69.72, H 6.51, N 4.19, requires: C 69.72, H 6.42, N 4.28%).

 $\nu_{\max}$  3610, 2979, 2835, 2822, 1840  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.92 (3 H, s, Me a), 2.15 (2 H, t,  $J=6.7$  Hz, Hd, He), 2.45-2.46 (2 H, m, Hc, Hb), 3.61 (3 H, s, Me b), 4.22 (1 H, t,  $J=6.2$  Hz, Ha.), 7.1-7.5 (10 H, m).


$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 15.33, 30.54, 33.59, 64.10, 127.82, 128.12, 128.57, 128.77, 130.99, 136.26, 171.23.

*m/z* EI (Found: 327.1293 C<sub>19</sub>H<sub>21</sub>NO<sub>2</sub>S requires: 327.1293).

$R^1=1\text{-Naph}$ ,  $R=1\text{-Naph}$

(24f) (2s)-2-[(1,t)Dinaphthalene-2-ylmethylidene]amino-4-(methylthio)butanoate (5.0 g 52% yield)

[1,1-Dinaphthalene-imine (6.36 g 22.6 mmol), L-methioninemethylester HCl (4.5 g 22.6 mmol)]



$\nu_{\max}$  3610, 2979, 2962, 2835, 2822, 1645  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 2.15 (3 H, s, Me a), 2.26-2.27 (2 H, m, He, Hd), 2.43-2.46 (2 H, m, Hb, Hc), 3.52 (1 H, t,  $J=4.7$  Hz, Ha), 3.6 (3 H, s, Me b), 7.1-7.9 (14 H, m).


$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 15.79, 30.89, 34.36, 53.81, 123.21, 125.32, 125.58, 126.19, 127.09, 127.49, 127.56, 128.31, 128.69, 130.29, 133.29, 176.62.

$m/z$  EI (Found: 427.1687  $\text{C}_{27}\text{H}_{25}\text{NO}_2\text{S}$  requires: 427.1684).

**R=2- Naph, R<sup>1</sup>=2- Naph**

**( 24 g) ( 2 s) - 2 - [( 1,t) Dinaphthalene- 2- ylmethylidene] amino- 4- ( methylthiol) butanoate ( 7.38 g 54% yield)**

[2,2- Dinaphthalene imine( 9 g 32 mmd), L- methioninemethylester HQ ( 6.37 g 32 mmd)]




$\nu_{\max}$  3615, 2982, 2965, 2880, 2815, 1615  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 2.17 (3 H, s, Me a), 2.36-2.37 (2 H, m, He, Hd), 2.61-2.62 (2 H, m, Hb, Hc), 3.52 (1 H, t,  $J=4.7$  Hz, Ha), 3.6 (3 H, s, Me b), 7.1-7.9 (14 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 15.28, 25.16, 33.85, 52.08, 53.22, 64.26, 125.03, 125.44, 125.87, 126.22, 126.89, 127.22, 127.59, 128.45, 128.95, 130.54, 143.36, 171.27.

$m/z$  EI (Found: 427.1685  $\text{C}_{27}\text{H}_{25}\text{NO}_2\text{S}$  requires: 427.1684).

## E- 1,3- Diphenyl- 3- hydroxy- 1- propene



Cerium (III) chloride heptahydrate (19.8 g, 53.2 mmol) was dissolved in methanol (250 ml) and chalcone (10g, 48 mmol) added. The resulting solution was cooled to 0 °C. Sodium borohydride (2.0 g, 53.2 mmol) was added to the solution in small portions over a period of 5 minutes such that effervescence was kept to a minimum. The reaction was stirred for a further 3 hours, water (100 ml) added, and the organic layer separated. The aqueous layer was extracted with ethyl acetate (3 × 100 ml), the combined organic extracts dried over magnesium sulphate and the solvent removed to yield an oil which turned to a white solid on standing, this was recrystallized from hexane to give needle-like crystals (9.9 g, 94%) mp 53-55 °C. <sup>76</sup>


(Found: C 85.62, H 6.75, C<sub>15</sub>H<sub>14</sub>O requires: C 85.68, H 6.71%)

$\nu_{\max}$  (CH<sub>2</sub>Cl<sub>2</sub> solution) 3348, 3033, 1501, 1450, 968 cm<sup>-1</sup>.

$\delta_{\text{H}}$  (CDCl<sub>3</sub>, 250 MHz) 2.71 (1 H, br, OH), 5.2 (1 H, d, *J*=6.4 Hz, Hc), 6.40 (1 H, dd, *J*=6.4, 16 Hz, Hb), 6.62 (1 H, d, *J*=16.0 Hz, Ha), 7.26-7.52 (10 H, m).

$\delta_{\text{C}}$  (CDCl<sub>3</sub>, 100 MHz), 79.16, 125.25, 126.15, 126.32, 127.52, 128.23, 130.26, 131.62, 136.26, 141.26, 141.37.

## E- 1,3- Diphenyl- 3- acetoxy- 1- propane



1, 3-Diphenyl-2-propan-1-ol (0.3 g, 1.43 mmol) was dissolved in DCM (20 ml) and pyridine (0.23 ml, 286 mmol, 2 eq) added. To the resulting clear solution was added 4-dimethylaminopyridine (100 mg, 0.82 mmol) and acetic anhydride (0.15 ml, 1.5 mmol, 1.05 eq), and the reaction was stirred for 3 hours. The mixture was concentrated, diethylether (100 ml) was added, and the solution was washed with aqueous copper sulphate (3 × 30 ml). The organic solutions were extracted and washed with sodium hydrogen carbonate solution (3 × 10 ml). The organic solution was dried over magnesium sulphate and the solvent removed to yield a colourless oil (0.3 g, 83%).<sup>76</sup>


(Found: C 80.90, H 6.41,  $\text{C}_{17}\text{H}_{16}\text{O}_2$  requires: C 80.92, H 6.40%)

$\nu_{\text{max}}$  (neat film) 3029, 1739, 1501, 1375, 1239  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.11 (3 H, s, Me), 6.34 (1 H, dd,  $J=16, 7$  Hz, Hb), 6.45 (1 H, d,  $J=7.0$  Hz, Hc), 6.63 (1 H, d,  $J=16$  Hz, Ha), 7.18-7.43 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz), 22.16, 76.72, 127.53, 127.82, 128.26, 129.42, 129.52, 129.73, 133.26, 136.16, 140.26, 171.26.

## Dimethyl-2- - 1,3- diphenyl prop-2- enyl propandioate



Allyl palladium chloride (22 mg, 0.06 mmol, 5% mol) was added to a solution of *anti*-2- (S)- (2-pyridyl)-1,3-dithiane 1-oxide (25 mg, 0.119 mmol, 10%) in DCM (1 ml) was added. The resulting solution was stirred for 30 minutes. 1, 3-Diphenylprop-2-enyl acetate (0.300g, 1.19 mmol) was added as a solution in DCM (5 ml), a catalytic amount of potassium acetate was added, and the resulting solution stirred for a further 30 minutes. Caesium carbonate (0.53 g, 2.38 mmol, 2 eq) and dimethyl malonate (0.31 g, 2.38 mmol, 2 eq) were added and the resulting solution stirred for 24 hours. The solution was filtered through Celite, the solvents removed and the residue purified by column chromatography using 5% ethyl acetate, 95% petroleum ether as eluent to yield the compound as a white solid (0.18 g, 93%), mp 123-124 °C.


(Found: C 73.59, H 6.18,  $\text{C}_{20}\text{H}_{20}\text{O}_4$  requires: C 74.06, H 6.21%).<sup>76</sup>

$\nu_{\text{max}}$  ( $\text{CHCl}_3$  solution) 3031, 2965, 1756, 1437  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 3.42 (3 H, s, Me a), 3.65 (3 H, s, Me b), 4.0 (1 H, d,  $J=16.0$  Hz, Hd), 4.21 (1 H, dd,  $J=16, 8$  Hz, Hc), 6.31 (1 H, dd,  $J=16, 8$  Hz, Hb), 6.95 (1 H, d,  $J=16$  Hz, Ha), 7.15-7.30 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz), 52.20, 52.50, 57.60, 126.42, 127.16, 127.62, 127.83, 128.82, 129.16, 132.19, 137.27, 140.26, 168.37, 168.26

## (-)-(Oxo camphor sulfonyl) imine



Reagent grade ammonium hydroxide (500 ml) was cooled to 0 °C. A solution of (+)-10-Camphorsulphonyl chloride (50 g 199.5 mmol) in DCM (500 ml) was added slowly to the stirred ammonium hydroxide solution. The mixture was stirred at 0 °C for 4 hours and allowed to reach room temperature. The organic layer was removed and the aqueous layer extracted with DCM (3 × 100 ml). The combined organic extracts were dried over magnesium sulphate and evaporated to dryness to yield the crude (+)-(1S)-10-camphorsulphonamide (35g, 76%), which was judged sufficiently pure for use in the preparation of (-)-(oxocamphorsulphonyl) imine.

The crude (+)-(1S)-camphorsulphonamide was added slowly to acetic acid (750 ml) and selenium dioxide (25 g, 225 mmol) at 0° in a 3 l flask. The reaction was heated to reflux overnight. The reaction was allowed to cool and filtered to remove the precipitated selenium. Water (250 ml) was added to the filtrate in a separating funnel, and the organic layer separated. The aqueous layer was extracted with DCM (3 × 125 ml) and the combined organic extracts dried over magnesium sulphate. Evaporation to dryness under reduced pressure yielded the crude product which was subsequently recrystallized from chloroform to yield the pure material as pale yellow crystals (23 g, 52%), mp 197-199 °C.<sup>75</sup>


(Found: C, 52.90; H, 5.68; N, 6.30  $C_{10}H_{13}NO_3S$ . requires: C, 52.86, H, 5.73, N, 6.18%).  $[\alpha]_D = -179^\circ$  ( $CCl_4$ , 10 mg/ml).

$\nu_{max}$  (slurry) 2957, 1772, 1359, 1169  $cm^{-1}$ .

$\delta_H$  ( $CDCl_3$ , 250 MHz) 0.99 (3 H, Me a, s, Me a), 1.17 (3 H, s, Me b), 1.8 – 2.1 (2 H, m, Hd, He), 2.2 – 2.4 (2 H, m, Hb, Hc), 2.78 (1 H, d,  $J=5.0$  Hz, Ha), 3.24 (1 H, d,  $J=13.6$  Hz, Hf), 3.45 (1 H, d,  $J=13.6$  Hz, Hg).

$\delta_C$  ( $CDCl_3$ , 100 MHz) 18.25, 20.14, 22.26, 27.95, 44.67, 50.67, 60.03, 62.95, 181.7, 198.26.

## (+)-[(8,8-Die thoxy camphoryl)sulfonyl] imine



(-)-(Oxocamphorsulphonyl)imine (2.0 g, 9 mmol) was dissolved in dry ethanol (60 ml). To this was added triethyl orthoformate (6.5 g, 7.3 ml, 43.95 mmol, 5 eq), concentrated hydrochloric acid (1 ml) and amberylst-15 ion exchange resin (1.0 g), and the reaction heated to reflux. After refluxing overnight, the reaction was allowed to cool to room temperature, water (30 ml) added and the organic layer separated. The aqueous layer was extracted with DCM ( $3 \times 100$  ml), the combined organic extract dried over magnesium sulphate, and the solvent removed. Recrystallization of the crude solid from methanol-water (1:2) gave the title material as a white crystalline solid (2.5 g, 65%), mp 144-146 °C.

(Found: C 55.92; H 7.72, N 4.72,  $\text{C}_{14}\text{H}_{23}\text{NO}_4\text{S}$ . requires: C 55.79, H 7.71, N 9.65%).


$\nu_{\text{max}}$  ( $\text{CH}_2\text{Cl}_2$  solution) 1662, 1345, 1169  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.03 (3 H, s, Me a), 1.08 (3 H, s, Me b), 1.21 (3 H, t,  $J=7.0$  Hz, Me c), 1.23 (3 H, t,  $J=7.0$  Hz, Me d), 1.76 – 2.05 (4 H, m, Hb, Hc, Hd, He), 2.1 (1H, d,  $J=5.0$  Hz, Ha), 2.81 (1 H, d,  $J=13.5$  Hz, He), 2.82 (1 H, d,  $J=13.5$  Hz, Hj), 2.87 (2 H, q,  $J=7.0$  Hz, Hh, Hi), 2.90 (2 H, q,  $J=7.2$  Hz, Hj,

Hk).

$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 14.80, 15.12, 20.82, 20.91, 20.99, 29.56, 46.28, 49.15, 53.02, 58.73, 58.92, 64.53, 103.26.

### (+)-[8,8-Diethoxy camphoryl sulphonyl] oxaziridine



Commercial aqueous hydrogen peroxide (0.6 ml, 5.12 mmol, 30% eq w/v) was added to a stirred suspension of potassium carbonate (0.37 g, 2.7 mmol) in methanol (15 ml) at room temperature. (+)-[8,8-Diethoxycamphoryl] sulphonyl] imine (0.35 g, 1.16 mmol) was added, and the reaction allowed to stir at room temperature for 8 hours. After this time, saturated brine (20 ml) and DCM (20 ml) was added and the organic later separated. The aqueous layer was extracted with DCM (2 × 20 ml) and the combined organic extracts washed with aqueous sodium sulphite solution (20 ml, 5% w/v) and dried over magnesium sulphate. Removal of the solvent with a bath temperature below 40 °C, followed by recrystallization from methanol furnished the title compound as a colourless crystalline solid (0.24 g, 65%), mp 117-119°C.

(Found: C 52.78, H 7.24, N 4.49. requires: C 52.80, H 7.16, N 4.39%).


$\nu_{\max}$  (slurry) 1370, 1162, 1069 cm<sup>-1</sup>.

$\delta_H$  (CDCl<sub>3</sub> 250 MHz) 1.04 (3 H, s, Me a.), 1.24 (3 H, t,  $J=6.0$  Hz, Me c), 1.34 (3

H, s, Me b), 1.42 (3 H, t,  $J=6.2$  Hz, Me d), 1.78-1.98 (4 H, m, Hb, Hc, Hd, He), 3.10 (1 H, d,  $J=14.0$  Hz, Ha), 3.29 (1 H, d,  $J=14.0$  Hz, Hf), 3.43 (1 H, d,  $J=14.0$  Hz, Hg), 3.55 (1 H, Hi, Hj, q,  $J=7.0$  Hz), 3.65 (2 H, Hk, Hl, q,  $J=7.2$  Hz).

$\delta_C$  (CDCl<sub>3</sub> 100 MHz) 14.95, 15.20, 20.70, 21.80, 28.30, 45.33, 47.65, 53.50, 54.65, 58.50, 59.10, 97.90, 102.52.

## 2-(2-Pyridyl)-1,3-dithiane



Distilled pyridine-2-carboxaldehyde (2.38 g, 22.2 mmol) and 1,3-propanedithiol (2.48 g, 23.15 mmol, 1.05 eq) were dissolved in toluene (100 ml), and *p*-toluene sulphonic acid (0.25 g) was added. The mixture was heated to reflux for 48 hours with a Dean-Stark trap attached, then allowed to cool to room temperature. The mixture was washed with 1.0 M sodium hydroxide (2 × 25 ml) and the organic portion extracted and dried over magnesium sulphate. The solvent was removed and the residue subjected to flash column chromatography with gradient elution from 10% ethyl acetate to 30% ethyl acetate in petroleum ether, to yield the title compound as a pale yellow oil (2.9 g, 66%).<sup>75</sup>


(Found: C 54.82, H 5.58, N 7.17, C<sub>9</sub>H<sub>11</sub>NS<sub>2</sub> requires: C 54.79, H 5.62, N 7.10%).

$\nu_{\text{max}}$  (neat) 2902, 1469, 1432, 1221, 1049 cm<sup>-1</sup>.

$\delta_{\text{H}}$  (CDCl<sub>3</sub>, 250 MHz) 1.8-2.26 (2 H, m, Hd, He), 2.9-3.15 (4 H, m, Hb, Hc, Hf, Hg), 5.36 (1 H, s, Ha), 7.18-7.72 (3 H, m), 8.45-8.60 (1 H, m).

$\delta_{\text{C}}$  (CDCl<sub>3</sub>, 100 MHz) 25.37, 31.26, 53.26, 53.65, 121.62, 123.46, 137.62, 150.27, 157.52.

## ***Anti*-2-(2-Pyridyl)-1,3 dithiane 1-(*S*)-oxide**



DCM (30 ml) was cooled to  $-20\text{ }^{\circ}\text{C}$  and 30% aqueous hydrogen peroxide (0.94 ml, 8.12 mmol, 4 eq) added, followed by diazabicycloundec-7-ene (DBU) (1.27 g, 8.12 mmol, 4 eq). After five minutes (+)-[8,8-(Diethoxycamphoryl)] sulfonylimine (0.67 g, 2.23 mmol, 1.1 eq) was added and the solution stirred for 20 minutes. 2-(2-Pyridyl)-1,3-dithiane (0.4 g, 2.03 mmol, 1 eq) was added as a solution in DCM (5 ml) dropwise over 5 minutes. The reaction was allowed to stir for 3 hours, then quenched by the addition of saturated aqueous sodium sulfide solution (10 ml) and allowed to reach room temperature. The organic layer was separated and the aqueous layer extracted with DCM ( $3 \times 50\text{ ml}$ ). The combined organic extracts were washed with 0.1 M hydrochloric acid and the acid washings extracted with DCM ( $2 \times 30\text{ ml}$ ). The combined organic extracts were dried over magnesium sulphate and evaporated. The residue was subjected to flash chromatography with gradient elution from 100% DCM to 2% methanol in DCM to yield first (+)-[(8,8-diethoxycamphoryl) sulphonylimine (0.55 g, 82%) and the title material (0.4 g, 92%) as a white solid mp  $123\text{--}126\text{ }^{\circ}\text{C}$


(Found: C 50.59, H 5.2, N 6.53,  $\text{C}_9\text{H}_{11}\text{NOS}_2$  requires: C 50.68, H 5.20, N 6.57%).  $[\alpha]_{\text{D}} = 54$  ( $\text{CCl}_4$ , 10 mg/ml). ee 98%.

$\nu_{\text{max}}$  (slurry), 3050, 1586, 1434,  $1035\text{ cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.20-2.89 (5 H, m), 3.6-3.75 (1 H, m), 4.75 (1 H, Ha, s, Ha), 7.26-7.72 (3 H, m), 8.6 (1 H, d,  $J=5$  Hz).

Absolute stereochemistry confirmed by X-ray as *trans* (Andrew Lund thesis)<sup>75</sup>

## Synthesis of 1,3-Dithianes from Para and Meta Nitro Substituted Benzaldehyde




Propane -1, 3-dithiol (1.1 eq) was added at 0 °C to a stirred solution of the aldehyde. The solution was stirred for two minutes and acetyl chloride (10 ml) added. After stirring overnight, the solution was evaporated and the residue recrystallized from methanol to yield the required compound. <sup>76a</sup>

$R_2 = NO_2$      $R_1 = H$

2- (4- Nitrophenyl) - 1,3- dithiane (5.91 g 93%)

(para nitro benzaldehyde (4.0 g 26.5 mmd), propane 1,3- dithiol (3.15 g 29.15 mmd))



(Found: C 49.58, H 4.51, N 5.71,  $C_{10}H_{11}NO_2S_2$  requires: C 49.79, H 4.56, N 5.30%). mp 120-122 °C

$\nu_{\text{max}}$  (slurry) 2935, 1451, 1392, 1276  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.0 (2 H, m, Hf, Hg), 2.72 (4 H, m, Ha, Hb, Hf, Hg), 5.22 (1 H, s), 7.4-7.5 (2 H, m, Hl, Hj), 8.2-8.9 (2 H, m, Hi, Hk).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 24.76, 31.76, 50.37, 123.94, 128.01, 140.13, 147.61.

$\text{R}_1=\text{NO}_2$   $\text{R}_2=\text{H}$

**2- (3- Nitrophenyl) - 1,3- dithiane** (5.8 g 91%)

(metanitrobenzaldehyde (4.0 g 26.5 mmd), propane1,3- dithiol (3.15 g 29.15 mmd)



**61 b**


(Found: C 50.13, H 4.55, N 5.78,  $\text{C}_{10}\text{H}_{11}\text{NO}_2\text{S}_2$  requires: C 49.79, H 4.56, N 5.80%). mp 146-148 °C.

$\nu_{\text{max}}$  (solution DCM) 2942, 1462, 1341, 1281  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.0-2.7 (1 H,m, Hb, Hc), 3.16-3.20 (4 H, m, Hd, He, Hf, Hg), 5.22 (1 H,s, Ha), 7.50-7.52 (1 H, m, Hh, Hi), 7.92-7.93 (1 H, m, Hk), 8.30-8.36 (1 H, m, Hl).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 25.01, 31.21, 51.26, 122.31, 131.26, 135.41, 141.27, 149.62.

## Synthesis of Benzaldehyde Amine Derivatives from the Corresponding Nitro




To a stirred mixture of (25a) or (25b) in a mixture of 50:50 ethanol and water, tin (3 eq) was added and the resulting mixture stirred for 10 minutes. Concentrated HCl (7 ml) was added, and the resulting mixture heated under reflux overnight. After cooling, sodium hydroxide pellets were added until the solution was basic. The mixture was separated and the aqueous later extracted with DCM (3  $\times$  30 ml). The combined organic solutions were dried over magnesium sulphate. The solvent was removed and residue purified by column chromatography in petrol 80% ethyl acetate 20%, to yield the desired compound as a yellow solid.

(p-nitro-1,3 dithiane(2.0 g, 8.3 mmol), Sn(28 g, 23.17 mmol)

$R_4=H, R_3=NH_2$

**2- (3- Aminophenyl) - 1,3- dithiane (1.7 g 97%)**

(3- nitro- 1,3 dthiane(2.0 g 8.3 mmd), Sn(2.5 g 23.17 mmd)




(Found: C 56.73, H 5.99, N 6.41,  $C_{10}H_{13}NS_2$  requires: C 50.87, H 6.13, N 6.6%). mp 138-140 °C.

$\nu_{\max}$  (slurry) 3492, 2965, 1592, 1497, 1398  $cm^{-1}$ .

$\delta_H$  ( $CDCl_3$ , 250 MHz), 2.1 (2 H, Hd, He, m), 3.26-3.29 (4 H, Hb, Hc, Hf, Hg, m), 3.52 (2 H, br,  $NH_2$ ), 5.16 (1 H, s, Ha), 6.52-6.54 (1 H, Hh, m), 6.81-6.83 (1 H, Hi, m), 7.16-7.18 (1 H, Hj, m), 7.3 (1 H, Hk, s).

$\delta_C$  ( $CDCl_3$ , 100 MHz) 29.25, 32.14, 51.26, 114.25, 115.62, 118.15, 130.25, 140.25, 149.29

## Synthesis of 1,3-Dithiane oxides from corresponding NO<sub>2</sub> and NH<sub>2</sub> derivatives




To a stirred solution of (25a-b) or (26a-b) in DCM, (+)-[ (8, 8-dimethoxy camphoryl) sulphonyl oxaziridine (1.1 eq) was added at -20 °C and the resulting solution stirred for 24 hours. The solution was concentrated and the residue chromatographed in 10% petrol 90% ethyl acetate as eluent to yield the sulfoxide.

R<sub>2</sub>=NO<sub>2</sub> - R<sub>1</sub>=H

2- ( 4- Nitr ophenyl) - 1,3- dithiane 1- oxide (0.26 g 82%)

(25a(0.3 g 1.24 mmd), oxaziridine(0.4 g 1.37 mmd))



(Found: C 46.69, H 4.28, N 5.45, C<sub>10</sub>H<sub>11</sub>NO<sub>3</sub>S<sub>2</sub> requires: C 46.69, H 4.22, N

5.39%). mp 145-147 °C.

$\nu_{\text{max}}$  (slurry) 2985, 1591, 1492, 1392, 1195  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.51-2.6 (2 H, m, Hd, He), 2.72-3.65 (2 H, m, Hf Hg), 3.7-4.4(2H, m, Hb, Hc), 4.62 (1 H, s, Ha), 7.5 (2 H, d,  $J=8.0$  Hz, Hh, Hk), 8.2 (2 H, d,  $J=8.0$  Hz, Hi, Hj).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 29.01, 31.06, 55.12, 70.25, 125.21, 129.32, 140.15, 148.32.

$\text{R}_1=\text{NO}_2$   $\text{R}_2=\text{H}$

**2- (3- Nitrophenyl) - 1,3- dithiane- 1- oxide** (0.31 g 97%)

(26b(0.3 g 1.24 mmd),  $\alpha$ aziridine(0.4 g 1.37 mmd))



**62 e**

(Found: C 46.55, H 4.19, N 5.38,  $\text{C}_{10}\text{H}_{11}\text{NO}_3\text{S}_2$  requires: C 46.69, H 4.28, N 5.45%). mp 166-168 °C.

$\nu_{\text{max}}$  2979, 1598, 1496, 1393, 1196  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.51-2.58 (2 H, m, Hd, He), 2.73-3.65 (2 H, m, Hf, Hg), 3.72-4.62 (2H, m, Hb, Hc), 4.82 (1 H, s, Ha), 7.52-7.55 (1 H, m, Hi), 7.8 (1


H, d,  $J=8.0$  Hz, Hn), 8.2 (1 H, d,  $J=8.0$  Hz, Hj), 8.7 (1 H, s, Hk).

$\delta_C$  (CDCl<sub>3</sub>, 100 MHz) 29.21, 31.25, 59.32, 69.25, 124.25, 125.32, 136.25, 148.35.

$R_2 = \text{NH}_2$   $R_1 = \text{H}$

2- (4- Aminophenyl) - 1,3- dithiane- 1- oxide (0.2 g 92%)

(26a(0.21 g 0.96 mmd), oxaziridine(0.39 g 1.34 mmd))



62b

(Found: C 52.89, H 5.71, N 5.98,  $\text{C}_{10}\text{H}_{13}\text{NS}_2\text{O}$  requires: C 52.86, H 5.72, N 6.17% mp 132-134 °C.  $\nu_{\text{max}}$  3425, 2985, 1586, 1496, 1398, 1176  $\text{cm}^{-1}$ ).


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 2.52-2.55 (2 H, m, Hd, He), 2.73-3.52 (4 H, m, Hb, Hc, Hf, Hg), 3.75 (2 H, br,  $\text{NH}_2$ ), 4.62 (1 H, s, Ha), 6.52 (2 H, d,  $J=8.0$  Hz, Hh, Hk), 7.44 (2 H, d,  $J=8.2$  Hz, Hi, Hj).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 29.21, 31.25, 55.16, 69.27, 115.26, 122.35, 129.27, 147.25.

$R_1 = H$   $R_2 = NH_2$

**2- (3- Aminophenyl) - 1,3- dithiane- 1- oxide** (0.24 g 93%)

(26b (0.24g 1.1 mmd),  $\alpha$ aziridine (0.35 g 1.21 mmd))




(Found: C 52.92, H 5.76, N 5.94,  $C_{10}H_{13}NS_2O$  requires: C 52.86, H 5.72, N 6.17%). mp 159-165 °C.

$\nu_{max}$  3435, 2976, 1572, 1496, 1392, 1162  $cm^{-1}$ .

$\delta_H$  ( $CDCl_3$ , 250 MHz), 2.32-2.34 (2 H, m, Hd, He), 2.43-2.63 (4 H, m, Hb, Hc, Hf, Hg), 3.62 (2 H, br,  $NH_2$ ), 4.85 (1 H, s, Ha), 6.92-6.96 (2 H, m, Hi), 7.25-7.28 (2 H, m, Hh, Hj), 7.8 (1 H, s, Hk).

$\delta_C$  ( $CDCl_3$ , 100 MHz) 29.36, 32.15, 56.25, 69.76, 116.26, 122.75, 131.25, 148.25.

## 2 -{[1,1 -Dimethyl thio]methyl}pyridine



2-Methyl-2-propanthiol (2.0 g, 22.2 mmol) was dissolved in dry THF (15 ml) and the solution was cooled to  $-78\text{ }^{\circ}\text{C}$ . *N*-butyllithium (19.4 ml, 31.04 mmol) was added slowly, and the resulting mixture left to stir for one hour. Purified 2-chloromethyl pyridine (3.6 g, 22 mmol) was added dropwise at  $-78\text{ }^{\circ}\text{C}$ . The resulting mixture left to stir for 3 hours. Water (30 ml) was added and the organic solution extracted with DCM ( $2 \times 40\text{ ml}$ ). The solvent was removed and the residue chromatographed in petrol ethyl acetate 70/30 as eluent to yield the required compound (2.3 g, 58%).


(Found: C 66.10, H 7.87, N 7.20,  $\text{C}_{10}\text{H}_{15}\text{NS}$  requires: C 66.29, H 8.29, N 7.73%). mp  $159\text{-}165\text{ }^{\circ}\text{C}$ .

$\nu_{\text{max}}$  (neat) 2960, 2898, 1590, 1568, 1475,  $1459\text{ cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.43 (9 H, s, Me), 3.9 (2 H, s, Hb), 7.1-7.16 (7 H, m, He), 7.4-7.42 (1 H, m, Hc), 7.6-7.65 (1 H, m, Hd), 8.5-8.53 (1 H, Hf, m, Hf).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 8.24, 26.32 53.35, 126.32, 139.35, 139.64, 140.21, 149.26.

### 3-Ethyl cyclohexan-1-one



Ethyl magnesium chloride in diethyl ether 1M (20 ml) was added to copper iodide (0.52 g, 2.93 mmol) under nitrogen and the resulting mixture stirred for 15 minutes at  $0^\circ\text{C}$ . Cyclohex-2-enone (1.9 g, 20.75 mmol) was added and the mixture stirred for 50 minutes. The mixture was added to ice cold 10% sulfuric acid, and left to stir for 30 minutes. The mixture was extracted with diethyl ether ( $3 \times 30\text{ ml}$ ), and the organic solution dried over magnesium sulfate. The organic solution was evaporated to dryness and the resulting yellow oil distilled under a pressure of 0.43 m bar to yield the product as a colourless oil (1.1 g, 42%).


$\nu_{\text{max}}$  3506, 2935, 1712, 1454, 1313, 1226  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 0.82 (3 H, s,  $J=7.4\text{ Hz}$ , Me), 1.26-1.28 (4 H, m, Hh, Hi, Hj, Hk), 1.65-1.66 (2 H, m, Hb), 2.16-2.18 (3 H, m, Hc, He, Hd), 2.7.2.73 (4 H, m, Hd, He, Hf, Hg).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.12, 23.52, 25.39, 25.45, 20.23, 35.97, 38.15, 211.15

$m/z$  EI (Found: 126.1041  $\text{C}_8\text{H}_{14}\text{O}$  requires: 126.1044).

### 4-Phenyl-hexan -2-one



Copper iodide (0.25 g, 1.3 mmol) was suspended in diethyl ether (30 ml) and cooled to 0 °C. Ethyl magnesium bromide 1M (9 ml, 8.2 mmol) was added and the reaction stirred for 15 minutes. *Trans*-4-phenylbut-3-en-2-one (1.5 g, 7.1 mmol) was added and the resulting solution stirred overnight. The reaction mixture was added to a mixture of ice cooled water (100 ml) and 10% sulphuric acid (30 ml) and stirred for 2 hours. The mixture was extracted with diethyl ether (3 × 30 ml), the solvent removed and the crude product chromatographed over silica in 9:1 petroleum ether ethylacetate to yield the required product (1.2 g, 95%).


$\nu_{\text{max}}$  (neat) 2963, 1716, 1495, 1412, 1356, 1163  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.73 (3 H, t,  $J=7.6$  Hz, Me a), 1.62-1.63 (2 H, m, Hb), 2.0 (3 H, s, Me b), 2.62-2.64 (2 H, m, He, Hd), 3.16-3.18 (1 H, m, Hc).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.95, 29.34, 30.47, 42.42, 50.42, 126.30, 127.53, 128.37, 128.41, 207.59.

$m/z$  FAB (Found 177.1239, ( $\text{M}^++1$ )  $\text{C}_{12}\text{H}_{17}\text{O}$ , requires 177.1201).

## 1,3 -Diphenylpentan -1 -one



Copper iodide (0.25 g, 1.3 mmol) was suspended in diethyl ether (30 ml) and the mixture cooled to 0 °C. Ethylmagnesium bromide 1.0M (9 ml, 8.2 mmol) was added and the reaction was stirred for 15 minutes. Chalcone (1.5 g, 7.1 mmol) was added and the resulting solution stirred overnight. The reaction mixture was added to a mixture of ice cooled water (100 ml) and 10% sulphuric acid (30 ml), and stirred for 2 hours. The mixture was extracted with diethyl ether (3 × 30 ml), the solvent removed and chromatographed over silica in 9:1 petroleum ether ethylacetate to yield the product (1.60 g, 95% ).


$\nu_{\text{max}}$  (neat) 2955, 1680, 1600, 1448, 1408 cm<sup>-1</sup>.

$\delta_{\text{H}}$  (CDCl<sub>3</sub>, 250 MHz) 0.75 (1 H, m, Hc), 0.82 (3 H, t,  $J=7.4$  Hz, Me), 1.63-1.64 (2 H, m, Hb), 3.16-3.18 (2 H, m, Hd, He), 7.2-7.9 (10 H, m).

$\delta_{\text{C}}$  (CDCl<sub>3</sub>, 100 MHz) 11.81, 29.09, 36.11, 41.39, 125.64, 126.64, 126.28, 127.97, 128.26, 128.42, 137.33, 144.70, 199.15.

$m/z$  FAB (Found 239.1430, ( $M^++1$ ) C<sub>17</sub>H<sub>19</sub>O, requires 239.1435).

### 3-Ethyl cyclopentan-2-one



Copper iodide (0.25 g, 1.3 mmol) was suspended in diethyl ether (30 ml) and cooled to  $0\text{ }^\circ\text{C}$ . Ethylmagnesium bromide 1M (9 ml, 8.2 mmol) was added and the reaction was stirred for 15 minutes. Cyclopenten-2-enone (1.5 g, 7.1 mmol) was added and the resulting solution stirred overnight. The reaction mixture was added to a mixture of ice cooled water (100 ml) and 10% sulphuric acid (30 ml) and stirred for 2 hours. The mixture was extracted with diethyl ether ( $3 \times 30\text{ ml}$ ), the solvent removed and the crude product chromatographed over silica in 9:1 petroleum ether ethylacetate to yield the required product (0.73 g, 95%).


$\nu_{\text{max}}$  1675, 1435, 1406, 1321,  $1198\text{ cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.83 (2 H, q,  $J=6.4\text{ Hz}$ , Hb), 0.92 (3 H, t,  $J=6.4\text{ Hz}$ , Me), 1.25-1.26 (1 H, m, Hc), 1.62-1.65 (2 H, m, Hh, Hi), 2.16-2.20 (4 H, m, Hd, He, Hf, Hg).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 12.53, 28.54, 28.91, 38.55, 38.95, 41.52, 219.99.

$m/z$  EI (Found 112.0885,  $\text{C}_7\text{H}_{12}\text{O}$ , requires 112.0888).

**General procedure for conjugate addition  
of diethyl zinc to cyclohexenone  
(S) -3 -Ethyl -cyclohexanone**



To a stirred solution of copper triflate (0.0045 g, 12 mmol, 10 mol %) and chiral ligand (20 mol %) at 0 °C in toluene, enone (0.097 g, 1.1 mmol, 1 eq) and diethylzinc (3 ml, 3 eq) were added sequentially. The resulting solution was stirred for 2 hours, then poured into 1 M HCl (20 ml), and the mixture extracted with ethyl acetate (3 × 40 ml). The organic layer was dried over magnesium sulfate and the solvent removed to leave the required product as a yellow oil, which was purified by chromatography to yield the title compound as a colourless oil.

$\nu_{\text{max}}$  3506, 2935, 1712, 1454, 1313, 1226  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.86 (3 H, t,  $J=7.4$  Hz, Me), 1.26-1.29 (2 H, m, Hb, Ha), 1.61-1.63 (2 H, m, Hj, Hi), 2.16 (1 H, m, Hc), 2.26-2.30 (6 H, m, Hd, He, Hg).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.12, 23.52, 25.45, 26.33, 35.97, 38.15, 211.10.

$m/z$  EI (Found: 126.1041,  $\text{C}_8\text{H}_{14}\text{O}$ , requires: 126.1044).

$[\alpha]_{\text{D}} = 69^\circ$  ( $\text{CCl}_4$ , 10 mg/ml).

## 2-Phenyl 2,5-dihydrofuran



Bromobenzene (79 mg, 0.5 mmol), 2,3 -dihydrofuran (175 mg, 2.5 mmol),  $\text{Pd}(\text{OAc})_2$  (30 mg, 5 mol %), triphenylphosphine (20 mg, 15 mol %) and silver carbonate (300 mg, 1.78 mmol), were added to acetonitrile (25 ml) and the reaction heated at 80 °C for 3 days. The solvent was removed and the residue chromatographed in ethyl acetate to yield the product as a pale yellow oil (0.05 g, 68%).

$\nu_{\text{max}}$  neat 2100, 1550, 1200, 1100  $\text{cm}^{-1}$ .


$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 4.4-4.6 (2 H, m, Hc, Hb), 5.6-5.8 (2 H, m, Hd, He), 5.6 (1 H, d,  $J=4.8$  Hz, Ha), 7.5-7.8 (5 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100MHz), 67.68, 100.43, 103.15, 118.51, 128.03, 128.35, 129.45,

130.55

$m/z$  EI (Found 146.0731,  $\text{C}_{10}\text{H}_{10}\text{O}$ , requires 146.0731).

## General Procedure for the Asymmetric Heck Reaction



To a stirred solution of  $\text{Pd}_2(\text{dba})_3$  (10 mmol%), benzene (30 ml) and triethylamine (10 ml) were added and the mixture stirred for 10 minutes, then the ligand (20% mmol) was added and the mixture heated at 40 °C for 2 hours.

Phenyltriflate (1 eq) and alkene (5 eq) were added and the mixture heated to 40°C for 3 days. The solvent was removed and the residue chromatographed in 7:3 mixture of ethyl acetate-petroleum ether to yield the product (60 mg, 70%).


$\nu_{\text{max}}$  2800, 2750, 2250, 1100, 1050  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 1.52 (1 H, t,  $J=10.3$  Hz, Ha), 1.63-1.64 (2 H, m, Hf, Hg), 1.72-1.73 (2 H, m, Hf, He), 2.26-2.18 (2 H, m, Hc, Hd), 6.16 (1 H, d,  $J=2.4$  Hz, Hb), 7.2-7.4 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 22.18, 22.43, 22.86, 24.86, 121.35, 124.76, 124.96, 125.46, 128.39, 128.41, 128.75, 128.86, 142.71, 143.33.

*m/z* EI (Found 234.1411, C<sub>18</sub>H<sub>18</sub>O, requires 234.1408).

## General procedure for the palladium catalysed allylic displacement reaction



To a stirred solution of the ligand (10 mol) in DCM (30 ml) at 0°C, allylpalladium chloride dimer (5 mol %) was added and the mixture stirred for 10 minutes.

1,3-Diphenyl-2-propylacetate (1 eq) was added and the resulting mixture stirred for 0.5 hour. Cesium carbonate (5 eq) and dimethylmalonate (5 eq) were added sequentially and the resulting mixture stirred for 24 hours at 0 °C. The solvent was filtered through celite, the celite was washed with DCM (3 × 30 ml), the combined organic solutions were evaporated and the residue chromatographed in 9:1 petroleum ether diethyl ether mixture, to yield the products.

(Found: C 73.92, H 6.18.  $\text{C}_{20}\text{H}_{25}\text{NO}_4$  requires: C 74.06, H 6.21%). mp 118-123 °C.


$\nu_{\text{max}}$  3031, 2952, 1735, 1452  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 3.45 (3 H, s, Me b), 3.66 (3 H, s, Me a), 3.99 (1 H, d,  $J=10.0$  Hz, Ha), 4.26 (1 H, dd,  $J=8.0, 10.0$  Hz, Hb), 6.31 (1 H, dd,  $J=8, 16$  Hz, Hc), 6.48 (1 H, d,  $J=16.0$  Hz, He), 97.20-7.35 (10 H, m).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 49.10, 52.31, 52.49, 126.21, 127.01, 127.42, 127.75,

128.35, 128.75, 129.01, 132.15, 136.72, 140.01, 167.72, 168.21.

### 3-Ethyl cyclohexanone 2,4 - dini trophenyl hydrazone



To a stirred solution of 3-ethyl cyclohexan-1-one (1 eq) in ethanol (30 ml), 2 or 3 drops of sulfuric acid were added followed by 2, 4 -dinitrophenylhydrazine (1 eq) and the solution was stirred for 20-30 minutes. The mixture was washed with sodium hydrogen carbonate (2 × 10 ml) and extracted with ethyl acetate (3 × 30 ml). The organic solutions were dried over magnesium sulfate and the solvent removed to yield the crude product as an oil which was chromatographed in petroleum ether -ethyl acetate 80:20 to yield an orange solid.


$\nu_{\max}$  3425, 2950, 1618, 1589, 1517, 1334, 1309  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz), 0.93 (3 H, t,  $J=7.5$  Hz, Me), 1.43-1.44 (3 H, m, Hb, Hc), 1.63-1.64 (2 H, m, Hd, He), 2.16-2.18 (3 H, m, Hf, Hg), 3.24-3.30 (4 H, m, Hh, Hi, Hk, Hl), 7.92 (1 H, d,  $J=4.0$  Hz, Hp), 8.1 (1 H, d,  $J=4.0$  Hz, Hn), 9.16 (1 H, s, Hm).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 11.68, 25.21, 26.19, 27.37, 29.53, 29.60, 40.77, 116.60, 124.01, 130.33, 145.72, 161.75.

$m/z$  FAB (Found: 307.1406 ( $\text{M}^++1$ )  $\text{C}_{14}\text{H}_{19}\text{N}_4\text{O}_4$  requires: 307.1328).

**(1S,2S)-2-[1,2-Dimethyl propyl (methyl) amino]  
-1-phenyl propyl -3,5-dinitrobenzoate**



To a solution of 3,5-dinitrobenzoyl chloride (160 mg, 0.7 mmol, 1.2 eq), in diethyl ether (20 ml) at 0°C amino alcohol (1S,2S)-2[1,2-dimethylpropyl(methyl) amino]-1-phenyl propan -1-ol (150 mg, 0.64 mmol) was added, followed immediately by DMAP (76 mg, 0.64 mmol). An immediate white precipitate appeared. After one hour the solid precipitate was collected and purified by column chromatography using petroleum ether- ethyl acetate-triethylamine (80:19:1) as eluent to yield the product as an oil (150 mg, 55%). Appendix 5

$\nu_{\text{max}}$  2970, 2461, 1739, 1542, 1458, 1346, 1265, 1157  $\text{cm}^{-1}$ .

$\delta_{\text{H}}$  ( $\text{CDCl}_3$ , 250 MHz) 0.70 (3 H, d,  $J=8.0$  Hz, Me a), 0.77 (3 H, d,  $J=8.0$  Hz, Me b), 0.80 (3 H, d,  $J=12.0$  Hz, Me c), 1.00 (3 H, d,  $J=8.0$  Hz, Me d), 1.60 (1 H, septet,  $J=8.0, 8.0$  Hz, Ha), 2.15 (3 H, s, Me e), 2.38 (1 H, dq,  $J=8.0, 12.0$  Hz, Hb), 7.2-7.5 (5 H, m), 9.15 (2 H, d,  $J=4.0$  Hz, He, Hg), 9.20 (1 H, s, Hf).

$\delta_{\text{C}}$  ( $\text{CDCl}_3$ , 100 MHz) 22.18, 12.96, 19.52, 20.96, 27.80, 32.00, 62.27, 66.21, 80.19, 122.09, 127.68, 128.68, 129.44, 134.78, 138.19, 148.56,

161.73.

*m/z* FAB (Found: 430.1972, ( $M^++1$ )  $C_{22}H_{18}N_3O_6$  requires 430.1899).

## **REFERENCES**

1. Trost, B.M.; Van Vranken, D.L.; Bingel, C. *J. Am. Chem. Soc.* **1992**, *114*, 9327.
2. Åkermark, B.; Zetterberg, K.; Hansson, S.; Krakenberger, B.; Vitagliano, A. *J. Organomet. Chem.* **1987**, *335*, 139.
3. Oslob, S.D.; Åkermark, B.; Helquist, P.; Norrly, P.O. *Organometallics*. **1997**, *16*, 3015.
4. Blanchadell, V.; Moreno-Manal, M.; Pajuelo, F.; Plexats, R. *Organometallics*. **1999**, *18*, 4934.
5. Shibasaki, M.; Sodeoka, M. *Syn. Org. Chem. Jpn.* **1994**, *52*, 9561.
6. Cabri, W.; Candiani, J.; Bedeschi, A.; Penco, S; Santi, R. *J. Org. Chem.* **1992**, *57*, 1471.
7. Dietev, R.K.; Tokles, M. *J. Am. Chem. Soc.* **1987**, *109*, 2040.
8. Stewart, K.R.; Lever, J.R.; Whangbo, M.H. *J. Org. Chem.* **1982**, *47*, 1472.
9. Naslund, J.; Welch, C. *Tetrahedron : Asymmetry*. **1991**, *2*, 1123.
10. Benson, C.S.; Snyder, J. *Tetrahedron Lett.* **1991**, *32*, 5885.
11. Koning, B.; Meetsma, A.; Kellogg, R.M. *J. Org. Chem.* **1998**, *63*, 5533.
12. Page, P.C.B.; Heaney, H.; Rassias, A.G.; Reignier, S.; Sampler, E.P.; Talib, S. *A Synlett*. **2000**, *1*, 104.
13. Crossland, R.K.; Noell, J.R. *J. Am Chem. Soc.* **1963**, *85*, 3231.
14. (a) Kelly, R.; van Rheenen, V. *Tetrahedron Lett.* **1973**, 1709.  
(b) Bergmann, M.; Vips, R.; Camacho, F. *Ber.* **1922**, *55*, 2796.  
(c) Zimkin, E.; Bergmann, E. *Rec. Trav. Chim.* **1952**, *71*, 229.
15. Ranaud, P.; Gerster, M. *Angew . Chem. Int. Ed.* **1988**, *37*, 2562.
16. Heaney, H.; Simcox, M.T.; Slawin, A.M.Z.; Giles, R.G. *Synlett*. **1998**, 640.
17. Heaney, H.; Taha, M.O.; Slawin, A.M.Z. *Tetrahedron Lett.* **1977**, *38*, 3051.

18. Cope, A.C.; Hancock, E.A. *J. Am. Chem. Soc.* **1942**, *66*, 1453.
19. Meltsneer, M.; Waldman, E.; Kremer, B. *J. Am. Chem. Soc.* **1940**, *62*, 3494.
20. Suwa, T.; Sugiyama, E.; Shibata, I.; Baba, A. *Synthesis*. **2000**, 789.
21. Mimoun, H.; de Saint Laumer, J.Y.; Giannini, L.; Scopelliti, R.; Floivani, C. *J. Am. Chem. Soc.* **1999**, *121*, 6158.
22. Yamamoto, H.; Maruoka, K. *J. Am. Chem. Soc.* **1981**, *103*, 4186.
23. (a) de Sousa, S.E.; O'Brien, P.; Poumellet, P. *J. Chem. Soc. , Perkin Trans. 1*. **1998**, 1483. (b) de Sousa, S.E.; O'Brien, P. *Tetrahedron Lett.* **1997**, *38*, 4885.
24. Anderson, J.C.; James, S.D.; Mathias, J.V. *Tetrahedron : Asymmetry*. **1988**, 753.
25. Fiore, K.; Martelli, G.; Honari, M.; Savvia, D. *Tetrahedron: Asymmetry*. **1999**, 4887.
26. Neelakantan, L.; Molin-case, J. *J. Org. Chem.* **1971**, *36*, 2261.
27. Robert, B.; Wilson, J.; Laine, M.L. *J. Am. Chem. Soc.* **1985**, *107*, 351.
28. Murahashi, S.; Yoshimura, N.; Tsumiyama, T.; Kojine, T. *J. Am. Chem. Soc.* **1983**, *105*, 927.
29. O'Donnell, M.J.; Polt, R.L. *J. Org. Chem.* **1982**, *47*, 2663.
30. Trost, B.M.; Paterson, D.E. *J. Org. Chem.* **1998**, *63*, 1339.
31. Morimoto, T.; Yamaguchi, Y.; Suzuki, M.; Saitoh, A. *Tetrahedron Lett*, **2000**, *41*, 1025.
32. Hiroi, H.; Suzuki, Y.; Abe, I. *Tetrahedron : Asymmetry*. **1999**, *10*, 1173.
33. Frost, C.G.; Howarth, J.; Williams, J.M.J. *Tetrahedron: Asymmetry*. **1992**, *3*, 1089.

34. Åkermark, B.; Hanson, S.; Vitagliano, A. *J. Am. Chem. Soc.* **1990**, *112*, 4587.
35. Åkermark, B.; Krakenberger, B.; Hanson, S.; Vitagliano, A. *Organometallics*. **1987**, *6*, 620.
36. Åkermark, B.; Hanson, S.; Krakenberger, B.; Vitagliano, A. Zetterberg, K. *Organometallics*. **1984**, *3*, 679.
37. Godleski, S.A.; Villhauer, E.B. *J. Org. Chem.* **1986**, *51*, 486.
38. Zhu, Z.; Lu, X. *Tetrahedron Lett.* **1987**, *28*, 1897.
39. Svenson, M.; Bremberg, V.; Hallman, K.; Csoregh, I.; Moberg, C. *Organometallics*. **1989**, *18*, 4900.
40. Saitoh, A.; Misawa, M.; Morimoto, T. *Synlett*. **1999**, *34*, 483.
41. Moreno, M.; Ribas, J. *Tetrahedron Lett.* **1989**, *30*, 3109.
42. Leggy, A.; Arnold, R.N.; Minnard, J.; Feringa, B.L. *J. Am. Chem. Soc.* **2001**, *123*, 5891.
43. Anderson, J.C.; James, S.D.; Mathias, J.P. *Tetrahedron : Asymmetry*. **1998**, *9*, 753.
44. Feringa, B.L.; Pineschi, M.; Arnold, L.; Imbos, R.; de Vries, A. *Angew. Chem. Int. Ed.* **1997**, *36*, 2620.
45. Alexakis, A.; Vastra, J.; Burton, J.; Benhaim, C.; Mangeney, P. *Tetrahedron Lett.* **1988**, *39*, 7869.
46. Schinnerl, M.; Seitz, M.; Kaiser, A.; Reiser, O. *Organic Letters*. **2001**, *3*.
47. Wu, Y. J.; Yun, H. Y.; Wu, Y. S.; Ding, K. L.; Zhou, Y. *Tetrahedron Asymmetry*. **2000**, *11*, 3543.
48. (a) Morimoto, T.; Yamaguchi, Y.; Suzuki, M.; Saitoh, A. *Tetrahedron Letters*, **2000**, *41*, 10025. (b) Knobel, A.K.H.; Escher, I.H.; Pfaltz, A. *Synlett*. **1997**, 1429.

49. (a) Dieter, K.R.; Tokles, M. *J. Am. Chem. Soc.* **1987**, *109*, 2040. (b) Stewart, J.R.K. ; Lever, J.R.; Whangbo, M.H. *J. Org.Chem.* **1982**,*47*,1472.
50. Houre, H.; Wilkins, S. *J. Org. Chem.* **1978**, *43*, 2443.
51. Degrado, S.J.; Mizutani, H.; Hoveyda, A.H. *J. Am. Chem. Soc.* **2001**, *123*, 755.
52. Anderson, J.C.; Cubbon, R.; Harding, M. *Tetrahedron : Asymmetry.* **1998**, *9*, 3461.
53. Mandoli, A.; Arnold, L.A.; de Vries, A.M.; Salvadori, P.; Feringa, B.L. *Tetrahedron : Asymmetry.* **2001**, *12*, 1924.
54. Krauss, S.R.; Smith, S.G. *J. Am. Chem. Soc.* **1981**, *103*, 141.
55. Nakamura, E.; Mori, S. *Angew. Chem. Int. Ed.* **2000**, *39*, 3750.
56. Torii, S.; Okumoto, H.; Akuhoshi, F.; Kotani, T. *J. Am. Chem. Soc.* **1989**, *111*, 8932.
57. Sato, Y.; Sodeoka, M.; Shibasaki, M. *J. Org. Chem.* **1989**, *54*, 4738.
58. Ozawa, F.; Kubo, A.; Hayashi, T. *Tetrahedron Lett.* **1992**, *11*, 1485.
59. Gilbertson, S; Xie, D. J.; Fu, Z. C. *J. Org. Chem.* **2001**, *66*, 7240.
60. Hennessy, A.J.; Malone, Y.M.; Guiry, P.J. *Tetrahedron Lett.* **1999**, *40*, 9163.
61. Larock, R.; Gung, W.; Parker, B. *Tetrahedron Lett.* **1989**, *30*, 2606.
62. Shibasaki, M.; Boden, C.; Kojima, A. *Tetrahedron* **1997**, *53*, 7371.
63. Young Cho, S.; Shibasaki, M. *Tetrahedron Lett.* **1998**, *39*, 1773.
64. Carmichael, A.J.; Earle, M.J.; Holbrey, J.D.; McCormac, P.B.; Seddon, K.R. *Organic Letters.* **1999**, *2*, 997.
65. Ozawa, F.; Kubo, A.; Matsumoto, Y.; Hayashi, T; Nishioka, E.; Yanagi, K.; Moriguchi, K. *Organometallics.* **1993**, *12*, 4188.


66. Carreno, M.C. *Chem Rev.* **1995**, *95*, 1717.
67. Pitchen, P.; Kagan, H.B.; Deshmukh, M.N.; Dunach, E. *J. Am. Chem. Soc.* **1984**, *106*, 8188.
68. Di Furia, D.; Modena, G.; Seraglia, R. *Synthesis.* **1984**, 325.
69. Davis, F.A.; Nadir, U.K.; Kluger, E.W. *J. Chem. Soc.; Chem. Commun.* **1977**, 25.
70. Page, P.C.B.; Heer, J.P.; Bethell, D.; Collington, E.W.; Andrews, D.M. *Tetrahedron Lett.* **1994**, *35*, 9629.
71. Donnoli, M.; Superchi, S.; Rosini, C. *Tetrahedron Lett.* **1998**, *39*, 8541.
72. Davis, F.A.; McCauley, J.P.; Chattopadhyay, S.; Harakal, M.E. *J. Am Chem. Soc.* **1987**, *109*, 3370.
73. Davis, F.A.; Jenkins, R.H.; Awad, S.B.; Stringer, O.D.; Watson, W.H.; Galloy, J. *J. Am Chem. Soc.* **1982**, *104*, 5412.
74. Davis, F.A.; Thimmareddy, R.; Han, W.; Carroll, P.J. *J. Am. Chem. Soc.* **1992**, *114*, 1428.
75. PhD Thesis. Andrew Lund, Loughborough University **1997**.
76. a) PhD Thesis Gerry Rassias, Loughborough University **1999**; b) PhD Thesis Serge Reignier, Loughborough University **2001**.


# **APPENDICES**

# **APPENDIX 1**


## **nOe Studies on Oxazolidines**

**(4S,5S)-2,3,4-trimethyl-2-(1-methyl ethyl)  
5-phenyl-1,3-oxazolidine**







# SAT 221



# SAT 221





F2 - Acquisition Parameters

Date_	20010604
Time	12.50
INSTRUM	dp400
PROBHD	5 mm Multinu
PULPROG	cosy45
TD	1024
SOLVENT	CDC13
NS	8
DS	16
SWH	3396.739 Hz
FIDRES	3.317128 Hz
AQ	0.1507828 sec
RG	35.9
DW	147.200 usec
DE	7.50 usec
TE	300.0 K
D1	2.00000000 sec
P1	15.00 usec
SFO1	400.1315265 MHz
NUC1	1H
PL1	-6.00 dB
DO	0.00000300 sec
IND	0.00029440 sec

F1 - Acquisition parameters

ND0	1
TD	256
SFO1	400.1315 MHz
FIDRES	13.268512 Hz
SW	8.489 ppm

F2 - Processing parameters


S1	512
SF	400.1300239 MHz
WDW	SINE
SSB	0
LB	0.00 Hz
GB	0
PC	1.00

F1 - Processing parameters

S1	512
MC2	OF
SF	400.1300239 MHz
WDW	SINE
SSB	0
LB	0.00 Hz
GB	0

2D NMR plot parameters

CX2	15.00 cm
CX1	15.00 cm
F2PLO	4.502 ppm
F2LO	1801.20 Hz
F2PH1	0.738 ppm
F2H1	295.22 Hz
F1PLO	4.502 ppm
F1LO	1801.20 Hz
F1PH1	0.738 ppm
F1H1	295.22 Hz
F2PPMCM	0.25091 ppm/cm
F2HZCM	100.39841 Hz/cm
F1PPMCM	0.25091 ppm/cm
F1HZCM	100.39841 Hz/cm



## F2 - Acquisition Parameters

Date\_ 20010604  
 Time 12.50  
 INSTRUM opx400  
 PROBHD 5 mm Multinu  
 PULPROG cosy45  
 TO 1024  
 SOLVENT CDCl<sub>3</sub>  
 NS 8  
 DS 16  
 S<sub>W</sub>H 3395.739 Hz  
 FIDRES 3.317128 Hz  
 AQ 0.1507828 sec  
 RG 35.9  
 D<sub>W</sub> 147.200 usec  
 DE 7.50 usec  
 TE 300.0 K  
 D: 2.00000000 sec  
 P: 15.00 usec  
 SFO1 400.1315265 MHz  
 NUC1 <sup>1</sup>H  
 PL1 -6.00 dB  
 DQ 0.00000300 sec  
 INQ 0.00029440 sec

## F1 - Acquisition parameters

NS 1  
 TO 256  
 SFO1 400.1315 MHz  
 FIDRES 13.268512 Hz  
 S<sub>W</sub> 8.489 ppm

## F2 - Processing parameters


SI 512  
 SF 400.1300239 MHz  
 WD<sub>W</sub> SINE  
 SSB 0  
 LB 0.00 Hz  
 GB 0  
 PC 1.00

## F1 - Processing parameters

SI 512  
 MC2 OF  
 SF 400.1300239 MHz  
 WD<sub>W</sub> SINE  
 SSB 0  
 LB 0.00 Hz  
 GB 0

## 2D NMR plot parameters

CX2 15.00 cm  
 CX1 15.00 cm  
 F2PLO 4.500 ppm  
 F2LO 1800.59 Hz  
 F2PHI 0.750 ppm  
 F2HI 300.10 Hz  
 F1PLO 4.500 ppm  
 F1LO 1800.59 Hz  
 F1PHI 0.750 ppm  
 F1HI 300.10 Hz  
 F2PPMCM 0.25000 ppm/cm  
 F2HZCM 100.03251 Hz/cm  
 F1PPMCM 0.25000 ppm/cm  
 F1HZCM 100.03251 Hz/cm



F2 - Acquisition Parameters

Date_	20010604
Time	12.50
INSTRUM	gpc400
PROBHD	5 mm Multinu
PULPROG	cosy45
TD	1024
SOLVENT	CDCl3
NS	6
DS	16
SWH	3396.739 Hz
FIDRES	3.317126 Hz
AQ	0.1507826 sec
RG	35.9
DW	147.200 usec
DE	7.50 usec
TE	300.0 K
D1	2.0000000 sec
P1	15.00 usec
SFO1	400.1315265 MHz
NUC1	1H
PL1	-6.00 dB
DD	0.00000300 sec
INO	0.0002944 sec

F1 - Acquisition parameters

NO0	!
TD	256
SFO1	400.1315 MHz
FIDRES	13.268512 Hz
SW	8.489 ppm

F2 - Processing parameters


SI	512
SF	400.1300239 MHz
WDW	SINE
SSB	0
LB	0.00 Hz
GB	0
PC	1.00

F1 - Processing parameters


SI	512
MC2	OF
SF	400.1300239 MHz
WDW	SINE
SSB	0
LB	0.00 Hz
GB	0

2D NMR plot parameters

CX2	15.00 cm
CX1	15.00 cm
F2PLO	8.000 ppm
F2LO	3201.03 Hz
F2PH1	-0.489 ppm
F2H1	-195.71 Hz
F1PLO	8.000 ppm
F1LO	3201.03 Hz
F1PH1	-0.489 ppm
F1H1	-195.71 Hz
F2PPMCM	0.56594 ppm/cm
F2HZCM	226.44928 Hz/cm
F1PPMCM	0.56594 ppm/cm
F1HZCM	226.44928 Hz/cm



Cx2	15.00	cm
Cx1	15.00	cm
F2PLO	7.995	ppm
F2LO	3198.87	Hz
F2PHI	-0.495	ppm
F2HI	-197.87	Hz
F1PLO	8.021	ppm
F1LO	3209.26	Hz
F1PHI	-0.469	ppm
F1HI	-187.48	Hz
F2PPMCM	0.56594	ppm/cm
F2HZCM	226.44928	Hz/cm
F1PPMCM	0.56594	ppm/cm
F1HZCM	226.44930	Hz/cm



## F2 - Acquisition Parameters

Date\_ 20010529  
Time 16.12  
INSTRUM dm400  
PROBHD 5 mm Multinu  
PULPROG noesytp  
TO 1024  
SOLVENT CD3CN  
NS 80  
DS 16  
SWH 3396.739 Hz  
FIDRES 3.317128 Hz  
AQ 0.1507828 sec  
RG 43.7  
OW 147.200 usec  
DE 7.50 usec  
TE 300.0 K  
D1 2.00000000 sec  
P1 15.00 usec  
SFO1 400.1315265 MHz  
NUC1 1H  
PL1 -6.00 dB  
D0 2.00000300 sec  
DB 2.99999999 sec  
TD 0.00014720 sec

## F1 - Acquisition Parameters

NOO 2  
TO 256  
SFO1 400.1315 MHz  
FIDRES 13.255512 Hz  
SW 5.489 ppm

## F2 - Processing parameters

SI 1024  
SF 400.1300262 MHz  
WDW SINE  
SSB 2  
LB 3.00 Hz  
GB 0  
PC 1.00


## F1 - Processing parameters

SI 512  
MC2 1PP1  
SF 400.1300262 MHz  
WDW SINE  
SSB 2  
LB 0.00 Hz  
GB 0


## 2D NMR plot parameters

CX2 15.00 cm  
CX1 15.00 cm  
F2PLO 4.612 ppm  
F2LO 1845.33 Hz  
F2PHI 3.591 ppm  
F2HI 236.52 Hz  
F1PLO 4.528 ppm  
F1LO 1851.97 Hz  
F1PHI 0.583 ppm  
F1HI 233.21 Hz  
F2PPMCM 0.26805 ppm/cm  
F2HZCM 107.25391 Hz/cm  
F1PPMCM 0.25971 ppm/cm  
F1HZCM 107.25391 Hz/cm

**(4S,5S)-2,3,4-trimethyl-5-phenyl-2 pyridin yl  
1,3-oxazolidine**



CASS Analysis 0950: Structure and Stereochemistry of ST84254-027A1 (CDC13)



```

Current Data Parameters
NAME          CASE_9858
LIFE         2
NOOBS        1

```

```

F2 - Acquisition Parameters
Date_      20180406
Time       14.51
INSTRUM    OM4400
PROBHD     5mm SBC 17KA
PULPROG    zgpg30
TD         20.48
SOLVENT    CDCl3
AQ         1.00
RG         18
SOLVENT    250.1 95.4 Hz
FIDRES     1.7530040 Hz
AQ         2.2815134 sec
RG         812.7
TD         138,200
DE         1.00
TE         300.2 K
D0         3.0000300 sec
d1         0.9000300 sec
d11        2.0000300 sec
d12        2.0001500 sec
d13        3.0002784 sec

```

```

----- CHANNEL 11 -----
NAME      IM
R2         10.50 USBC
T1         10.50 USBC
R11        7.00 CB
FCL        430.1320007 Mc

```

----- GRADIENT CHANNEL -----  
#16 1500.00 WSEC

```

FI - Acquisition Parameters
1000          1
7             128
SF01         400.132 MHz
FREQ01       28.042144 Hz
Ed           0.077 pcu


```

F2 - Processing parameters	
IT	1024
GF	430.1293906 Hz
WOW	TIME
SSD	G
LB	0.00 Hz
FB	G
PC	1.00

F1 - Processing parameters	
SI	1024
W2	25
3F	400.1292888 Hz
40u	57AC
52u	0
18	0.00 Hz
20	0

2D NMR plot parameters	
COZ	28.00 cm
CH1	28.00 cm
FAPLO	0.000 ppm
FZLO	3601.17 Hz
FPM1	1.000 ppm
FMI	400.13 Hz
FAPLO	0.000 ppm
FZLO	3601.17 Hz
FPM1	1.000 ppm
FMI	400.13 Hz
FAPLO	0.0000 ppm/cm
FZLO	150.05208 Hz/cm
FAPLO	0.00000 ppm/cm
FZLO	150.05208 Hz/cm

CASS Analysis 0950: Structure and Stereochemistry of ST84254-027A1 (CDC13)



Current Data Parameters  
NAME CASS\_0950  
EXPNO 8  
PROCNO 1

F2 - Acquisition Parameters  
Date\_ 20010405  
Time 16.39  
INSTRUM cps400  
PROCNO 5 nu T1 31P  
PULPROG zgpg30  
TD 65536  
SOLVENT DMSO  
NS 512  
DS 4  
SHH 8278.148 Hz  
FIDRES 0.126314 Hz  
AQ 3.9584243 sec  
RG 2048  
DE 60.400 usec  
DEL 8.00 usec  
TE 300.0 K  
D1 2.00000000 sec  
D2 0.50000000 sec  
D1E 0.00010000 sec  
d20 0.24890000 sec


----- CHANNEL f1 -----  
NUC1 31P  
P1 8.40 usec  
D2 15.80 usec  
P12 90000.00 usec  
PL0 120.00 dB  
PL1 0.00 dB  
RF01 400.130158 MHz  
SP1 60.00 dB  
SFOFF1 0.00 Hz  
SFOFF2 0.00 Hz

----- GRADIENT CHANNEL -----  
GPMAX1 sine 100  
GPMAX2 sine 100  
GPMAX3 sine 100  
GPMAX4 sine 100  
GPX1 0.00 %  
GPX2 0.00 %  
GPX3 0.00 %  
GPX4 0.00 %  
GPT1 0.00 %  
GPT2 0.00 %  
GPT3 0.00 %  
GPT4 0.00 %  
GPZ1 51.00 %  
GPZ2 57.00 %  
GPZ3 48.00 %  
GPZ4 -40.00 %  
P16 1000.00 usec

F2 - Processing parameters  
SI 32768  
SF 400.130158 MHz  
WDW EM  
SSB 0  
LB 1.00 Hz  
GB 0  
PC 1.00

10 1000 plot parameters  
C1 30.00 CA  
F1P 0.000 ppm  
F1 3501.37 Hz  
F2P 0.500 ppm  
F2 205.07 Hz  
PPMCH 0.26333 ppm/Hz  
HZCH 113.37017 MHz/Hz


CASS Analysis 0950: Structure and Stereochemistry of ST84254-027A1 (CDC13)




Current Data Parameters  
 NAME: CASS\_0950  
 EXPNO: 1  
 PROCNO: 1

F2 - Acquisition Parameters  
 Date\_: 20190605  
 Time: 11.51  
 INSTRUM: spect  
 PROBR: 5mm QNP 1H/13  
 PULPROG: zgpg30  
 VS: 300.13  
 SOLVENT: CDCl3  
 NS: 1024  
 DS: 4  
 SWH: 2001.854 Hz  
 FIDRES: 0.753864 Hz  
 AQ: 0.2681310 sec  
 RG: 13004  
 CH: 128.000 MHz  
 DE: 5.00 mmol  
 TE: 300.2 K  
 DWT2: 1.06800000  
 GB: 0.0000000 sec  
 HY: 0.0000000 sec  
 HX: 0.0000000 sec  
 GY: 0.0000000 sec  
 GX: 0.0000000 sec  
 Y1: 0.0000000 sec  
 Y2: 0.0000000 sec  
 Y3: 0.0000000 sec  
 Y4: 0.0000000 sec  
 Y5: 0.0000000 sec  
 Y6: 0.0000000 sec  
 Y7: 0.0000000 sec  
 Y8: 0.0000000 sec  
 Y9: 0.0000000 sec  
 Y10: 0.0000000 sec  
 Y11: 0.0000000 sec  
 Y12: 0.0000000 sec  
 Y13: 0.0000000 sec  
 Y14: 0.0000000 sec  
 Y15: 0.0000000 sec  
 Y16: 0.0000000 sec  
 Y17: 0.0000000 sec  
 Y18: 0.0000000 sec  
 Y19: 0.0000000 sec  
 Y20: 0.0000000 sec  
 Y21: 0.0000000 sec  
 Y22: 0.0000000 sec  
 Y23: 0.0000000 sec  
 Y24: 0.0000000 sec  
 Y25: 0.0000000 sec  
 Y26: 0.0000000 sec  
 Y27: 0.0000000 sec  
 Y28: 0.0000000 sec  
 Y29: 0.0000000 sec  
 Y30: 0.0000000 sec  
 Y31: 0.0000000 sec  
 Y32: 0.0000000 sec  
 Y33: 0.0000000 sec  
 Y34: 0.0000000 sec  
 Y35: 0.0000000 sec  
 Y36: 0.0000000 sec  
 Y37: 0.0000000 sec  
 Y38: 0.0000000 sec  
 Y39: 0.0000000 sec  
 Y40: 0.0000000 sec  
 Y41: 0.0000000 sec  
 Y42: 0.0000000 sec  
 Y43: 0.0000000 sec  
 Y44: 0.0000000 sec  
 Y45: 0.0000000 sec  
 Y46: 0.0000000 sec  
 Y47: 0.0000000 sec  
 Y48: 0.0000000 sec  
 Y49: 0.0000000 sec  
 Y50: 0.0000000 sec  
 Y51: 0.0000000 sec  
 Y52: 0.0000000 sec  
 Y53: 0.0000000 sec  
 Y54: 0.0000000 sec  
 Y55: 0.0000000 sec  
 Y56: 0.0000000 sec  
 Y57: 0.0000000 sec  
 Y58: 0.0000000 sec  
 Y59: 0.0000000 sec  
 Y60: 0.0000000 sec  
 Y61: 0.0000000 sec  
 Y62: 0.0000000 sec  
 Y63: 0.0000000 sec  
 Y64: 0.0000000 sec  
 Y65: 0.0000000 sec  
 Y66: 0.0000000 sec  
 Y67: 0.0000000 sec  
 Y68: 0.0000000 sec  
 Y69: 0.0000000 sec  
 Y70: 0.0000000 sec  
 Y71: 0.0000000 sec  
 Y72: 0.0000000 sec  
 Y73: 0.0000000 sec  
 Y74: 0.0000000 sec  
 Y75: 0.0000000 sec  
 Y76: 0.0000000 sec  
 Y77: 0.0000000 sec  
 Y78: 0.0000000 sec  
 Y79: 0.0000000 sec  
 Y80: 0.0000000 sec  
 Y81: 0.0000000 sec  
 Y82: 0.0000000 sec  
 Y83: 0.0000000 sec  
 Y84: 0.0000000 sec  
 Y85: 0.0000000 sec  
 Y86: 0.0000000 sec  
 Y87: 0.0000000 sec  
 Y88: 0.0000000 sec  
 Y89: 0.0000000 sec  
 Y90: 0.0000000 sec  
 Y91: 0.0000000 sec  
 Y92: 0.0000000 sec  
 Y93: 0.0000000 sec  
 Y94: 0.0000000 sec  
 Y95: 0.0000000 sec  
 Y96: 0.0000000 sec  
 Y97: 0.0000000 sec  
 Y98: 0.0000000 sec  
 Y99: 0.0000000 sec  
 Y100: 0.0000000 sec


CASS Analysis 0950: Structure and Stereochemistry of ST84254-027A1 (CDC13)

[illegible]


**(4S,5S)-2,3,4-trimethyl-2,5-diphenyl-  
1,3-oxazolidine**



JS 0992 Stereochemistry of STB4254-006 in CDCl3

[illegible]

S 0992 Stereochemistry of ST84254-006 in CDC13




Source: Bruker Avance	
NAME	100
EXPNO	1
PROCNO	1
F2 - Acquisition Parameters	
DATE_	20070720
TIME	1.00
INSTRUM	QNP400
PROBHD	5 mm QNP 1H/13
PULPROG	zgpg30
TD	32768
F2 - Acquisition Parameters	625.13
Q1	0.00
Q2	0.00
Q3	0.00
Q4	0.00
Q5	0.00
Q6	0.00
Q7	0.00
Q8	0.00
Q9	0.00
Q10	0.00
Q11	0.00
Q12	0.00
Q13	0.00
Q14	0.00
Q15	0.00
Q16	0.00
Q17	0.00
Q18	0.00
Q19	0.00
Q20	0.00
Q21	0.00
Q22	0.00
Q23	0.00
Q24	0.00
Q25	0.00
Q26	0.00
Q27	0.00
Q28	0.00
Q29	0.00
Q30	0.00
Q31	0.00
Q32	0.00
Q33	0.00
Q34	0.00
Q35	0.00
Q36	0.00
Q37	0.00
Q38	0.00
Q39	0.00
Q40	0.00
Q41	0.00
Q42	0.00
Q43	0.00
Q44	0.00
Q45	0.00
Q46	0.00
Q47	0.00
Q48	0.00
Q49	0.00
Q50	0.00
Q51	0.00
Q52	0.00
Q53	0.00
Q54	0.00
Q55	0.00
Q56	0.00
Q57	0.00
Q58	0.00
Q59	0.00
Q60	0.00
Q61	0.00
Q62	0.00
Q63	0.00
Q64	0.00
Q65	0.00
Q66	0.00
Q67	0.00
Q68	0.00
Q69	0.00
Q70	0.00
Q71	0.00
Q72	0.00
Q73	0.00
Q74	0.00
Q75	0.00
Q76	0.00
Q77	0.00
Q78	0.00
Q79	0.00
Q80	0.00
Q81	0.00
Q82	0.00
Q83	0.00
Q84	0.00
Q85	0.00
Q86	0.00
Q87	0.00
Q88	0.00
Q89	0.00
Q90	0.00
Q91	0.00
Q92	0.00
Q93	0.00
Q94	0.00
Q95	0.00
Q96	0.00
Q97	0.00
Q98	0.00
Q99	0.00
Q100	0.00


10 KHz slot parameters	
CS	10.00 GS
F1P	8.000 MHz
F1	1201.84 MHz
F2P	8.000 MHz
F2	0.00 MHz
RFCH	6.20667 GHz/GS
4ZCH	108.76133 MHz/GS

10 top list parameters	
CX	13.00 cd
FIP	0.000 dpu
F1	3201.04 Hz
F2P	0.000 ppm
FZ	0.00 Hz
FXCH	3.26667 pc/c2
YCH	103.75133 Hz/c2

**(4S,5R)-2,3,4-trimethyl-5-phenyl-2-pyridyl  
1,3-oxazolidine**




SS 0991 Stereochemistry of STB4254-002 in COCL3



Current Data Parameters  
NAME STB4254-002  
EXPNO 10  
PROCNO 1

F1 - Acquisition Parameters  
CH1 2001001  
T1 10.00  
SOLVENT 0.00  
PULPROG 1.00  
F1 1.00  
F2 1.00  
F3 1.00  
F4 1.00  
F5 1.00  
F6 1.00  
F7 1.00  
F8 1.00  
F9 1.00  
F10 1.00  
F11 1.00  
F12 1.00  
F13 1.00  
F14 1.00  
F15 1.00  
F16 1.00  
F17 1.00  
F18 1.00  
F19 1.00  
F20 1.00  
F21 1.00  
F22 1.00  
F23 1.00  
F24 1.00  
F25 1.00  
F26 1.00  
F27 1.00  
F28 1.00  
F29 1.00  
F30 1.00  
F31 1.00  
F32 1.00  
F33 1.00  
F34 1.00  
F35 1.00  
F36 1.00  
F37 1.00  
F38 1.00  
F39 1.00  
F40 1.00  
F41 1.00  
F42 1.00  
F43 1.00  
F44 1.00  
F45 1.00  
F46 1.00  
F47 1.00  
F48 1.00  
F49 1.00  
F50 1.00  
F51 1.00  
F52 1.00  
F53 1.00  
F54 1.00  
F55 1.00  
F56 1.00  
F57 1.00  
F58 1.00  
F59 1.00  
F60 1.00  
F61 1.00  
F62 1.00  
F63 1.00  
F64 1.00  
F65 1.00  
F66 1.00  
F67 1.00  
F68 1.00  
F69 1.00  
F70 1.00  
F71 1.00  
F72 1.00  
F73 1.00  
F74 1.00  
F75 1.00  
F76 1.00  
F77 1.00  
F78 1.00  
F79 1.00  
F80 1.00  
F81 1.00  
F82 1.00  
F83 1.00  
F84 1.00  
F85 1.00  
F86 1.00  
F87 1.00  
F88 1.00  
F89 1.00  
F90 1.00  
F91 1.00  
F92 1.00  
F93 1.00  
F94 1.00  
F95 1.00  
F96 1.00  
F97 1.00  
F98 1.00  
F99 1.00  
F100 1.00

ASS 0991 Stereochemistry of ST84254-002 in CDCL3



Current Data Parameters

NAME	VALUE
EXPNO	10
PROCNO	1

F1 - Acquisition Parameters

NAME	VALUE
DATE_	20100811
TIME	15.34
INSTRUM	QNP-400
PROBHD	5 mm QNP 1H/13
PULPROG	zgpg30
TD	65536
SF	400.146
WDW	EM
SSB	0
LB	3.00
GB	0
PC	1.00
RG	327.68
RT	0.00
CT	0.00
TE	300.2
DE	0.00
CH	0.00
SI	327.68
SC	0.00
SD	0.00
SE	0.00
SM	0.00
SN	0.00
SO	0.00
SP	0.00
SR	0.00
SS	0.00
ST	0.00
SV	0.00
SW	0.00
TX	0.00
TY	0.00
TZ	0.00
U1	0.00
U2	0.00
U3	0.00
U4	0.00
U5	0.00
U6	0.00
U7	0.00
U8	0.00
U9	0.00
U0	0.00
V1	0.00
V2	0.00
V3	0.00
V4	0.00
V5	0.00
V6	0.00
V7	0.00
V8	0.00
V9	0.00
V0	0.00
W1	0.00
W2	0.00
W3	0.00
W4	0.00
W5	0.00
W6	0.00
W7	0.00
W8	0.00
W9	0.00
W0	0.00
X1	0.00
X2	0.00
X3	0.00
X4	0.00
X5	0.00
X6	0.00
X7	0.00
X8	0.00
X9	0.00
X0	0.00
Y1	0.00
Y2	0.00
Y3	0.00
Y4	0.00
Y5	0.00
Y6	0.00
Y7	0.00
Y8	0.00
Y9	0.00
Y0	0.00
Z1	0.00
Z2	0.00
Z3	0.00
Z4	0.00
Z5	0.00
Z6	0.00
Z7	0.00
Z8	0.00
Z9	0.00
Z0	0.00

F2 - Processing Parameters

NAME	VALUE
DATE_	20100811
TIME	15.34
INSTRUM	QNP-400
PROBHD	5 mm QNP 1H/13
PULPROG	zgpg30
TD	65536
SF	400.146
WDW	EM
SSB	0
LB	3.00
GB	0
PC	1.00
RG	327.68
RT	0.00
CT	0.00
TE	300.2
DE	0.00
CH	0.00
SI	327.68
SC	0.00
SD	0.00
SE	0.00
SM	0.00
SN	0.00
SO	0.00
SP	0.00
SR	0.00
SS	0.00
ST	0.00
SV	0.00
SW	0.00
TX	0.00
TY	0.00
TZ	0.00
U1	0.00
U2	0.00
U3	0.00
U4	0.00
U5	0.00
U6	0.00
U7	0.00
U8	0.00
U9	0.00
U0	0.00
V1	0.00
V2	0.00
V3	0.00
V4	0.00
V5	0.00
V6	0.00
V7	0.00
V8	0.00
V9	0.00
V0	0.00
W1	0.00
W2	0.00
W3	0.00
W4	0.00
W5	0.00
W6	0.00
W7	0.00
W8	0.00
W9	0.00
W0	0.00
X1	0.00
X2	0.00
X3	0.00
X4	0.00
X5	0.00
X6	0.00
X7	0.00
X8	0.00
X9	0.00
X0	0.00
Y1	0.00
Y2	0.00
Y3	0.00
Y4	0.00
Y5	0.00
Y6	0.00
Y7	0.00
Y8	0.00
Y9	0.00
Y0	0.00
Z1	0.00
Z2	0.00
Z3	0.00
Z4	0.00
Z5	0.00
Z6	0.00
Z7	0.00
Z8	0.00
Z9	0.00
Z0	0.00

F3 - Processing Parameters

NAME	VALUE
DATE_	20100811
TIME	15.34
INSTRUM	QNP-400
PROBHD	5 mm QNP 1H/13
PULPROG	zgpg30
TD	65536
SF	400.146
WDW	EM
SSB	0
LB	3.00
GB	0
PC	1.00
RG	327.68
RT	0.00
CT	0.00
TE	300.2
DE	0.00
CH	0.00
SI	327.68
SC	0.00
SD	0.00
SE	0.00
SM	0.00
SN	0.00
SO	0.00
SP	0.00
SR	0.00
SS	0.00
ST	0.00
SV	0.00
SW	0.00
TX	0.00
TY	0.00
TZ	0.00
U1	0.00
U2	0.00
U3	0.00
U4	0.00
U5	0.00
U6	0.00
U7	0.00
U8	0.00
U9	0.00
U0	0.00
V1	0.00
V2	0.00
V3	0.00
V4	0.00
V5	0.00
V6	0.00
V7	0.00
V8	0.00
V9	0.00
V0	0.00
W1	0.00
W2	0.00
W3	0.00
W4	0.00
W5	0.00
W6	0.00
W7	0.00
W8	0.00
W9	0.00
W0	0.00
X1	0.00
X2	0.00
X3	0.00
X4	0.00
X5	0.00
X6	0.00
X7	0.00
X8	0.00
X9	0.00
X0	0.00
Y1	0.00
Y2	0.00
Y3	0.00
Y4	0.00
Y5	0.00
Y6	0.00
Y7	0.00
Y8	0.00
Y9	0.00
Y0	0.00
Z1	0.00
Z2	0.00
Z3	0.00
Z4	0.00
Z5	0.00
Z6	0.00
Z7	0.00
Z8	0.00
Z9	0.00
Z0	0.00

F4 - Processing Parameters

NAME	VALUE
DATE_	20100811
TIME	15.34
INSTRUM	QNP-400
PROBHD	5 mm QNP 1H/13
PULPROG	zgpg30
TD	65536
SF	400.146
WDW	EM
SSB	0
LB	3.00
GB	0
PC	1.00
RG	327.68
RT	0.00
CT	0.00
TE	300.2
DE	0.00
CH	0.00
SI	327.68
SC	0.00
SD	0.00
SE	0.00
SM	0.00
SN	0.00
SO	0.00
SP	0.00
SR	0.00
SS	0.00
ST	0.00
SV	0.00
SW	0.00
TX	0.00
TY	0.00
TZ	0.00
U1	0.00
U2	0.00
U3	0.00
U4	0.00
U5	0.00
U6	0.00
U7	0.00
U8	0.00
U9	0.00
U0	0.00
V1	0.00
V2	0.00
V3	0.00
V4	0.00
V5	0.00
V6	0.00
V7	0.00
V8	0.00
V9	0.00
V0	0.00
W1	0.00
W2	0.00
W3	0.00
W4	0.00
W5	0.00
W6	0.00
W7	0.00
W8	0.00
W9	0.00
W0	0.00
X1	0.00
X2	0.00
X3	0.00
X4	0.00
X5	0.00
X6	0.00
X7	0.00
X8	0.00
X9	0.00
X0	0.00
Y1	0.00
Y2	0.00
Y3	0.00
Y4	0.00
Y5	0.00
Y6	0.00
Y7	0.00
Y8	0.00
Y9	0.00
Y0	0.00
Z1	0.00
Z2	0.00
Z3	0.00
Z4	0.00
Z5	0.00
Z6	0.00
Z7	0.00
Z8	0.00
Z9	0.00
Z0	0.00


The figure is a 10x10 grid plot. The horizontal axis is labeled 'ppm' and ranges from 0 to 10. The vertical axis is labeled 'ppm' and ranges from 0 to 140. The plot shows a high concentration of particles in the bottom-left corner (low ppm, low ppm) and a few scattered particles in the top-right corner (high ppm, high ppm).

```


DETECT DATA ACQUISITION
=====
DETECT 1
=====
=====
#1 - AMPLITUDE PARAMETERS
=====
DATA 13.00
GAIN 0.0000
0 to 100.000
FREQ 10.000
F2 10.000
F3 10.000
F4 10.000
F5 10.000
F6 10.000
F7 10.000
F8 10.000
F9 10.000
F10 10.000
F11 10.000
F12 10.000
F13 10.000
F14 10.000
F15 10.000
F16 10.000
F17 10.000
F18 10.000
F19 10.000
F20 10.000
F21 10.000
F22 10.000
F23 10.000
F24 10.000
F25 10.000
F26 10.000
F27 10.000
F28 10.000
F29 10.000
F30 10.000
F31 10.000
F32 10.000
F33 10.000
F34 10.000
F35 10.000
F36 10.000
F37 10.000
F38 10.000
F39 10.000
F40 10.000
F41 10.000
F42 10.000
F43 10.000
F44 10.000
F45 10.000
F46 10.000
F47 10.000
F48 10.000
F49 10.000
F50 10.000
F51 10.000
F52 10.000
F53 10.000
F54 10.000
F55 10.000
F56 10.000
F57 10.000
F58 10.000
F59 10.000
F60 10.000
F61 10.000
F62 10.000
F63 10.000
F64 10.000
F65 10.000
F66 10.000
F67 10.000
F68 10.000
F69 10.000
F70 10.000
F71 10.000
F72 10.000
F73 10.000
F74 10.000
F75 10.000
F76 10.000
F77 10.000
F78 10.000
F79 10.000
F80 10.000
F81 10.000
F82 10.000
F83 10.000
F84 10.000
F85 10.000
F86 10.000
F87 10.000
F88 10.000
F89 10.000
F90 10.000
F91 10.000
F92 10.000
F93 10.000
F94 10.000
F95 10.000
F96 10.000
F97 10.000
F98 10.000
F99 10.000
F100 10.000
F101 10.000
F102 10.000
F103 10.000
F104 10.000
F105 10.000
F106 10.000
F107 10.000
F108 10.000
F109 10.000
F110 10.000
F111 10.000
F112 10.000
F113 10.000
F114 10.000
F115 10.000
F116 10.000
F117 10.000
F118 10.000
F119 10.000
F120 10.000
F121 10.000
F122 10.000
F123 10.000
F124 10.000
F125 10.000
F126 10.000
F127 10.000
F128 10.000
F129 10.000
F130 10.000
F131 10.000
F132 10.000
F133 10.000
F134 10.000
F135 10.000
F136 10.000
F137 10.000
F138 10.000
F139 10.000
F140 10.000
F141 10.000
F142 10.000
F143 10.000
F144 10.000
F145 10.000
F146 10.000
F147 10.000
F148 10.000
F149 10.000
F150 10.000
F151 10.000
F152 10.000
F153 10.000
F154 10.000
F155 10.000
F156 10.000
F157 10.000
F158 10.000
F159 10.000
F160 10.000
F161 10.000
F162 10.000
F163 10.000
F164 10.000
F165 10.000
F166 10.000
F167 10.000
F168 10.000
F169 10.000
F170 10.000
F171 10.000
F172 10.000
F173 10.000
F174 10.000
F175 10.000
F176 10.000
F177 10.000
F178 10.000
F179 10.000
F180 10.000
F181 10.000
F182 10.000
F183 10.000
F184 10.000
F185 10.000
F186 10.000
F187 10.000
F188 10.000
F189 10.000
F190 10.000
F191 10.000
F192 10.000
F193 10.000
F194 10.000
F195 10.000
F196 10.000
F197 10.000
F198 10.000
F199 10.000
F200 10.000
F201 10.000
F202 10.000
F203 10.000
F204 10.000
F205 10.000
F206 10.000
F207 10.000
F208 10.000
F209 10.000
F210 10.000
F211 10.000
F212 10.000
F213 10.000
F214 10.000
F215 10.000
F216 10.000
F217 10.000
F218 10.000
F219 10.000
F220 10.000
F221 10.000
F222 10.000
F223 10.000
F224 10.000
F225 10.000
F226 10.000
F227 10.000
F228 10.000
F229 10.000
F230 10.000
F231 10.000
F232 10.000
F233 10.000
F234 10.000
F235 10.000
F236 10.000
F237 10.000
F238 10.000
F239 10.000
F240 10.000
F241 10.000
F242 10.000
F243 10.000
F244 10.000
F245 10.000
F246 10.000
F247 10.000
F248 10.000
F249 10.000
F250 10.000
F251 10.000
F252 10.000
F253 10.000
F254 10.000
F255 10.000
F256 10.000
F257 10.000
F258 10.000
F259 10.000
F260 10.000
F261 10.000
F262 10.000
F263 10.000
F264 10.000
F265 10.000
F266 10.000
F267 10.000
F268 10.000
F269 10.000
F270 10.000
F271 10.000
F272 10.000
F273 10.000
F274 10.000
F275 10.000
F276 10.000
F277 10.000
F278 10.000
F279 10.000
F280 10.000
F281 10.000
F282 10.000
F283 10.000
F284 10.000
F285 10.000
F286 10.000
F287 10.000
F288 10.000
F289 10.000
F290 10.000
F291 10.000
F292 10.000
F293 10.000
F294 10.000
F295 10.000
F296 10.000
F297 10.000
F298 10.000
F299 10.000
F300 10.000
F301 10.000
F302 10.000
F303 10.000
F304 10.000
F305 10.000
F306 10.000
F307 10.000
F308 10.000
F309 10.000
F310 10.000
F311 10.000
F312 10.000
F313 10.000
F314 10.000
F315 10.000
F316 10.000
F317 10.000
F318 10.000
F319 10.000
F320 10.000
F321 10.000
F322 10.000
F323 10.000
F324 10.000
F325 10.000
F326 10.000
F327 10.000
F328 10.000
F329 10.000
F330 10.000
F331 10.000
F332 10.000
F333 10.000
F334 10.000
F335 10.000
F336 10.000
F337 10.000
F338 10.000
F339 10.000
F340 10.000
F341 10.000
F342 10.000
F343 10.000
F344 10.000
F345 10.000
F346
```

11/11/2001  
 11/11/2001  
 11/11/2001

**(4S,5R)-2,3,4-trimethyl-2-(1-methyl ethyl)  
5-phenyl-1,3-oxazolidine**




S 0992 Stereochemistry of ST84254-006 in CDCl3

[illegible]



CASS 0992 Stereochemistry of ST84254-006 in CDCL3



Current Data Parameters  
NAME CASS\_0992  
EXPER 3  
PROCNO 1

F2 - Acquisition Parameters  
Date\_ 20010425  
Time 0.12  
INSTRUM dx-400  
PROBHD 5 CD T1 31P  
PULPROG selmgp.3  
TD 65536  
SOLVENT CDCL3  
NS 150  
DS 4  
SWH 6278.140 Hz  
FIDRES 0.123314 Hz  
AQ 3.5564243 sec  
RG 512  
OU 80.000 usec  
CZ 8.00 usec  
TE 323.5 K  
D1 2.0000000 sec  
C8 0.2000000 sec  
D18 0.0001000 sec  
S20 0.2000000 sec

===== CHANNEL f1 =====  
NUC1 1H  
P1 7.00 usec  
a2 13.00 usec  
P12 20000.00 usec  
PL0 120.00 dB  
PL1 0.00 dB  
SFO1 400.1300000 MHz  
SP1 09.00 dB  
SFOA01 90.00 MHz  
SCFF1 0.00 Hz

===== GRADIENT CHANNEL =====  
SFOA01 90.00 MHz  
SFOA02 90.00 MHz  
SFOA03 90.00 MHz  
SFOA04 90.00 MHz  
GPR1 0.00 Hz  
GPR2 0.00 Hz  
GPR3 0.00 Hz  
GPR4 0.00 Hz  
GPR5 0.00 Hz  
GPR6 0.00 Hz  
GPR7 0.00 Hz  
GPR8 0.00 Hz  
GPR9 13.00 Hz  
GPR10 17.00 Hz  
GPR11 40.00 Hz  
GPR12 -40.00 Hz  
P15 1000.00 usec

F2 - Processing parameters  
SI 32768  
SF 400.1300000 MHz  
WDW EM  
SSB 0  
LB 1.00 Hz  
GB 0  
PC 1.00

1D NMR plot parameters  
CH 30.00 cm  
FIP 0.000 sec  
FI 3201.04 Hz  
F2P 0.000 sec  
F2 0.00 Hz  
F2CH 0.2000000 sec/CD  
A2CH 100.70133 Hz/CD

CASS 0992 Stereochemistry of ST84254-006 in CDCL3

Current Data Parameters  
NAME CASS\_0992  
EXPRO 2  
PROCNO 1


F2 - Acquisition Parameters  
Date\_ 20010425  
Time 8.57  
INSTRUM spect  
PROBHD 5 mm T1 31P  
PULPROG zgpg30  
TD 65536  
SOLVENT CDCl3  
NS 84  
DS 4  
SWH 6278.148 Hz  
FIDRES 0.126314 Hz  
AQ 3.9584213 sec  
RG 512  
DQ 80.400 usec  
ZC 0.00 usec  
TE 300.0 K  
D1 2.00000000 sec  
C3 0.55000000 sec  
D18 0.00010000 sec  
D20 0.20000000 sec

===== CHANNEL f1 =====  
NUC1 31P  
P1 7.00 usec  
P2 15.20 usec  
P12 5000.00 usec  
PL0 120.00 dB  
PL1 0.00 dB  
SFO1 400.1326327 MHz  
SP1 60.00 dB  
SFOH1 99.91 MHz  
SFOF1 0.00 MHz

===== GRADIENT CHANNEL =====  
GPMAX1 500.00  
GPMAX2 500.00  
GPMAX3 500.00  
GPMAX4 500.00  
GPK1 0.00 G  
GPK2 0.00 G  
GPK3 0.00 G  
GPK4 0.00 G  
GPY1 0.00 G  
GPY2 0.00 G  
GPY3 0.00 G  
GPY4 0.00 G  
GPT1 11.00 G  
GPT2 17.00 G  
GPT3 40.00 G  
GPT4 -40.00 G  
P10 1000.00 usec


F2 - Processing parameters  
SI 32768  
SF 400.1300024 MHz  
WDW EM  
SSB 0  
LB 1.00 Hz  
GB 0  
PC 1.00

1D NMR list parameters  
CH 15.00 cm  
F1P 0.000 mm  
F1 1201.84 Hz  
F2P 0.000 mm  
F2 0.00 Hz  
FPCW 3.26687 sec/c3  
FCH 105.79133 Hz/c3



## APPENDIX 2

**Methyl (2S)-2-[diphenyl methyldene)  
amino]-4-methyl thiol butanoate**






Table 1. Crystal data and structure refinement for pcb1.

Identification code	pcb1
Chemical formula	C <sub>19</sub> H <sub>21</sub> NO <sub>2</sub> S
Formula weight	327.43
Temperature	150(2) K
Radiation, wavelength	MoK $\alpha$ , 0.71073 Å
Crystal system, space group	monoclinic, P2 <sub>1</sub>
Unit cell parameters	a = 9.8599(7) Å $\alpha$ = 90° b = 9.0747(7) Å $\beta$ = 101.609(2)° c = 10.1733(7) Å $\gamma$ = 90°
Cell volume	891.64(11) Å <sup>3</sup>
Z	2
Calculated density	1.220 g/cm <sup>3</sup>
Absorption coefficient $\mu$	0.190 mm <sup>-1</sup>
F(000)	348
Crystal colour and size	colourless, 0.93 × 0.79 × 0.55 mm <sup>3</sup>
Reflections for cell refinement	6985 ( $\theta$ range 2.63 to 28.60°)
Data collection method	Bruker SMART 1000 CCD diffractometer $\omega$ -rotation with narrow frames
$\theta$ range for data collection	2.04 to 28.60°
Index ranges	h -12 to 12, k -11 to 11, l -13 to 13
Completeness to $\theta$ = 26.00°	99.6 %
Intensity decay	0%
Reflections collected	7527
Independent reflections	3942 ( $R_{int}$ = 0.0125)
Reflections with $F^2 > 2\sigma$	3854
Absorption correction	semi-empirical from equivalents
Min. and max. transmission	0.843 and 0.903
Structure solution	direct methods
Refinement method	Full-matrix least-squares on $F^2$
Weighting parameters a, b	0.0572, 0.1265
Data / restraints / parameters	3942 / 1 / 211
Final R indices [ $F^2 > 2\sigma$ ]	R1 = 0.0311, wR2 = 0.0842
R indices (all data)	R1 = 0.0317, wR2 = 0.0846
Goodness-of-fit on $F^2$	1.030
Absolute structure parameter	-0.03(6)
Extinction coefficient	0.009(3)
Largest and mean shift/su	0.001 and 0.000
Largest diff. peak and hole	0.381 and -0.200 e Å <sup>-3</sup>

Table 2. Atomic coordinates and equivalent isotropic displacement parameters ( $\text{\AA}^2$ ) for pcb1.  $U_{eq}$  is defined as one third of the trace of the orthogonalized  $U^{\mu}$  tensor.

	x	y	z	$U_{eq}$
N(1)	0.83218(11)	0.48875(12)	0.64871(11)	0.0248(2)
C(2)	0.85380(12)	0.33241(15)	0.62404(12)	0.0249(2)
C(3)	0.88564(13)	0.31405(16)	0.48399(13)	0.0284(2)
C(4)	0.76099(16)	0.35105(18)	0.37638(14)	0.0362(3)
S(1)	0.78674(4)	0.32516(5)	0.20723(3)	0.04072(11)
C(5)	0.7977(2)	0.1278(2)	0.1981(2)	0.0524(5)
C(6)	0.97535(14)	0.27720(15)	0.72971(15)	0.0284(3)
O(1)	1.07541(11)	0.21585(15)	0.70680(12)	0.0425(3)
O(2)	0.95356(11)	0.30059(15)	0.85253(11)	0.0436(3)
C(7)	1.0596(2)	0.2458(3)	0.96163(19)	0.0561(5)
C(8)	0.72551(13)	0.53156(13)	0.69198(12)	0.0217(2)
C(9)	0.71317(13)	0.69263(14)	0.71632(13)	0.0227(2)
C(10)	0.81242(15)	0.78902(16)	0.68554(16)	0.0328(3)
C(11)	0.80393(17)	0.93833(18)	0.70879(19)	0.0415(4)
C(12)	0.69649(17)	0.99478(17)	0.76277(18)	0.0383(3)
C(13)	0.59750(16)	0.90065(16)	0.79336(17)	0.0332(3)
C(14)	0.60447(14)	0.75031(15)	0.76987(15)	0.0284(3)
C(15)	0.61405(12)	0.43313(14)	0.72357(13)	0.0227(2)
C(16)	0.50245(15)	0.38865(17)	0.62460(15)	0.0314(3)
C(17)	0.40009(14)	0.29896(19)	0.65773(15)	0.0375(3)
C(18)	0.40759(14)	0.25346(16)	0.78822(16)	0.0330(3)
C(19)	0.51787(14)	0.29726(15)	0.88694(14)	0.0310(3)
C(20)	0.62088(14)	0.38714(15)	0.85487(14)	0.0263(3)

Table 3. Bond lengths [Å] and angles [°] for pcbI.

N(1)–C(8)	1.2796(16)	N(1)–C(2)	1.4638(18)
C(2)–C(6)	1.5246(18)	C(2)–C(3)	1.5286(17)
C(3)–C(4)	1.5101(19)	C(4)–S(1)	1.8046(14)
S(1)–C(5)	1.798(2)	C(6)–O(1)	1.1959(17)
C(6)–O(2)	1.3268(18)	O(2)–C(7)	1.451(2)
C(8)–C(9)	1.4915(17)	C(8)–C(15)	1.5003(17)
C(9)–C(10)	1.3943(18)	C(9)–C(14)	1.3980(17)
C(10)–C(11)	1.381(2)	C(11)–C(12)	1.386(2)
C(12)–C(13)	1.379(2)	C(13)–C(14)	1.3891(19)
C(15)–C(20)	1.3881(18)	C(15)–C(16)	1.3934(19)
C(16)–C(17)	1.390(2)	C(17)–C(18)	1.378(2)
C(18)–C(19)	1.382(2)	C(19)–C(20)	1.3916(18)
C(8)–N(1)–C(2)	120.76(11)	N(1)–C(2)–C(6)	108.58(10)
N(1)–C(2)–C(3)	109.26(11)	C(6)–C(2)–C(3)	109.98(10)
C(4)–C(3)–C(2)	111.20(10)	C(3)–C(4)–S(1)	114.34(9)
C(5)–S(1)–C(4)	101.56(9)	O(1)–C(6)–O(2)	123.66(14)
O(1)–C(6)–C(2)	125.30(13)	O(2)–C(6)–C(2)	110.99(11)
C(6)–O(2)–C(7)	115.95(13)	N(1)–C(8)–C(9)	117.05(11)
N(1)–C(8)–C(15)	125.53(11)	C(9)–C(8)–C(15)	117.40(11)
C(10)–C(9)–C(14)	118.71(12)	C(10)–C(9)–C(8)	119.55(11)
C(14)–C(9)–C(8)	121.74(11)	C(11)–C(10)–C(9)	120.51(13)
C(10)–C(11)–C(12)	120.54(14)	C(13)–C(12)–C(11)	119.52(14)
C(12)–C(13)–C(14)	120.52(14)	C(13)–C(14)–C(9)	120.20(13)
C(20)–C(15)–C(16)	119.12(12)	C(20)–C(15)–C(8)	119.30(11)
C(16)–C(15)–C(8)	121.58(11)	C(17)–C(16)–C(15)	120.05(13)
C(18)–C(17)–C(16)	120.55(13)	C(17)–C(18)–C(19)	119.73(12)
C(18)–C(19)–C(20)	120.19(13)	C(15)–C(20)–C(19)	120.36(12)

Table 4. Anisotropic displacement parameters ( $\text{\AA}^2$ ) for pcb1. The anisotropic displacement factor exponent takes the form:  $-2\pi^2[h^2a^{*2}U^{11} + \dots + 2hka^*b^*U^{12}]$


	$U^{11}$	$U^{22}$	$U^{33}$	$U^{23}$	$U^{13}$	$U^{12}$
N(1)	0.0240(5)	0.0248(5)	0.0274(5)	-0.0005(4)	0.0094(4)	-0.0004(4)
C(2)	0.0226(5)	0.0247(6)	0.0291(6)	-0.0020(5)	0.0092(4)	-0.0010(5)
C(3)	0.0261(6)	0.0312(6)	0.0297(6)	-0.0018(5)	0.0099(4)	0.0031(5)
C(4)	0.0349(7)	0.0467(9)	0.0285(6)	0.0003(6)	0.0099(5)	0.0123(6)
S(1)	0.0419(2)	0.0538(2)	0.02698(17)	-0.00052(16)	0.00826(13)	0.00927(18)
C(5)	0.0538(11)	0.0485(11)	0.0534(11)	-0.0166(8)	0.0077(9)	0.0028(8)
C(6)	0.0236(6)	0.0264(6)	0.0356(7)	0.0004(5)	0.0068(5)	-0.0038(5)
O(1)	0.0275(5)	0.0559(7)	0.0437(6)	-0.0032(5)	0.0065(4)	0.0108(5)
O(2)	0.0357(5)	0.0635(8)	0.0315(5)	0.0041(5)	0.0067(4)	0.0129(5)
C(7)	0.0471(10)	0.0807(15)	0.0363(9)	0.0089(9)	-0.0012(7)	0.0128(10)
C(8)	0.0215(6)	0.0250(6)	0.0192(5)	0.0008(4)	0.0050(4)	-0.0013(4)
C(9)	0.0214(5)	0.0248(6)	0.0224(6)	0.0007(4)	0.0055(4)	-0.0004(5)
C(10)	0.0312(7)	0.0271(7)	0.0451(8)	-0.0007(5)	0.0200(6)	-0.0013(5)
C(11)	0.0401(8)	0.0278(7)	0.0637(11)	-0.0007(7)	0.0274(8)	-0.0060(6)
C(12)	0.0447(9)	0.0232(6)	0.0519(9)	-0.0006(6)	0.0215(7)	0.0016(6)
C(13)	0.0329(7)	0.0304(7)	0.0407(8)	0.0017(6)	0.0179(6)	0.0054(5)
C(14)	0.0266(6)	0.0283(6)	0.0331(7)	0.0018(5)	0.0127(5)	0.0003(5)
C(15)	0.0199(5)	0.0241(5)	0.0258(6)	-0.0015(5)	0.0087(4)	0.0001(4)
C(16)	0.0272(6)	0.0410(7)	0.0259(6)	-0.0014(5)	0.0055(5)	-0.0041(5)
C(17)	0.0238(6)	0.0484(9)	0.0396(8)	-0.0067(6)	0.0042(5)	-0.0094(6)
C(18)	0.0247(6)	0.0317(7)	0.0468(8)	-0.0013(6)	0.0169(6)	-0.0039(5)
C(19)	0.0320(6)	0.0330(7)	0.0323(7)	0.0035(5)	0.0164(5)	-0.0005(5)
C(20)	0.0249(6)	0.0296(6)	0.0250(6)	-0.0006(5)	0.0067(5)	-0.0022(5)

Table 5. Hydrogen coordinates and isotropic displacement parameters ( $\text{\AA}^2$ ) for pcb1.



	x	y	z	U
H(2)	0.7685	0.2755	0.6304	0.030
H(3A)	0.9636	0.3794	0.4748	0.034
H(3B)	0.9142	0.2111	0.4722	0.034
H(4A)	0.7352	0.4551	0.3873	0.043
H(4B)	0.6823	0.2889	0.3896	0.043
H(5A)	0.8855	0.0944	0.2534	0.079
H(5B)	0.7932	0.0980	0.1048	0.079
H(5C)	0.7205	0.0833	0.2313	0.079
H(7A)	1.0747	0.1406	0.9482	0.084
H(7B)	1.0296	0.2600	1.0469	0.084
H(7C)	1.1460	0.2998	0.9635	0.084
H(10)	0.8864	0.7517	0.6483	0.039
H(11)	0.8723	1.0029	0.6876	0.050
H(12)	0.6911	1.0976	0.7786	0.046
H(13)	0.5240	0.9389	0.8308	0.040
H(14)	0.5352	0.6865	0.7903	0.034
H(16)	0.4963	0.4197	0.5344	0.038
H(17)	0.3243	0.2688	0.5898	0.045
H(18)	0.3372	0.1922	0.8102	0.040
H(19)	0.5234	0.2659	0.9770	0.037
H(20)	0.6963	0.4172	0.9232	0.032

## APPENDIX 3

**Methyl (2S)-2-[(E)-2-methyl-1-phenyl propylidene  
amino]-4-(methyl thiol) butanoate**



NCE



Current Data Parameters  
NAME J11  
EXPNO 10  
PROCNO 1

F2 - Acquisition Parameters  
Date\_ 20010308  
Time 10.57  
INSTRUM spect  
PROBHD 5 mm TCI 31P  
PULPROG zgpg30  
TD 65536  
SOLVENT DMSO  
NS 38  
DS 4  
SWH 8278.148 Hz  
FIDRES 0.125314 Hz  
AQ 3.9584247 sec  
RG 2548  
SN 40.400 usec  
SE 8.00 usec  
TE 300.0 K  
D1 2.0000000 sec  
D2 0.5000000 sec  
D18 2.0000000 sec  
D20 0.2480000 sec


----- CHANNEL f1 -----  
NUC1 31P  
P1 7.50 usec  
P2 13.20 usec  
P3 30000.00 usec  
PL1 120.00 dB  
PL2 0.00 dB  
SFO1 400.1312845 MHz  
TO1 10.00 sec  
SFOH1 30000.00 MHz  
SFOFF1 0.00 MHz

----- GRADIENT CHANNEL -----  
GPMAX1 sine.100  
GPMAX2 sine.100  
GPMAX3 sine.100  
GPMAX4 sine.100  
GP11 0.00 %  
GP12 0.00 %  
GP13 0.00 %  
GP14 0.00 %  
GP15 0.00 %  
GP16 0.00 %  
GP17 0.00 %  
GP18 0.00 %  
GP19 0.00 %  
GP20 0.00 %  
GP21 11.00 %  
GP22 17.00 %  
GP23 40.00 %  
GP24 40.00 %  
G16 1000.00 usec

F2 - Processing parameters  
SI 32768  
SF 400.1308824 MHz  
WDW EM  
SSB 0  
GB 1.00 Hz  
DB 0  
RC 1.00

1D post proc parameters  
GX 30.00 sec  
FIP 10.000 sec  
F1 4001.30 Hz  
F2 1.000 sec  
F3 400.13 Hz  
SNUCH 0.30000 sec/cm  
-ZCN 120.02500 Hz/cm

JL



Current Data Parameters  
 NAME JL11  
 EXPNO 1  
 PROCNO 1

F2 - Acquisition Parameters  
 Date\_ 20010308  
 Time 11.03  
 INSTRUM drx400  
 PROBHD 5 mm TXI 31P  
 PULPROG zg30  
 TO 65536  
 SOLVENT CDCl3  
 NS 9  
 DS 2  
 SWH 8278.146 Hz  
 FIDRES 0.126314 Hz  
 AQ 3.5564243 sec  
 RG 32  
 CW 50.400 usec  
 DE 5.00 usec  
 TE 300.0 K  
 SI 2.00000000 sec


----- CHANNEL f1 -----  
 NUC1 1H  
 P1 8.40 usec  
 PL1 0.00 dB  
 SFO1 400.1324710 MHz

F2 - Processing parameters  
 SI 32768  
 SF 400.1300528 MHz  
 WDW EM  
 SSB 0  
 LB 0.30 Hz  
 GB 0  
 PC 1.00

1D NMR plot parameters  
 CX 30.00 cm  
 FIP 11.000 ppm  
 F1 4401.43 Hz  
 F2P 0.000 ppm  
 F2 0.00 Hz  
 DPMCH 0.36667 ppm/cm  
 FZCH 146.71434 Hz/cm

## APPENDIX 4

(4S,5S)-2-[2-(1,1-diphenyl phosphenyl)  
3,4-dimethyl-5-phenyl-1,3-oxazolidine






Table 1. Crystal data and structure refinement for pcbp3.

Identification code	pcbp3	
Chemical formula	$C_{29}H_{28}NOP$	
Formula weight	437.49	
Temperature	150(2) K	
Radiation, wavelength	MoK $\alpha$ , 0.71073 Å	
Crystal system, space group	monoclinic, $P2_1$	
Unit cell parameters	$a = 7.2969(5)$ Å	$\alpha = 90^\circ$
	$b = 10.9083(7)$ Å	$\beta = 95.113(2)^\circ$
	$c = 15.2493(10)$ Å	$\gamma = 90^\circ$
Cell volume	1208.97(14) Å <sup>3</sup>	
Z	2	
Calculated density	1.202 g/cm <sup>3</sup>	
Absorption coefficient $\mu$	0.135 mm <sup>-1</sup>	
$F(000)$	464	
Crystal colour and size	colourless, 0.59 × 0.32 × 0.30 mm <sup>3</sup>	
Reflections for cell refinement	8635 ( $\theta$ range 1.29 to 28.61°)	
Data collection method	Bruker SMART 1000 CCD diffractometer $\omega$ rotation with narrow frames	
$\theta$ range for data collection	1.34 to 28.77°	
Index ranges	$h -9$ to 9, $k -14$ to 14, $l -19$ to 20	
Completeness to $\theta = 26.00^\circ$	100.0 %	
Intensity decay	0%	
Reflections collected	10256	
Independent reflections	5378 ( $R_{int} = 0.0245$ )	
Reflections with $F^2 > 2\sigma$	5093	
Absorption correction	semi-empirical from equivalents	
Min. and max. transmission	0.925 and 0.961	
Structure solution	direct methods	
Refinement method	Full-matrix least-squares on $F^2$	
Weighting parameters $a, b$	0.0787, 0.1322	
Data / restraints / parameters	5378 / 1 / 291	
Final R indices [ $F^2 > 2\sigma$ ]	$R1 = 0.0408$ , $wR2 = 0.1073$	
R indices (all data)	$R1 = 0.0425$ , $wR2 = 0.1090$	
Goodness-of-fit on $F^2$	1.030	
Absolute structure parameter	-0.07(7)	
Largest and mean shift/su	0.001 and 0.000	
Largest diff. peak and hole	0.367 and -0.333 e Å <sup>-3</sup>	

Table 2. Atomic coordinates and equivalent isotropic displacement parameters ( $\text{\AA}^2$ ) for pcbp3.  $U_{\text{eq}}$  is defined as one third of the trace of the orthogonalized  $U^{\text{ij}}$  tensor.

	x	y	z	$U_{\text{eq}}$
O(1)	0.77962(16)	0.64964(15)	0.58090(8)	0.0357(3)
C(2)	0.8328(2)	0.65321(18)	0.49268(11)	0.0282(3)
C(3)	1.0265(2)	0.7090(2)	0.50152(11)	0.0324(4)
N(4)	1.09393(19)	0.66605(17)	0.58954(10)	0.0331(3)
C(5)	0.9358(2)	0.67950(18)	0.64062(11)	0.0293(4)
C(6)	0.6968(2)	0.72449(18)	0.43251(11)	0.0312(4)
C(7)	0.6252(3)	0.8350(2)	0.45898(16)	0.0449(5)
C(8)	0.5075(3)	0.9026(3)	0.3994(2)	0.0664(9)
C(9)	0.4626(3)	0.8587(3)	0.3150(2)	0.0688(9)
C(10)	0.5314(3)	0.7497(3)	0.28944(14)	0.0590(7)
C(11)	0.6482(3)	0.6819(2)	0.34791(13)	0.0419(5)
C(12)	1.1449(3)	0.6666(3)	0.43057(14)	0.0519(6)
C(13)	1.2555(3)	0.7318(3)	0.62786(14)	0.0522(7)
C(14)	0.9448(2)	0.59548(16)	0.71993(10)	0.0257(3)
C(15)	1.0545(2)	0.49067(17)	0.72113(11)	0.0299(3)
C(16)	1.0667(2)	0.41161(16)	0.79257(12)	0.0299(3)
C(17)	0.9683(2)	0.43636(16)	0.86440(12)	0.0285(3)
C(18)	0.8584(2)	0.54093(15)	0.86409(11)	0.0260(3)
C(19)	0.8456(2)	0.62188(15)	0.79308(11)	0.0235(3)
P(1)	0.69434(5)	0.75805(4)	0.78798(2)	0.02506(11)
C(20)	0.5830(2)	0.74279(16)	0.89086(11)	0.0254(3)
C(21)	0.4401(2)	0.65714(18)	0.89216(12)	0.0314(4)
C(22)	0.3447(2)	0.64429(19)	0.96677(13)	0.0353(4)
C(23)	0.3862(3)	0.71889(19)	1.03921(13)	0.0367(4)
C(24)	0.5253(3)	0.8051(2)	1.03836(12)	0.0355(4)
C(25)	0.6243(2)	0.81640(17)	0.96446(11)	0.0296(3)
C(26)	0.8612(2)	0.88031(16)	0.81642(11)	0.0251(3)
C(27)	0.8157(2)	0.99829(17)	0.78670(11)	0.0301(3)
C(28)	0.9335(3)	1.09611(17)	0.80809(13)	0.0344(4)
C(29)	1.1002(3)	1.07663(19)	0.85779(13)	0.0364(4)
C(30)	1.1469(3)	0.95938(19)	0.88650(14)	0.0386(4)
C(31)	1.0291(3)	0.86236(17)	0.86596(13)	0.0339(4)

Table 3. Bond lengths [Å] and angles [°] for pcbp3.

O(1)–C(5)	1.432(2)	O(1)–C(2)	1.434(2)
C(2)–C(6)	1.506(2)	C(2)–C(3)	1.533(2)
C(3)–N(4)	1.465(2)	C(3)–C(12)	1.516(3)
N(4)–C(5)	1.456(2)	N(4)–C(13)	1.457(3)
C(5)–C(14)	1.514(2)	C(6)–C(11)	1.387(3)
C(6)–C(7)	1.388(3)	C(7)–C(8)	1.403(3)
C(8)–C(9)	1.385(5)	C(9)–C(10)	1.361(5)
C(10)–C(11)	1.391(3)	C(14)–C(15)	1.395(2)
C(14)–C(19)	1.413(2)	C(15)–C(16)	1.386(3)
C(16)–C(17)	1.389(3)	C(17)–C(18)	1.394(2)
C(18)–C(19)	1.394(2)	C(19)–P(1)	1.8481(17)
P(1)–C(26)	1.8319(18)	P(1)–C(20)	1.8374(17)
C(20)–C(25)	1.391(2)	C(20)–C(21)	1.402(2)
C(21)–C(22)	1.393(3)	C(22)–C(23)	1.384(3)
C(23)–C(24)	1.385(3)	C(24)–C(25)	1.397(2)
C(26)–C(31)	1.394(2)	C(26)–C(27)	1.395(2)
C(27)–C(28)	1.391(3)	C(28)–C(29)	1.391(3)
C(29)–C(30)	1.385(3)	C(30)–C(31)	1.382(3)
C(5)–O(1)–C(2)	108.89(12)	O(1)–C(2)–C(6)	111.52(14)
O(1)–C(2)–C(3)	104.65(14)	C(6)–C(2)–C(3)	113.63(15)
N(4)–C(3)–C(12)	112.98(16)	N(4)–C(3)–C(2)	100.69(13)
C(12)–C(3)–C(2)	113.33(18)	C(5)–N(4)–C(13)	112.79(16)
C(5)–N(4)–C(3)	103.91(13)	C(13)–N(4)–C(3)	113.83(16)
O(1)–C(5)–N(4)	105.11(13)	O(1)–C(5)–C(14)	110.22(14)
N(4)–C(5)–C(14)	112.73(15)	C(11)–C(6)–C(7)	119.3(2)
C(11)–C(6)–C(2)	119.57(18)	C(7)–C(6)–C(2)	121.04(18)
C(6)–C(7)–C(8)	119.5(2)	C(9)–C(8)–C(7)	120.0(3)
C(10)–C(9)–C(8)	120.4(2)	C(9)–C(10)–C(11)	120.1(2)
C(6)–C(11)–C(10)	120.6(2)	C(15)–C(14)–C(19)	119.38(15)
C(15)–C(14)–C(5)	119.38(15)	C(19)–C(14)–C(5)	121.25(15)
C(16)–C(15)–C(14)	121.14(16)	C(15)–C(16)–C(17)	119.82(17)
C(16)–C(17)–C(18)	119.61(16)	C(19)–C(18)–C(17)	121.31(15)
C(18)–C(19)–C(14)	118.74(15)	C(18)–C(19)–P(1)	122.70(13)
C(14)–C(19)–P(1)	118.47(12)	C(26)–P(1)–C(20)	101.48(8)
C(26)–P(1)–C(19)	101.14(7)	C(20)–P(1)–C(19)	101.48(8)
C(25)–C(20)–C(21)	118.77(16)	C(25)–C(20)–P(1)	123.79(13)
C(21)–C(20)–P(1)	117.30(14)	C(22)–C(21)–C(20)	120.35(18)
C(23)–C(22)–C(21)	120.22(17)	C(22)–C(23)–C(24)	120.03(17)
C(23)–C(24)–C(25)	119.96(18)	C(20)–C(25)–C(24)	120.64(17)
C(31)–C(26)–C(27)	118.55(16)	C(31)–C(26)–P(1)	124.01(14)
C(27)–C(26)–P(1)	117.44(13)	C(28)–C(27)–C(26)	120.50(16)
C(29)–C(28)–C(27)	120.18(17)	C(30)–C(29)–C(28)	119.48(18)
C(31)–C(30)–C(29)	120.35(17)	C(30)–C(31)–C(26)	120.92(17)

Table 4. Anisotropic displacement parameters ( $\text{\AA}^2$ ) for pchp3. The anisotropic displacement factor exponent takes the form:  $-2\pi^2[h^2a^{*2}U^{11} + \dots + 2hka^*b^*U^{12}]$


	$U^{11}$	$U^{22}$	$U^{33}$	$U^{23}$	$U^{13}$	$U^{12}$
O(1)	0.0231(6)	0.0599(9)	0.0244(6)	0.0084(6)	0.0038(4)	-0.0049(6)
C(2)	0.0256(7)	0.0353(9)	0.0240(7)	0.0027(7)	0.0030(6)	-0.0010(7)
C(3)	0.0261(8)	0.0474(10)	0.0240(7)	0.0073(7)	0.0036(6)	-0.0035(7)
N(4)	0.0211(7)	0.0526(10)	0.0257(7)	0.0076(7)	0.0028(5)	-0.0056(6)
C(5)	0.0270(8)	0.0371(10)	0.0244(8)	0.0028(7)	0.0049(6)	-0.0029(7)
C(6)	0.0217(8)	0.0420(10)	0.0301(8)	0.0099(7)	0.0035(6)	-0.0030(6)
C(7)	0.0347(10)	0.0431(11)	0.0578(13)	0.0087(10)	0.0101(9)	0.0036(8)
C(8)	0.0364(11)	0.0552(15)	0.110(3)	0.0338(16)	0.0198(13)	0.0126(10)
C(9)	0.0327(11)	0.099(2)	0.0739(18)	0.0587(18)	-0.0001(11)	-0.0009(13)
C(10)	0.0329(10)	0.103(2)	0.0393(10)	0.0307(14)	-0.0048(8)	-0.0164(14)
C(11)	0.0309(9)	0.0647(14)	0.0301(9)	0.0096(9)	0.0017(7)	-0.0094(9)
C(12)	0.0330(10)	0.0924(19)	0.0320(10)	0.0089(11)	0.0132(8)	0.0060(11)
C(13)	0.0327(9)	0.083(2)	0.0391(10)	0.0146(11)	-0.0050(8)	-0.0218(11)
C(14)	0.0267(8)	0.0287(8)	0.0216(7)	-0.0006(6)	0.0013(6)	-0.0040(6)
C(15)	0.0289(8)	0.0324(9)	0.0294(8)	-0.0018(7)	0.0072(6)	-0.0004(7)
C(16)	0.0265(8)	0.0263(8)	0.0373(9)	-0.0014(7)	0.0018(6)	-0.0002(6)
C(17)	0.0303(8)	0.0256(8)	0.0296(8)	0.0037(6)	0.0027(6)	-0.0022(6)
C(18)	0.0272(8)	0.0273(8)	0.0234(7)	-0.0008(6)	0.0021(6)	-0.0019(6)
C(19)	0.0221(7)	0.0247(8)	0.0236(7)	-0.0018(6)	0.0003(6)	-0.0019(6)
P(1)	0.02449(19)	0.02588(19)	0.02435(18)	0.00130(16)	-0.00044(13)	0.00078(16)
C(20)	0.0217(7)	0.0249(8)	0.0293(7)	0.0016(7)	0.0005(5)	0.0029(6)
C(21)	0.0264(8)	0.0310(9)	0.0361(9)	0.0039(7)	-0.0001(7)	-0.0009(7)
C(22)	0.0250(8)	0.0353(9)	0.0458(10)	0.0125(8)	0.0047(7)	-0.0007(7)
C(23)	0.0290(9)	0.0477(11)	0.0345(9)	0.0117(8)	0.0081(7)	0.0048(7)
C(24)	0.0365(9)	0.0414(10)	0.0289(9)	-0.0003(7)	0.0046(7)	0.0023(8)
C(25)	0.0266(8)	0.0317(9)	0.0307(8)	0.0006(7)	0.0038(6)	-0.0012(7)
C(26)	0.0262(7)	0.0267(8)	0.0231(8)	0.0006(6)	0.0053(6)	-0.0004(6)
C(27)	0.0298(8)	0.0303(8)	0.0302(8)	0.0029(7)	0.0022(6)	0.0023(7)
C(28)	0.0440(10)	0.0239(8)	0.0364(10)	0.0026(7)	0.0096(8)	-0.0012(7)
C(29)	0.0371(10)	0.0352(10)	0.0378(10)	-0.0030(8)	0.0080(8)	-0.0101(8)
C(30)	0.0284(9)	0.0404(10)	0.0463(11)	-0.0015(8)	-0.0016(8)	-0.0027(8)
C(31)	0.0304(9)	0.0285(9)	0.0419(10)	0.0044(7)	-0.0025(7)	0.0027(7)

Table 5. Hydrogen coordinates and isotropic displacement parameters ( $\text{\AA}^2$ ) for pcbp3.


	x	y	z	U
H(2)	0.8401	0.5674	0.4701	0.034
H(3)	1.0180	0.8005	0.5012	0.039
H(5)	0.9264	0.7666	0.6603	0.035
H(7)	0.6558	0.8646	0.5170	0.054
H(8)	0.4586	0.9786	0.4170	0.080
H(9)	0.3832	0.9048	0.2748	0.083
H(10)	0.4996	0.7199	0.2315	0.071
H(11)	0.6952	0.6056	0.3298	0.050
H(12A)	1.1542	0.5770	0.4318	0.078
H(12B)	1.2681	0.7024	0.4412	0.078
H(12C)	1.0890	0.6931	0.3728	0.078
H(13A)	1.2953	0.6968	0.6856	0.078
H(13B)	1.2248	0.8185	0.6347	0.078
H(13C)	1.3550	0.7243	0.5891	0.078
H(15)	1.1219	0.4732	0.6721	0.036
H(16)	1.1422	0.3407	0.7924	0.036
H(17)	0.9759	0.3823	0.9135	0.034
H(18)	0.7909	0.5573	0.9133	0.031
H(21)	0.4081	0.6076	0.8419	0.038
H(22)	0.2509	0.5841	0.9679	0.042
H(23)	0.3193	0.7109	1.0896	0.044
H(24)	0.5533	0.8566	1.0880	0.043
H(25)	0.7208	0.8749	0.9645	0.036
H(27)	0.7034	1.0119	0.7516	0.036
H(28)	0.9001	1.1765	0.7887	0.041
H(29)	1.1813	1.1432	0.8719	0.044
H(30)	1.2605	0.9456	0.9205	0.046
H(31)	1.0629	0.7823	0.8859	0.041

## APPENDIX 5

(1R,2S)-2-[1,2-dimethyl propyl) methyl amino]-1, -phenyl propyl-3, -5-dinitro-  
benzene-1-carboxylate



Sa357



## F2 - Acquisition Parameters

Date\_ 20011214  
Time 14.16  
INSTRUM opx400  
PROBHD 5 mm Multinu  
PULPROG cosy45  
TD 1024  
SOLVENT D2O  
NS 28  
DS 1E  
SWH 4194.631 Hz  
FIDRES 4.096319 Hz  
AQ 0.1221108 sec  
RG 287.4  
DW 119.200 usec  
DE 7.50 usec  
TE 300.0 K  
D1 1.00000000 sec  
P1 15.00 usec  
SF01 400.1319304 MHz  
NUC1 1H  
PL1 -6.00 dB  
D0 0.00000300 sec  
IN0 0.00023840 sec

## F1 - Acquisition parameters

MD0 1  
TO 243  
SF01 400.1319 MHz  
FIDRES 17.261856 Hz  
SW 10.483 ppm

## F2 - Processing parameters


S1 512  
SF 400.130097 MHz  
WDW SINE  
SSB 0  
LB 0.00 Hz  
GB 0  
PC 1.00

## F1 - Processing parameters

S1 512  
MC2 OF  
SF 400.130097 MHz  
WDW SINE  
SSB 0  
LB 0.00 Hz  
GB 0

## 2D NMR plot parameters

CX2 15.00 cm  
CX1 15.00 cm  
F2PLD 4.002 ppm  
F2LD 1601.13 Hz  
F2PH1 0.193 ppm  
F2H1 77.30 Hz  
F1PLD 4.002 ppm  
F1LD 1601.13 Hz  
F1PH1 0.193 ppm  
F1H1 77.30 Hz  
F2PPMCH 0.25389 ppm/cm  
F2HZCM 101.58871 Hz/cm  
F1PPMCH 0.25389 ppm/cm  
F1HZCM 101.58871 Hz/cm



ppm

## F2 - Acquisition Parameters

Date\_ 20011214  
Time 14.16  
INSTRUM opx400  
PROBHD 5 mm Multinu  
PULPROG cosy45  
TD 1024  
SOLVENT D2O  
NS 26  
DS 16  
SWH 4194.631 Hz  
FIDRES 4.095319 Hz  
AQ 0.1221108 sec  
RG 287.4  
DW 119.200 usec  
DE 7.50 usec  
TE 300.0 K  
D1 1.0000000 sec  
P1 15.00 usec  
SFO1 400.1319304 MHz  
NUC1 1H  
PL1 -6.00 dB  
D0 0.00000300 sec  
IN0 0.00023840 sec

## F1 - Acquisition parameters

ND0 1  
TD 243  
SFO1 400.1319 MHz  
FIDRES 17.261856 Hz  
SW 10.483 ppm

## F2 - Processing parameters


SI 512  
SF 400.1300097 MHz  
WDW SINE  
SSB 0  
LB 0.00 Hz  
GB 0  
PC 1.00

## F1 - Processing parameters

SI 512  
MC2 OF  
SF 400.1300097 MHz  
WDW SINE  
SSB 0  
LB 0.00 Hz  
GB 0

## 2D NMR plot parameters

CX2 15.00 cm  
CX1 15.00 cm  
F2PLO 4.000 ppm  
F2L0 1600.52 Hz  
F2PH1 0.200 ppm  
F2H1 80.03 Hz  
F1PLO 4.000 ppm  
F1L0 1600.52 Hz  
F1PH1 0.200 ppm  
F1H1 80.03 Hz  
F2PPHCH 0.25333 ppm/cm  
F2HZCH 101.36626 Hz/cm  
F1PPHCH 0.25333 ppm/cm  
F1HZCH 101.36626 Hz/cm



## F2 - Acquisition Parameters

Date\_ 20011214  
 Time 14.16  
 INSTRUM gp400  
 PROBHD 5 mm Multinu  
 PULPROG cosy45  
 TO 1024  
 SOLVENT D2O  
 NS 28  
 DS 16  
 SWH 4194.631 Hz  
 FIDRES 4.096319 Hz  
 AQ 0.1221108 sec  
 RG 287.4  
 DW 119.200 usec  
 DE 7.50 usec  
 TE 300.0 K  
 D1 1.00000000 sec  
 P1 15.00 usec  
 SFO1 400.1319304 MHz  
 NUC1 1H  
 PL1 -6.00 dB  
 D0 0.00000300 sec  
 IN0 0.00023840 sec

## F1 - Acquisition parameters

MD0 1  
 TO 243  
 SFO1 400.1319 MHz  
 FIDRES 17.261856 Hz  
 SW 10.483 ppm

## F2 - Processing parameters


SI 512  
 SF 400.1300097 MHz  
 WDW SINE  
 SSB 0  
 LB 0.00 Hz  
 GB 0  
 PC 1.00

## F1 - Processing parameters

SI 512  
 MC2 OF  
 SF 400.1300097 MHz  
 WDW SINE  
 SSB 0  
 LB 0.00 Hz  
 GB 0

## 2D NMR plot parameters

CX2 15.00 cm  
 CX1 15.00 cm  
 F2PLO 10.042 ppm  
 F2LO 4017.96 Hz  
 F2PHI -0.442 ppm  
 F2HI -176.67 Hz  
 F1PLO 10.042 ppm  
 F1LO 4017.96 Hz  
 F1PHI -0.442 ppm  
 F1HI -176.67 Hz  
 F2PPMCH 0.69888 ppm/cm  
 F2HZCM 279.64206 Hz/cm  
 F1PPMCH 0.69888 ppm/cm  
 F1HZCM 279.64206 Hz/cm



## F2 - Acquisition Parameters

Date\_ 20011214  
 Time 14.16  
 INSTRUM opx400  
 PROBHD 5 mm Multinu  
 PULPROG cosy42  
 TD 1024  
 SOLVENT D2O  
 NS 26  
 DS 16  
 SWH 4194.631 Hz  
 FIDRES 4.096319 Hz  
 AQ 0.1221108 sec  
 RG 267.4  
 OW 119.200 usec  
 DE 7.50 usec  
 TE 300.0 K  
 D1 1.00000000 sec  
 P1 15.00 usec  
 SFO1 400.1319304 MHz  
 NUC1 1H  
 PL1 -6.00 dB  
 DO 0.0000300 sec  
 INO 0.00023840 sec

## F1 - Acquisition parameters

MD0 1  
 TD 243  
 SFO1 400.1319 MHz  
 FIDRES 17.261856 Hz  
 SW 10.483 ppm

## F2 - Processing parameters

SI 512  
 SF 400.130097 MHz  
 WDW SINE  
 SSB 0  
 LB 0.00 Hz  
 GB 0  
 PC 1.00

## F1 - Processing parameters

SI 512  
 MC2 OF  
 SF 400.130097 MHz  
 WDW SINE  
 SSB 0  
 LB 0.00 Hz  
 GB 0

## 2D NMR plot parameters

CX2 15.00 cm  
 CX1 15.00 cm  
 F2PLO 6.500 ppm  
 F2LO 2600.84 Hz  
 F2PHI 0.200 ppm  
 F2HI 80.03 Hz  
 F1PLO 6.500 ppm  
 F1LO 2600.84 Hz  
 F1PHI 0.200 ppm  
 F1HI 80.03 Hz  
 F2PPMCH 0.42000 ppm/cm  
 F2HZCM 168.05461 Hz/cm  
 F1PPMCH 0.42000 ppm/cm  
 F1HZCM 168.05461 Hz/cm

