

This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Hydrological droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1073/pnas.1222473110>

PUBLISHER

© National Academy of Sciences

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Prudhomme, Christel, Ignazio Giuntoli, Emma L. Robinson, Douglas B. Clark, Nigel W. Arnell, Rutger Dankers, Balazs M. Fekete, et al.. 2019. "Hydrological Droughts in the 21st Century, Hotspots and Uncertainties from a Global Multimodel Ensemble Experiment". figshare. <https://hdl.handle.net/2134/22127>.

Hydrological droughts in the 21st century: hotspots and uncertainties from a global multi-model ensemble experiment

Authors: Christel Prudhomme¹, Ignazio Giuntoli^{1,2}, Emma L. Robinson¹, Douglas B. Clark¹, Nigel W. Arnell³, Rutger Dankers⁴, Balázs Fekete⁵, Wietse Franssen⁶, Dieter Gerten⁷, Simon N. Gosling⁸, Stefan Hagemann⁹, David M. Hannah², Hyungjun Kim¹⁰, Yoshimitsu Masaki¹¹, Yusuke Satoh¹², Tobias Stacke⁹, Yoshihide Wada¹³, Dominik Wisser^{13,14}

¹ Centre for Ecology and Hydrology, Maclean Building, Wallingford, Oxfordshire, OX10 8BB, United Kingdom ² School of Geography, Earth and Environment Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom ³ Walker Institute for Climate System Research, University of Reading, Reading, RG6 6AR, United Kingdom ⁴ Met Office Hadley Centre, Exeter, United Kingdom ⁵ Civil Engineering Department, The City College of New York, New York, USA ⁶ Earth System Science, Wageningen University and Research Centre, P.O. Box 47, 6700 AA Wageningen, Netherlands ⁷ Potsdam Institute for Climate Impact Research, Germany ⁸ School of Geography, University of Nottingham, Nottingham, NG7 2RD, United Kingdom ⁹ Max Planck Institute for Meteorology, Bundesstraße 53, 20146 Hamburg, Germany ¹⁰ Institute of Industrial Science, The University of Tokyo, Tokyo, Japan ¹¹ National Institute for Environmental Studies, Japan ¹² Department of Civil Engineering, The University of Tokyo, Japan ¹³ Department of Physical Geography, Utrecht University, Heidelberglaan 2, 3584 CS Utrecht, Netherlands ¹⁴ Center for Development Research, University of Bonn, Germany

Submitted to Proceedings of the National Academy of Sciences of the United States of America

Increasing concentrations of greenhouse gases in the atmosphere are expected to modify the global water cycle with significant consequences for terrestrial hydrology. We assess the impact of climate change on hydrological droughts in a multi-model experiment including seven Global Impact Models (GIMs) driven by bias-corrected climate from five Global Climate Models (GCMs) under four Representative Concentration Pathways (RCPs). Drought severity is defined as the fraction of land under drought conditions. Results show a likely increase in the global severity of hydrological drought at the end of the 21st century, with systematically greater increases for RCPs describing stronger radiative forcings. Under RCP8.5, droughts exceeding 40% of analysed land area are projected by nearly half of the simulations. This increase in drought severity has a strong Signal-to-Noise ratio at the global scale, and Southern Europe, Middle East, South East United States, Chile and South West Australia are identified as possible hotspots for future water security issues. The uncertainty due to GIMs is greater than that from GCMs, particularly if including a GIM that accounts for the dynamic response of plants to CO₂ and climate, as this model simulates little or no increase in drought frequency. Our study demonstrates that different representations of terrestrial water cycle processes in GIMs are responsible for a much larger uncertainty in the response of hydrological drought to climate change than previously thought. When assessing the impact of climate change on hydrology it is hence critical to consider a diverse range of GIMs to better capture the uncertainty.

Drought | Climate impact | global hydrology | evaporation | CO₂

Introduction

The global water cycle is expected to change over the 21st century due to the combined effects of climate change and increasing human intervention. In a warmer world the water holding capacity of the atmosphere will increase, resulting in a change in the frequency of precipitation extremes, increased evaporation and dry periods (1) and intensification of droughts (2). This is represented by most Global Climate Models (GCMs) by increased summer dryness and winter wetness over large areas of continental mid-to high-latitudes in the Northern Hemisphere (3), associated with a reduction in water availability at continental (4, 5) and global scales (6, 7). Because such changes have potentially very serious implications in some regions of the world, identifying areas where there is agreement in the direction and magnitude of changes in drought characteristics (hotspots) in response to climate change is essential information for water resource management aimed at ensuring water security in a changing climate.

Most GCMs, however, are not able to reproduce the fine-scale processes governing terrestrial hydrology (and hence runoff) and suffer from systematic biases (8). As land-atmospheric feedbacks are not yet fully understood and reproduced by global models (9), and because full coupling of GCMs and Global Impact Models (GIMs) is not straightforward, GIMs forced by data from GCMs have been used as tools to quantify the impact of changed climate on the water cycle and droughts (10), despite by definition ignoring important feedbacks and their possible modification with climate change (11). GIMs vary in the types of processes represented and the parameterisations employed. Some GIMs, particularly those designed to quantify water resources, only calculate the water balance (e.g. 12), while others consider coupled water and energy balances, sometimes also representing the dynamic response of plants to changes in atmospheric CO₂ and climate (e.g. 13). Until recently the uncertainty in the simulation of the terrestrial water cycle related to the choice of a particular GIM had not been investigated. However, the Water Model Intercomparison

Significance

Increasing concentrations of greenhouse gases in the atmosphere are widely expected to influence global climate over the coming century. The impact on drought is uncertain because of the complexity of the processes, but can be estimated using outputs from an ensemble of global models (hydrological and climate models). Using an ensemble of 35 simulations we show a likely increase in the global severity of drought by the end of 21st century, with regional hotspots including South America and Central and Western Europe in which the frequency of drought increases by more than 20%. The main source of uncertainty in the results comes from the hydrological models, with climate models contributing to a substantial but smaller amount of uncertainty.

Reserved for Publication Footnotes

137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204




Fig. 1. : Percentage change in the occurrence of days under drought conditions for the period 2070-2099 relative to 1976-2005, based on a multi-model ensemble MME experiment under RCP8.5 from five Global Climate Models and seven Global Impact Models: MME Mean change (left) and associated Signal-to-Noise ratio (S2N, MME mean change divided by its Inter-Quartile range, right). See Methods for definition of drought, S2N and masking procedure.




Fig. 2. Mean change in drought severity (Change in GDI, y-axis) as measured by the daily Global Deficit Index GDI for the period 2070-2099 relative to 1976-2005 based on a multi-model ensemble MME experiment calculated over the whole year (left), December to February (DJF middle) and June to August (JJA right). Changes are given for each MME member and are organised by radiative forcing (from left to right RCP2.6 R2, RCP4.5 R4, RCP6.0 R6, RCP8.5 R8). In each RCP panel, results are organised according to driving GCMs from left to right: HadGEM2-ES; IPSLCM5-ARL; MIROC-ESM-CHEM; GFDL-ESM2M and NorESM1-M. CO₂ effect in GIMs is described as colour: black/open symbols No CO₂; cyan/filled symbols: CO₂. GIMs are indicated by symbols: up triangle HO8; circle JULES; x Mac-PDM.09; +MPI-HM; pentagon PRCGLOB-WB; down triangle VIC; square WBM.

Project (Water MIP; 14) highlighted that simulated hydrological averages can vary substantially between GIMs, even when driven with the same bias-corrected climatic forcing (14, 15), and uncertainty in future projection due to GIMs can be as large as that from GCMs in some regions (16, 17). While in the climate-to-impact modelling chain much effort has been directed to better understand the uncertainty due to GCMs, studies of the impact of climate change on water availability and drought have often been based on one or a few GIMs, potentially underestimating the overall uncertainty.

This study focuses on identifying regions where the impact of climate change on hydrological drought (henceforth simply 'drought') shows a strong signal of change between the end of the 20th and 21st centuries. We define drought as occurring when total runoff is less than a given threshold. Drought represents the time-integrated effect of several interlinked processes and stores, including precipitation, evaporation and soil moisture storage (10); because some of these processes are represented by GCMs and some by GIMs, it is vital to quantify the relative uncertainty introduced by both GCMs and GIMs when assessing climate change impacts.

We use outputs from the ISI-MIP multi-model ensemble (MME) experiment (18) of 35 members (for RCPs 2.6 and 8.5;




Fig. 3. : Cumulative Density Functions CDFs of daily Global Deficit Index GDI calculated over 30-year periods (1976-2005 for historical forcing and 2070-2099 for RCP forcings) for each Multi-Model Ensemble member.

only 27 members available for RCP4.5 and 6.0) in which GIMs of different types were driven by bias-corrected (8) climate from state-of-the-art CMIP5 GCMs (19). These GIMs describe the terrestrial water cycle at global scale and include current understanding of hydrological systems (20). Note that statistical bias-correction can influence the signal of runoff changes but this generally remains smaller than uncertainty from GCMs and GIMs (21). The simulations we use did not consider water management or changes of land use, so they represent the effects of climate change alone. We quantify changes in the space-time variability of drought that are projected to occur under four Representative Concentration Pathways (RCPs) that span a wide range of ra-

205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272




Fig. 4. : Cumulative Density Functions CDFs of daily Global Deficit Index GDI calculated over 30-year periods (1976-2005 for historical forcing and 2070-2099 for RCP forcings) for simulations accounting for the dynamic response of plants to CO₂ and climate (CO₂) and not (No CO₂) for three Global Impact Models driven by HadGEM2-ES. Note that all CDFs in this figure sample a smaller set of locations than those in Fig. 3.




Fig. 5. : Mean percentage changes in Regional Deficit Index RDI between 30-year simulations of reference (1976-2005) and future (2070-2099) under RCP8.5 for 17 world regions. Values are averaged over all the MME members (All), by GCMs and by GIMs. JULES includes CO₂ and vegetation effects. See SI and (27) for GEO region descriptions and acronyms.

diative forcing (22) (Methods). We also evaluate the uncertainty associated with both GCMs and GIMs, so as to identify hotspots of change where we have more confidence in the projections of future drought severity.

Results

We analyse future droughts by comparing the temporal and spatial patterns of simulated runoff in the years 2070-2099 (RCP forcings) with those from a reference period 1976-2005 (historical forcings). For each land cell and simulation we define a runoff Deficit Index (DI) which is equal to 1 if the daily runoff (not river discharge) is less than a daily drought threshold (10th percentile) calculated from the reference period (Methods), and is zero otherwise; a grid cell with DI=1 is hence under drought conditions. As a measure of the severity of drought, Global and Regional Deficit Index (GDI and RDI) time series are calculated as the fraction of land area on the globe or a given region that is under drought conditions. Arid grid cells in which runoff is equal to zero more than 90% of time in the reference or future periods are excluded (white areas on Fig. 1). Also excluded were GIMs for which on average fewer than 75% of the remaining land cells could be used to calculate a DI because of a high proportion of zero runoff values; hence a total of seven GIMs driven by five GCMs are included in the MME for RCP2.6 and RCP8.5 (Methods; Figs. S2 and S3 in Supplementary Information SI). The MME mean changes in GDI and RDI are examined at the annual and seasonal scale, with Signal-to-Noise ratios (S2N; MME mean divided by its Inter-Quartile range, Methods) used to quantify the relative sizes of variability due to GIMs and GCMs, and to identify hotspots where the signal is strongest.

Global changes

Under RCP8.5, the MME mean change in the frequency of drought (i.e. DI=1) shows a widespread increase of drought conditions across the globe and in particular in most parts of South and North America, large parts of tropical and southern Africa, the Mediterranean region, South East China and Australia; little change or reduced occurrence of drought conditions are found in northern Canada, North East Russia, the Horn of Africa and parts of Indonesia (Fig. 1). There is strong seasonality across many mid- to high-latitude regions in the Northern Hemisphere, with small changes or reductions in DJF and larger increases in JJA (Fig. S4). For 25 members (i.e. 70% of the ensemble) the frequency of drought increases in 60.3% of unmasked land cells, falling to 44.9% in DJF when there is the largest degree of disagreement between ensemble members as to the direction of changes. Over the whole year, S2N is largest in the Mediterranean and the Middle East, Chile, South East US, and Western Australia (Fig. 1).

In Fig. 2 we calculate the mean change in GDI for the four RCPs. The results show a likely increase in drought severity with a MME mean increase of 3.9% under RCP2.6, 6.3% for RCP4.5, 7.4% for RCP6.0 and reaching 13% under RCP8.5 (see (23) for method and SI for detailed results); changes are largest in JJA (17.6%) and smallest in DJF (10.6%) under RCP8.5. The systematic increase in drought severity with radiative forcing (Fig. 2) is associated with considerable variation in the magnitude of the changes ranging from -1.7% to +11.2% under RCP2.6 and -4.8% to 25.4% under RCP8.5. S2N associated with GCMs and GIMs shows a stronger signal (less uncertainty) for GCMs (mean S2N=2.44) than for GIMs (1.82) primarily due to smaller IQ for GCMs (mean IQ=0.049) than for GIMs (0.070) (SI for details). This indicates that, at the global scale, the variability due to different GIMs is larger than that due to different GCMs.

There is a statistically significant (see Methods) increase in the frequency of severe events (large GDI) for all RCP/GCM/GIM combinations except for JULES which shows a consistently smaller change signal in all simulations but one for RCP 8.5 (Cumulative Density Function CDF, Fig. 3). Under historical forcing, drought affects less than 21% of the global land area at any one time (GDI<0.21; black lines) but this is exceeded for 23 out of 35 simulations under RCP2.6 (dark blue) and for 30 under RCP8.5 (red). Largest increases are seen for RCP8.5, with maximum drought severity exceeding 40% of land in 16 simulations. There is greater temporal variability in the GDI in many simulations of the RCPs (flatter CDFs in Fig. 3), increasing with radiative forcing, and associated with more pronounced variability between GIMs.

Effects of model structure

All the models shown in Fig. 3 calculate the water balance of the land, but only H08 and JULES consider the energy balance (Table S1 in SI), and only JULES represents the effects of CO₂ on stomatal opening and includes a dynamic vegetation model that allows vegetation to grow in response to its environment. To examine whether the different behaviour of JULES was at-

tributable to one of these differences in model structure, we used results from two further GIMs (LPJmL and MATSIRO) that were excluded from the MME because of a high proportion of zero runoff values (Methods and Figs. S2 and S3 in SI) but which share some similarities with JULES in terms of model structure: MATSIRO and JULES are energy (and water) balance models, while LPJmL and JULES both represent varying CO₂ and dynamic vegetation effects. For this analysis the GDI was calculated using the smaller sample of land cells dictated by LPJmL and MATSIRO (Fig. S3 in SI) after the masking procedure was applied to those GIMs. CDFs from the energy balance GIMs show a strong response of H08 and MATSIRO to climate change (Fig. 4), broadly similar to those from the water balance models of Fig. 3, which is evidence that it is not the inclusion of an energy balance that makes JULES different. (The CDFs in Fig. 4 cannot be compared directly with those in Fig. 3 because of the different locations sampled. However, as the sampling does not substantially alter the distributions for JULES between these figures, qualitative comparisons of the figures can be made.)

In contrast, the CDFs for the models that include CO₂ and vegetation effects (JULES and LPJmL; solid lines in Fig. 4 right) show a weaker response to climate change. The effect on plants of increased CO₂ concentration is often considered to consist of physiological and structural components. The former results in the stomata opening less widely in a CO₂-enriched atmosphere, leading to less water loss through transpiration (24). However, increased growth can alter the structure of the vegetation, potentially resulting in increased leaf area (and increased transpiration) (25). Sensitivity experiments in which CO₂ was allowed to vary only until the year 2000, after which it remained constant, showed increased response to climate from both JULES and LPJmL (dashed lines in Fig. 4 right), albeit the increase is much more pronounced in JULES. Both models gave less transpiration under higher CO₂ than when CO₂ was constant despite having increased biomass (Fig. S5 in SI) indicating a strong physiological effect of CO₂. Further runs of JULES in which all structural aspects of the vegetation (fractional coverage, leaf area and height) were kept constant in time showed that the largest increase in drought occurred when both CO₂ and vegetation were constant (Fig. S6 in SI) – that is for the configuration most similar to that of the other GIMs shown in Figs. 2 and 3. This suggests that accounting for the dynamic response of plants to CO₂ and climate is largely responsible for the outlying (small) response of JULES in Fig. 3. This is consistent with a study showing that a substantially lower irrigation water demand under climate change is simulated by GIMs including CO₂ effect than those without (26).

Regional changes

We calculate the RDI for the 17 GEO sub-regions (27) for which the DI could be calculated at more than 50% of the points in the region; note however the wide variation in size between regions (27). Average changes under RCP8.5 (top line in Fig. 5) vary between no change (Eastern Africa) to 28% (Central Europe), with 5 regions showing increases of at least 20% (South and Meso-America, Caribbean, Central and Western Europe).

S2N is larger than 1 in 6 regions, and is larger than 1.5 in Western and Central Europe. Uncertainty is largest in Eastern Europe, South East Asia, South America and Eastern Africa where S2N is less or equal to 0.7. When calculated separately for GCMs and GIMs ensembles, the average S2N resulting from different GCMs is larger than that from GIMs in 11 regions, and in 3 of those it is more than 1.5 times as large (Meso America, Central Europe and Caribbean; see Table S3 in SI for details). This is seen in Fig. 5 where the variation in mean change between the GCMs (lines near top of figure) is smaller than that between the GIMs (lines at bottom of figure). In North and South America, Eastern Europe, East and South Asia, Central and Eastern Africa, S2N from GCMs and GIMs are comparable. When JULES is excluded

from the ensemble (i.e. only 6 GIMs included) the mean change remains relatively similar to that from the 7GIMs ensemble but the S2N ratio increases, with magnitude depending on the region (Tables S2 and S3 in SI for details). Note however that even with JULES excluded from the MME, the uncertainty from GIMs generally remains greater than that from GCMs, suggesting that JULES is not the dominant source of uncertainty within GIMs.

Discussion

Previous global modelling experiments have suggested that under climate change soil moisture and runoff would decrease (see e.g. 4-7, 10, 28, 29-32), albeit with large regional uncertainty in the magnitude of changes. However most studies only include one or a few Global Impact Models (GIMs), while recent work (17, 26) has shown that the uncertainty associated with the response of terrestrial hydrology to climate change simulated by different GIMs could be as large as the uncertainty in the response of the climate to greenhouse gas forcings simulated by Global Climate Models (GCMs). The uncertainty associated with GIMs has been attributed to differences in the number and type of processes represented in the GIMs (e.g. water balance, energy balance), and to differences in the details of their implementations. We used a Multi-Model Ensemble (MME) experiment including a relatively large number of GIMs of diverse types (which are able to reproduce the main characteristics of water deficits in terms of regional extent and duration (15)), forced by the same bias-corrected climate from simulations of five state-of-the-art GCMs, to assess changes in the frequency and severity of droughts, at the global and regional scales, under four different RCPs for the end of the 21st century.

At the regional scale, our results show that drought frequency (proportion of time under drought conditions) and severity (proportion of land under drought conditions) is very likely (i.e. more than 90% of ensemble members) to increase in the Caribbean, South America, Western and Central Europe, Central Africa, Australia and New Zealand and Western Indian Ocean under RCP8.5; this reinforces earlier findings based on CMIP3/ SRES projections (2). In Eastern Africa, the variation between GCMs is large and the Signal-to-Noise ratio (S2N) is close to zero. Both GCMs and GIMs contribute to the overall uncertainty in the response; improving the representation of regional changes in droughts is dependent on improved process representation in the models, for example through analyses of GIMs biases when forced by observed climate. It is however beyond the scope of this paper to diagnose the reasons for the differences between particular GCMs and GIMs.

The MME shows a likely increase in the spatial extent (severity) of dry episodes under all four RCPs, with increasingly large changes under greater radiative forcing and S2N greater than 1 at the global scale and in some regions. Under RCP8.5, all five GCMs show a substantially warmer climate which will tend to drive increased evaporation. There is more variation in projected precipitation changes, both between regions and between GCMs (Fig. S1 of SI). Where evaporation increases and precipitation decreases, soil moisture deficit can build up, resulting in increased drought. In our results, areas with the largest signal of drought increase are generally located where precipitation is projected to decrease. However, even in those areas where precipitation increases, drought can still increase if this extra water is lost through greater evaporation. This is the case in tropical areas where GCMs indicate increased precipitation (e.g. parts of Central Africa; Fig. S1 in SI), but increase in evaporation leads to more drought being simulated by GIMs (Fig. 1). Using S2N calculated over different sub-sets of the MME describing range of GCMs and GIMs, we showed that the total uncertainty associated with projected changes in drought is larger from GIMs compared to GCMs. GIMs uncertainty is particularly affected by an outlying

GIM, JULES, that shows systemically lower response to climate change, but remains larger than the uncertainty in GCMs even when excluding it from the ensemble, e.g. GDI S2N is 2.48 from GIMs and 3.01 from GCMs when excluding JULES (Table S3 in SI; numbers in bracket for details).

By investigating JULES simulations further, we show that its outlying signal is largely the result of the inclusion of a description of the plant response to enhanced CO₂, a process that is not represented in most GIMs used to simulate global water resources. The effects of CO₂ and dynamic vegetation on plant evapotranspiration and mean runoff have been studied before (e.g. 24, 25, 33, 34) but the effect on drought and a direct comparison with hydrological models has not been presented before. When atmospheric CO₂ increases, the stomata can partially close (35) conserving the water and resulting in smaller changes of evapotranspiration in a warmer climate (26, 36). This leaves a wetter soil and thereby a less likely drought occurrence, as found in our results. At the leaf-scale the physiological effect of increased CO₂ is well characterised by laboratory and field studies (37) but models differ substantially in the predicted response of transpiration at the ecosystem level (38) and the net effect of physiological and structural changes is also highly uncertain (39). Our results suggest that the inclusion of CO₂ and vegetation dynamics can fundamentally change the drought response to climate change but the magnitude of these changes remain uncertain. This underlines the importance of including a diverse range of GIMs describing various processes when designing multi-model experiments, and that more research should be conducted to better understand the response of vegetation water use to CO₂ increase.

Our MME only considered the impact of climate change, with no representation of water management or changes in land use. Climate (including CO₂ effects on vegetation) is not the only forcing relevant to assessments of future droughts and water scarcity as water demand can generate water stress (40) and the projected future population increase will likely result in further increases in water stress (41). For a thorough investigation of water availability, the combined effect of climate, land use and water management should be taken into account, using a range of GCMs and GIMs to capture the uncertainty.

Methods

In this paper we have analysed simulations from the Global Impact Models of the ISI-MIP ensemble experiments for which daily runoff data were available.

The experiments considered five different worlds: one representative of historical radiative forcing and four possible future worlds. These future scenarios included: a very high baseline (rising radiative forcing reaching 8.5Wm⁻² by 2100, RCP8.5), a very low forcing level (radiative forcing peaking at 3Wm⁻² before declining to reach 2.6Wm⁻² by 2100 RCP2.6), and two medium stabilization scenarios (stabilization without overshoot pathway to 4.5/6.0 Wm⁻² at 2100 RCP4.5/RCP6.0) (22, 42). Each radiative forcing scenario was implemented by five Global Climate Models (GCMs): HadGEM2-ES; IPSL-CM5A-LR, MIROC-ESM-CHEM, GFDL-ESM2M and NorESM1-M (18). Transient GCM outputs were re-gridded to a common 0.5° latitude x 0.5° longitude grid and a 2-step bias correction procedure implemented for each month independently (8) based on the WATCH Forcing data (43).

The bias-corrected GCM outputs (8) were used as inputs for nine Global Impact Models: H08, JULES, LPJmL, Mac-PDM.09, MATSIRO, MPI-hm, PRC-GlobWB, VIC and WBM (see SI for references). For RCP4.5 and 6.0, VIC and Mac-PDM.09 were only driven by HadGEM2-ES. The GIMs were run on 0.5° grids (except JULES, which was run with grid cell size 1.875° longitude x 1.25° latitude, then regridded to 0.5° for analysis). GIMs were spun up to a quasi steady state by repeated use of detrended meteorology for 1951-1980, followed by a simulation of the period 1951-2005. Simulations for each RCP covered 2006-2099 (2005-2099 for HadGEM2-ES-forced runs). All GIMs considered contemporary patterns of land use, except JULES which modelled natural vegetation only, with no land use. No anthropogenic storage (e.g. dams and reservoir) or water management were represented.

We have not investigated the extent to which the drought results from JULES depend on the grid scale, as no simulations at other scales are available. However, our best assessment from other work with JULES (not specifically on drought) is that results are generally not very sensitive to the size of the grid cells, at least for modest changes in resolution (say 0.5 to 3 degrees) and for regionally or globally averaged statistics.

Daily total runoff is the sum of surface and subsurface runoff. It is an integrated response to all basin input, storage and transfer processes, and the useable output of river basins for various water sectors. The daily total runoff outputs from the RCP/GCM/GIM combinations were extracted and analysed for two time slices: 1976-2005 (historical forcing hist or reference period) and 2070-2099 (future forcing or RCP).

The ISI-MIP dataset also includes experiments of JULES and LPJmL in which CO₂ was allowed to vary only until the year 2000, after which it was kept constant ("noCO₂" runs), while the meteorological forcing included the climate change signal as before. "No CO₂" runs of JULES were only used in the sensitivity analysis of Fig. 4; all other analysis used JULES runs with varying CO₂.

Following (6), drought episodes were defined relative to a time-varying threshold corresponding to the 10th percentile of total runoff (Q90) simulated under hist (notation follows convention from (44) with Q90 being the runoff value exceeded or equalled 90% of the time). We calculated a daily Q90 value using a 30-day moving window to capture intra-monthly temporal patterns, as in (15). This was used as a threshold to calculate daily runoff deficit indices (DIs), such that DI=1 when runoff < Q90 and DI=0 otherwise. All years were treated as having 365 days, with data for 29 Feb removed from models that included that day. The difference in the frequency and fraction of land cells for which runoff is below this threshold between the future and historical runs quantifies any signal of increase in severity and frequency of drought. To capture geographical and seasonal characteristics of the runoff, thresholds were calculated for each land-cell independently. To reduce uncertainty due to climate-impact modelling biases in the results, thresholds were also calculated for each climate-impact model combination as in (45).

For each land cell of each simulation, we calculated a measure of drought frequency to be the fraction of days under drought for each hist and RCP scenario. We quantified a signal as the difference between the fraction of days under drought between each RCP scenario and the corresponding historical scenario. We calculated the Signal-to-Noise ratio (S2N) as the ensemble mean change divided by the Inter-Quartile IQ range of changes for the full MME and certain sub-sets of RCP/GCM/GIM combinations (i.e. S2N associated with GCMs is the S2N associated with each GIM driven by all GCMs, then averaged across all GCMs. For GIMs uncertainty it is the S2N associated with each GCM for all GIMs, then averaged across all GIMs). Sensitivity of S2N to the definition of spread was tested (SI). It showed that values based on IQ range are similar but slightly more conservative (i.e. smaller S2N) than those based on Standard Deviation and hence S2N-IQ was chosen.

Similarly to (46), we removed arid regions from the analysis. We defined arid cells to be those which had more than 90% of the runoff time series equal to zero in any single climate-impact model combination; the land-mask of these cells is displayed in SI. Of the remaining cells, the seasonal variation in the runoff resulted in some very dry periods. We therefore also implemented a daily veto for each cell of each GCM/GIM combination, which discarded days for which the value of Q90 was zero. There were two GIMs (LPJmL and MATSIRO) for which the masking and vetoing removed too many points to be able to calculate global averages (see SI for details), so we did not consider them in the global analysis. After applying the masking procedure to the remaining seven GIMs, 82% of the total land cells (55051 out of 67420 grid cells) was included in the analysis. For JULES, this corresponds to 83% of the total land cells (6270 out of 7558 grid cells). Note that we investigated the effects of model structure (inclusion of dynamic effect of CO₂ on plants) using the two discarded GIMs (LPJmL and MATSIRO) along with JULES, using the full 9GIMs mask, which retained only 64% of the total land cells (68% of the JULES land cells; see SI for details). Therefore, these sensitivity tests are not global results.

The global impact of changing drought was studied by calculating a daily Global Deficit Index (GDI) for each GIM/GCM/RCP combination over the un-masked land cells. This is the weighted average of the number of land cells under drought conditions, with weights proportional to the area of each grid cell. It represents the global proportion of land (or spatial extent) under drought and gives a measure of the global severity of a dry episode; it varies between 0 (no land cells under drought conditions that day) to 1 (all land cells under drought conditions that day). The method differs from that of (47) who calculated spatial extent of droughts over contiguous cells but is similar to that of (6) to avoid potential discontinuity introduced by minor events. Seasonal GDI were derived by extracting GDI time series for two specific 3-month periods: December to February (DJF) and June to August (JJA).

We also calculated a daily Regional Deficit Index (RDI) for 17 of the Geo regions defined in (27). These regions were those for which we could calculate a DI value for at least 50% of the land cells.

Differences between hist and RCP GDIs were assessed using the 1-sided Kolmogorov-Smirnov (48) test which measures the distance between the empirical cumulative distribution functions of two samples of *n*1 observations (here *n*1=365x30=10950). Results are presented at the 95% level.

Acknowledgements.

This work has been conducted under the framework of ISI-MIP. The ISI-MIP Fast Track project was funded by the German Federal Ministry of Education and Research (BMBF) with project funding reference number

01L51201A. Responsibility for the content of this publication lies with the authors. We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the HadGEM2, IPSL and MIROC climate modelling groups for producing and making available their model output. For CMIP the U.S. Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals. The work has been part funded by the CEH-NERC water programme. IG was

1. Trenberth KE (1999) Conceptual framework for changes of extremes of the hydrological cycle with climate change. *Clim. Change* 42:327-339.
2. Field CB, *et al.* eds (2012) *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A special report of Working Groups I and II of the Intergovernmental Panel on Climate Change* (Cambridge University Press, Cambridge, UK, and New York, NY, USA), p 582.
3. Meehl GA, *et al.* (2007) Global Climate Projections. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, & Miller HL (Cambridge University Press, Cambridge, UK), pp 747-845.
4. Cayan DR, *et al.* (2010) Future dryness in the southwest US and the hydrology of the early 21st century drought. *Proceedings of the National Academy of Sciences* 107(50):21271-21276.
5. Lloyd-Hughes B, Shaffrey LC, Vidale PL, & Arnell NW (2012) An evaluation of the spatiotemporal structure of large-scale European drought within the HiGEM climate model. *International Journal of Climatology*:n/a-n/a.
6. Sheffield J & Wood EF (2008) Projected changes in drought occurrence under future global warming from multi-model, multi-scenario, IPCC AR4 simulations. *Climate Dynamics* 13:79-105.
7. Dai A (2011) Drought under global warming: A review. *Wiley Interdisciplinary Reviews: Climate Change* 2:45-65, DOI: 10.1002/wcc.1081.
8. Hempel S, Frieler K, Warszawski L, Schewe J, & Piontek F (2013) A trend-preserving bias correction; the ISI-MIP approach. *Earth Syst. Dynam. Discuss.* 4(1):49-92.
9. Taylor CM, de Jeu RAM, Guichard F, Harris PP, & Dorigo WA (2012) Afternoon rain more likely over drier soils. *Nature* 489(7416):423-426.
10. Sheffield J & Wood EF (2011) *Drought - Past problems and future scenarios* (Earthscan, London, Washington) p 210.
11. Seneviratne SI, *et al.* (2010) Investigating soil moisture-climate interactions in a changing climate: A review. *Earth-Science Reviews* 99(3-4):125-161.
12. Gosling SN & Arnell NW (2011) Simulating current global river runoff with a global hydrological model: model revisions, validation, and sensitivity analysis. *Hydrological Processes* 25:1129-1145.
13. Clark DB, *et al.* (2011) The Joint UK Land Environment Simulator (JULES), model description - Part 2: Carbon fluxes and vegetation dynamics. *Geosci. Model Dev.* 4(3):701-722.
14. Haddeland I, *et al.* (2011) Multimodel Estimate of the Global Terrestrial Water Balance: Setup and First Results. *Journal of Hydrometeorology* 12(5):869-884.
15. Prudhomme C, *et al.* (2011) How well do large-scale models reproduce regional hydrological extremes in Europe? *Journal of Hydrometeorology* 12(6):1181-1204.
16. Gudmundsson L, *et al.* (2011) Comparing Large-Scale Hydrological Model Simulations to Observed Runoff Percentiles in Europe. *Journal of Hydrometeorology* 13(2):604-620.
17. Hagemann S, *et al.* (2012) Climate change impact on available water resources obtained using multiple global climate and hydrology models. *Earth Syst. Dynam. Discuss.* 3(2):1321-1345.
18. Warszawski L (submitted this issue) Research Design of the Intersectoral Impact Model Intercomparison Project (ISI-MIP). *Proceedings of the National Academy of Sciences*.
19. Taylor KE, Stouffer RJ, & Meehl GA (2011) An Overview of CMIP5 and the Experiment Design. *Bulletin of the American Meteorological Society* 93(4):485-498.
20. Gudmundsson L, Wagener T, Tallaksen LM, & Engeland K (2012) Evaluation of nine large-scale hydrological models with respect to the seasonal runoff climatology in Europe. *Water Resources Research* 48:W11504.
21. Hagemann S, *et al.* (2011) Impact of a Statistical Bias Correction on the Projected Hydrological Changes Obtained from Three GCMs and Two Hydrology Models. *Journal of Hydrometeorology* 12(4):556-578.
22. van Vuuren D, *et al.* (2011) The representative concentration pathways: an overview. *Clim. Change* 109(1):5-31.
23. IPCC (2007) Summary for Policymakers. *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Parry ML, Canziani OF, Palutikof JP, Van der Linden PJ, & Hanson CE (Cambridge University Press, Cambridge, UK), pp 7-22.
24. Cao L, Bala G, Caldeira K, Nemani R, & Ban-Weiss G (2010) Importance of carbon dioxide physiological forcing to future climate change. *Proceedings of the National Academy of Sciences* 107(21):9513-9518.

funded by a PhD scholarship from the UK Natural Environment Research Council (NE/YXS1270382). RD was supported by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). YM was supported by the Environment Research and Technology Development Fund (S-10) of the Ministry of the Environment, Japan. Original codes were written by Dr. George Goodsell who is gratefully acknowledged. The manuscript could be improved thanks to constructive comments from two anonymous reviewers who are gratefully acknowledged.

25. Alo CA & Wang GL (2008) Potential future changes of the terrestrial ecosystem based on climate projections by eight general circulation models. *Journal of Geophysical Research-Biogeosciences* 113(G1).
26. Wada Y, *et al.* (2013) Multi-model projections and uncertainties of irrigation water demand under climate change. *Geophysical Research Letters*:n/a-n/a.
27. UNEP (2012) GEO 5 Global Environment Outlook 5 - Environment for the future we want. p 551.
28. Dai A (2013) Increasing drought under global warming in observations and models. *Nature Clim. Change* 3(1):52-58.
29. Burke EJ & Brown SJ (2008) Evaluating uncertainties in the projection of future drought. *Journal of Hydrometeorology* 9(2):292-299.
30. Burke EJ, Brown SJ, & Christidis N (2006) Modeling the Recent Evolution of Global Drought and Projections for the Twenty-First Century with the Hadley Centre Climate Model. *Journal of Hydrometeorology* 7(5):1113-1125.
31. Kirono DGC, Kent DM, Hennessy KJ, & Mpelasoka F (2011) Characteristics of Australian droughts under enhanced greenhouse conditions: Results from 14 global climate models. *Journal of Arid Environments* 75(6):566-575.
32. Orlowsky B & Seneviratne SI (2012) Elusive drought: uncertainty in observed trends and short- and long-term CMIP5 projections. *Hydrol. Earth Syst. Sci. Discuss.* 9(12):13773-13803.
33. Alkama R, Kageyama M, & Ramstein G (2010) Relative contributions of climate change, stomatal closure, and leaf area index changes to 20th and 21st century runoff change: A modelling approach using the Organizing Carbon and Hydrology in Dynamic Ecosystems (ORCHIDEE) land surface model. *Journal of Geophysical Research: Atmospheres* 115(D17):n/a-n/a.
34. Betts RA, *et al.* (2007) Projected increase in continental runoff due to plant responses to increasing carbon dioxide. *Nature* 448(7157):1037-1041.
35. Leakey ADB, *et al.* (2009) Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany* 60(10):2859-2876.
36. Gerten D, *et al.* (2011) Global Water Availability and Requirements for Future Food Production. *Journal of Hydrometeorology* 12(5):885-899.
37. Wang K & Dickinson RE (2012) A review of global terrestrial evapotranspiration: Observation, modeling, climatology, and climatic variability. *Reviews of Geophysics* 50(2):RG2005.
38. De Kauwe MG, *et al.* (2013) Forest water use and water use efficiency at elevated CO₂: a model-data intercomparison at two contrasting temperate forest FACE sites. *Global Change Biology* 19(6):1759-1779.
39. Sitch S, *et al.* (2008) Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs). *Global Change Biology* 14(9):2015-2039.
40. Wada Y, *et al.* (2011) Global monthly water stress: 2. Water demand and severity of water stress. *Water Resources Research* 47(7):W07518.
41. Vörösmarty CJ, Green P, Salisbury J, & Lammers RB (2000) Global Water Resources: Vulnerability from Climate Change and Population Growth. *Science* 289(5477):284-288.
42. Moss RH, *et al.* (2010) The next generation of scenarios for climate change research and assessment. *Nature* 463(7282):747-756.
43. Weedon GP, *et al.* (2011) Creation of the WATCH Forcing Data and Its Use to Assess Global and Regional Reference Crop Evaporation over Land during the Twentieth Century. *Journal of Hydrometeorology* 12(5):823-848.
44. Tallaksen LM & Van Lanen HAJ eds (2004) *Hydrological drought - processes and estimation methods for streamflow and groundwater* (Elsevier), p 579.
45. Orlowsky B & Seneviratne S (2012) Global changes in extreme events: regional and seasonal dimension. *Clim. Change* 110(3):669-696.
46. Corzo Perez GA, van Huijgevoort MHJ, Voß F, & van Lanen HAJ (2011) On the spatiotemporal analysis of hydrological droughts from global hydrological models. *Hydrol. Earth Syst. Sci.* 15(9):2963-2978.
47. Andreadis KM, Clark EA, Wood AW, Hamlet AF, & Lettenmaier DP (2005) Twentieth-Century Drought in the Conterminous United States. *Journal of Hydrometeorology* 6(6):985-1001.
48. Wilks DS (2006) *Statistical methods in the atmospheric science* (Elsevier) p 627.