

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.



For the full text of this licence, please go to:  
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

**Title: Associations of objectively measured sedentary behaviour and physical activity with markers of cardio-metabolic health**

J. Henson, Leicester Diabetes Centre, Leicester General Hospital, UK  
T. Yates, Department of Cardiovascular Sciences, University of Leicester, UK  
S. J. H. Biddle, School of Sport, Exercise and Health Sciences, Loughborough University, UK  
C. L. Edwardson, Leicester Diabetes Centre, Leicester General Hospital, UK  
K. Khunti, Department of Health Sciences, University of Leicester, UK  
E. G. Wilmot, Department of Cardiovascular Sciences, University of Leicester, UK  
L. J. Gray, Department of Health Sciences, University of Leicester, UK  
T. Gorely, School of Sport, Exercise and Health Sciences, Loughborough University, UK  
M. A. Nimmo, School of Sport, Exercise and Health Sciences, Loughborough University, UK  
M. J. Davies, Department of Cardiovascular Sciences, University of Leicester, UK

\* All authors are affiliated with The NIHR Leicester-Loughborough Diet, Lifestyle and Physical Activity Biomedical Research Unit which is a partnership between University Hospitals of Leicester NHS Trust, Loughborough University and the University of Leicester.

**Corresponding Author:**

Mr Joseph Henson  
Leicester Diabetes Centre  
Leicester General Hospital  
Leicester  
LE5 4PW  
UK

**Email address:** [jjh18@le.ac.uk](mailto:jjh18@le.ac.uk)

**Tel:** +44 116 258 4436

**Fax:** +44116 258 4053.

Abstract word count= 240

Main text word count = 3,101

Number of Tables = 2

Number of Figures = 2

## Abstract

*Aims/hypothesis* To examine the associations between objectively measured sedentary time, breaks in sedentary time, moderate-to-vigorous physical activity (MVPA) and total physical activity with markers of cardio-metabolic health in a population at a high risk of type 2 diabetes mellitus.

*Methods* This study reports data from two ongoing diabetes prevention programmes. Participants at a high risk of type 2 diabetes were recruited from primary care practices located within the East Midlands, United Kingdom, 2010-2011, using either a validated risk score or medical records. Sedentary time ( $<100$ counts/min), MVPA ( $\geq 1952$ counts/min) and total physical activity (counts) were measured using Actigraph GT3X accelerometers (15s epochs). A break was considered as any interruption in sedentary time ( $\geq 100$ counts/min). Linear regression analysis examined the associations with markers of cardio-metabolic health.

*Results* 878 participants comprising 153 from 'Project STAND' (age= $32.9 \pm 5.6$ years; female=71.2%) and 725 from 'Walking Away from Diabetes' (age= $63.7 \pm 7.8$ years; female=35.2%). Following adjustment for various covariates, including MVPA, there were detrimental linear associations of sedentary time with 2-hour plasma glucose (standardised beta co-efficient ( $\beta$ )) ( $\beta=0.220$ ,  $p<0.001$ ), triglycerides ( $\beta=0.206$ ,  $p=0.001$ ) and HDL cholesterol ( $\beta=-0.123$ ,  $p=0.029$ ). In contrast, after adjustment for sedentary time, MVPA was significantly inversely associated with BMI ( $\beta=-0.215$ ,  $p<0.001$ ) and waist circumference ( $\beta=-0.228$ ,  $p<0.001$ ), but not with measured biochemical variables. Breaks in sedentary time were inversely associated with waist circumference ( $\beta=-0.215$ ,  $p<0.001$ ) and BMI ( $\beta=-0.151$ ,  $p=0.001$ ). Total physical activity was

*Conclusions/Interpretation* In adults at high risk of type 2 diabetes mellitus, time spent sedentary is strongly and adversely associated with cardio-metabolic health. Furthermore, sedentary time may be a more important indicator of poor health than MVPA.

Walking Away from Type 2 Diabetes Study - **ISRCTN31392913**.

Project STAND (Sedentary Time And Diabetes) - **ISRCTN08434554**.

**Keywords:** Breaks in sedentary time, High-risk, Physical activity, Primary care, Sedentary behaviour, Type 2 diabetes mellitus.

### Abbreviations:

$\beta$ - Beta-coefficient

cpm- Counts per minute

CVD- Cardiovascular disease

FPG- Fasting plasma glucose

IFG- Impaired fasting glycaemia

IGR- Impaired glucose regulation

IGT- Impaired glucose tolerance

IMD- Index of multiple deprivation

MVPA- Moderate-to-vigorous physical activity

METs- Metabolic equivalents

STAND- Sedentary Time And Diabetes

UK- United Kingdom

## Introduction

Sedentary behaviour has previously been characterised as  $\leq 1.5$  metabolic equivalents (METs) [1, 2]. METs are the energy cost of physical activity and are expressed as multiples of resting metabolic rate- where one MET (or  $3.5 \text{ ml min}^{-1} \text{ kg}^{-1}$ ) is equivalent to a typical metabolism at rest for an average person. Given the fact it is impractical to measure energy expenditure in most studies and there are limited behaviours that involve both sitting and energy expenditure ( $>1.5$  METs), a more operational behavioural interpretation has been recommended, where sedentary behaviour is defined as any non-exercise sitting time [3]. Over the last decade, sedentary behaviour has emerged as a distinctive behavioural paradigm with detrimental effects on chronic disease risk, independent of moderate- to vigorous-intensity physical activity (MVPA) [4-8]. This new paradigm is conceptualised around two constructs: total time spent sedentary and the number of breaks in sedentary behaviour (e.g. rising from a sitting/lying position to standing). Both expressions show strong associations with markers of cardio-metabolic health independent of each other and other lifestyle behaviours [4-6, 8, 9].

Epidemiological evidence examining the effect of time spent sedentary has tended to focus on self-report measures [6, 10-12], which are prone to bias and have poor levels of validity [13]. Although several studies employing objective measures of sedentary behaviour have been reported, most have been conducted in the general population [4, 5, 8, 14, 15]. It is therefore unclear to what extent the reported associations are generalisable to those at high risk of chronic disease. This is an important limitation as international recommendations and policies specify that chronic disease prevention strategies should include targeted interventions aimed at the identification and management of high risk individuals [16-18]. Therefore the importance of sedentary behaviour in this group needs to be better understood in order to inform the content and structure of prevention programmes.

The effect of age on the association between sedentary time and cardio-metabolic risk also remains unclear. Many studies involving children and adolescents have reported minimal or no associations between sedentary time and markers of cardio-metabolic health and evidence in young adults remains limited, particularly in high risk populations [19, 20].

In this study, we examined the extent to which sedentary time, breaks in sedentary time, and MVPA are independently associated with cardio-metabolic risk factors in a population identified with a high risk of developing type 2 diabetes mellitus. We hypothesised that all three constructs would be independently associated with health.

## Methods

*Participants* This study used combined baseline data two prevention studies, Walking Away from Type 2 Diabetes Study (WA) (ISRCTN31392913) and Project STAND (Sedentary Time And Diabetes) (ISRCTN08434554), 2010-2011. Both trial protocols have been published elsewhere [21, 22]. Briefly, WA is a randomised controlled trial investigating whether a lifestyle intervention programme can promote behaviour change in those identified at high risk of type 2 diabetes mellitus. Similarly, Project STAND is a randomised controlled trial investigating the effect of structured education and self monitoring on reducing sedentary time in young adults identified at high risk of developing type 2 diabetes mellitus.

Individuals were unaware of their diabetes risk status before entering both studies and all participants were excluded if they had known type 2 diabetes mellitus or were taking steroids. Baseline measurements across both studies were performed before treatment allocation by the same team of trained staff who followed identical standard operating procedures. Informed consent obtained from all eligible participants and both studies gained full ethical and governance approvals from the Nottingham Research Ethics Committee and Leicestershire, Northamptonshire and Rutland Comprehensive Local Research Network.

*Walking Away* Middle age and older adults (aged up to 74) were recruited from 10 primary care practices within Leicestershire, United Kingdom (UK). Individuals at high risk of impaired glucose regulation (IGR) (composite of impaired glucose tolerance (IGT) and/or impaired fasting glycaemia (IFG)) or type 2 diabetes mellitus were identified using a modified version of the automated Leicester Risk Score, specifically designed to be administered in primary care [23]. An automated platform using medical records was used to rank individuals for diabetes risk using predefined weighted variables (age, gender, BMI, family history of type 2 diabetes mellitus and use of antihypertensive medication). Those scoring within the 90<sup>th</sup> percentile in each practice were invited to take part in the study. This approach has been shown to have reasonable sensitivity and specificity for identifying participants at a high risk of IGR [23].

*Project STAND* Young adults who were at risk of developing type 2 diabetes mellitus from across Leicestershire and the South East Midlands region were recruited from primary care practices. Practices databases were searched for participants meeting the following inclusion criteria: a) aged 18-40 years with a BMI in the obese range ( $\geq 30\text{kg/m}^2$ ;  $\geq 27.5\text{kg/m}^2$  for south Asians) or b) aged 18-40 years with a BMI in the overweight range  $\geq 25\text{kg/m}^2$  ( $\geq 23\text{kg/m}^2$  for south Asians) plus one additional risk factor of a family history of type 2 diabetes mellitus or cardiovascular disease (CVD), previous gestational diabetes, polycystic ovarian syndrome, HbA1c  $\geq 5.8\%$  or IGR [24].

*Cardiovascular, metabolic and anthropometric outcomes* Markers of metabolic and cardiovascular health were measured, including fasting plasma glucose (FPG) and 2-hour plasma glucose (via an OGTT), HbA1c, total cholesterol, HDL cholesterol and triglycerides. Venous blood samples were obtained following an overnight fast and all assays were measured in the same laboratory located within the Leicester Royal Infirmary, UK. Analysis was conducted by individuals blinded to the patients' identity, using stable methodologies, standardised to external quality assurance values. Plasma glucose was analysed in venous samples via the hexokinase method. HbA1c was analysed using the Bio-Rad Variant II HPLC system (Bio-Rad Clinical Diagnostics, Hemel Hempstead, UK). HDL-cholesterol and triglycerides were measured using standard enzymatic techniques.

Body weight (Tanita TBE 611, Tanita, West Drayton, UK) and waist circumference (midpoint between the lower costal margin and iliac crest) were measured to the nearest 0.1 kg and 0.5 cm respectively. Information on current smoking status, medication and ethnicity was obtained following an interview administered protocol with a health care professional. Social deprivation was determined by assigning an Index of Multiple Deprivation (IMD) score to the participant's resident area [25]. IMD scores are publically available continuous measures of compound social and material deprivation which are calculated using a variety of data including current income, employment, education and housing.

*Accelerometer measures* Participants were asked to wear a tri-axial accelerometer (Actigraph GT3X, Florida, USA) on the right midaxillary line of the hip (attached via a waistband), for a minimum of seven consecutive days during waking hours. These accelerometers translate raw accelerations into activity counts and Freedson cut-offs were used to categorise MVPA ( $\geq 1952$  counts per minute (cpm)) and sedentary behaviour ( $< 100$ cpm) [26]. A break was considered as an interruption in sedentary time in which the accelerometer count rose up to or above 100cpm [4, 8]. Accelerometers were initialised to record activity in 5s epochs in the STAND cohort and 15s epochs in the WA cohort. In order to allow for direct comparison, all data was analysed using 15s epochs, following epoch preprocessor conversion in the STAND cohort.

Non-wear time was defined as a minimum of 60 minutes of continuous zero counts and days with at least 600 minutes wear time were considered valid [4, 5, 14]. In order for data to be included in the analysis, participants required at least four valid days [27].

A physical activity data analysis tool (ActiSCi, Suffolk, UK) was used to derive wear time, apply cut-points for sedentary time and MVPA and undertake epoch preprocessor conversions (5s to 15s).

*Statistical analysis* Statistical analyses were performed using PASW Statistics v18.0. Due to their skewed distribution, FPG, 2-hour glucose, HDL-cholesterol, triglycerides and HDL: total cholesterol ratio were log-transformed.

Forced-entry linear regression analysis was used on the combined study cohorts to examine independent associations of sedentary time, breaks in sedentary time and MVPA with markers of metabolic (FPG, 2-hour glucose, waist circumference, BMI, HbA1c) and cardiovascular health (triglycerides, HDL cholesterol and HDL: total cholesterol ratio).

Model 1 was adjusted for age, gender, smoking status, ethnicity, social deprivation, lipid lowering and beta-blocker medication, family history of type 2 diabetes mellitus, days accelerometer worn and time accelerometer worn (average minutes per day). Model 2 was additionally adjusted for MVPA time (minutes per day) and the associations for breaks and MVPA were examined having also adjusted for sedentary time (minutes per day). In order to examine the extent to which adiposity may attenuate these relationships, model 3 was further adjusted for BMI.

Significant associations were followed up with interaction terms to assess differences in the strength of the associations between sedentary time, breaks in sedentary time and MVPA by study group and sex, using a model adjusted for the above covariates. To further represent the strength of sedentary time and breaks in sedentary time with cardio-metabolic markers, variables were also examined as tertiles using analysis of covariance (ANCOVA) procedures.

In order to enable direct comparison to previous published studies, a sensitivity analysis was conducted to investigate whether results were affected by integrating the measure of sedentary time to 60s epochs.

Two-tailed  $p$  values of 0.05 or less were considered statistically significant for main effects. Adjustment was not made for multiple comparisons, therefore data were viewed with caution and in relation to the overall pattern of results.  $p < 0.1$  was considered significant for interactions. Due to log-transformation, and to allow for direct comparisons across cardio-

metabolic markers, results of the linear regression analysis are presented as the standardised beta co-efficient ( $\beta$ ) $\pm$ SE.

## Results

In total, 153 younger participants from Project STAND (age=32.9 $\pm$ 5.6years; female=71.2%) and 725 older participants from WA (age=63.7 $\pm$ 7.8years; female=35.2%) had valid measures of both objective activity and biochemical variables. This equated to 87% of the combined cohort. The majority of excluded participants failed to meet the minimum accelerometer wear time requirement. Those included in this analysis had a similar ethnic breakdown and social deprivation score compared to those who did not reach the minimum accelerometer criteria. However, those excluded were more likely to be younger (51.3 $\pm$ 14.6 vs. 58.4 $\pm$ 13.8years,  $p$ <0.001) with a larger waist circumference (105.4 $\pm$ 15.4 vs. 101.6 $\pm$ 12.0cm,  $p$ <0.001) and BMI (34.6 $\pm$ 6.7 vs. 32.5 $\pm$ 5.2kg/m<sup>2</sup>,  $p$ <0.001). Table 1 reports the demographic, cardio-metabolic, anthropometric and accelerometer characteristics of included participants.

Accelerometer wear time (14.5 $\pm$ 1.4 vs. 14.4 $\pm$ 1.4hours per day) and sedentary time (10.3 $\pm$ 1.5 vs. 10.3 $\pm$ 1.5hours) were similar between study cohorts. The younger STAND cohort spent a longer time engaged in MVPA (interquartile range; 0.7 (0.4-0.9) vs. 0.5 (0.3-0.9hours)). Sedentary time and breaks showed small correlations in the STAND ( $r$ =-0.208,  $p$ <0.01) and WA ( $r$ =-0.106,  $p$ =0.01) cohorts. MVPA had an inverse correlation with sedentary time (STAND;  $r_s$ =-0.328,  $p$ <0.001; WA;  $r_s$ =-0.363,  $p$ <0.001) and a small association with breaks (STAND;  $r_s$ =0.246,  $p$ <0.01; WA;  $r_s$ =0.215,  $p$ <0.001).

*Sedentary Time* Table 2 displays the overall associations in the combined cohort of total sedentary time with biomedical and anthropometric markers. After adjustments for known confounders, including MVPA and BMI, sedentary time showed a detrimental association with 2-hour glucose ( $\beta$ =0.220 $\pm$ 0.060,  $p$ =<0.001), HDL-cholesterol ( $\beta$ =-0.123 $\pm$ 0.056,  $p$ =0.029) and triglycerides ( $\beta$ =0.206 $\pm$ 0.061,  $p$ =0.001).

*MVPA* MVPA was significantly inversely associated with 2-hour glucose ( $\beta$ =-0.121 $\pm$ 0.035,  $p$ =<0.001), triglycerides ( $\beta$ =-0.149 $\pm$ 0.036,  $p$ =<0.001), total cholesterol:HDL ratio ( $\beta$ =-0.124 $\pm$ 0.034,  $p$ =<0.001), HDL cholesterol ( $\beta$ =0.150 $\pm$ 0.033,  $p$ =<0.001), BMI ( $\beta$ =-0.241 $\pm$ 0.031,  $p$ =<0.001) and waist circumference ( $\beta$ =-0.270 $\pm$ 0.033,  $p$ =<0.001). However, after adjustment for sedentary time, only BMI ( $\beta$ =-0.215 $\pm$ 0.041,  $p$ =<0.001) and waist circumference ( $\beta$ =-0.228 $\pm$ 0.043,  $p$ =<0.001) remained significant.

*Breaks in sedentary time* Table 2 also reports the associations of breaks in sedentary time with the same markers of health. Independent of known confounders (including sedentary time), breaks in sedentary behaviour were significantly inversely associated with 2-hour glucose ( $\beta$ =-0.111 $\pm$ 0.055,  $p$ =0.046) waist circumference ( $\beta$ =-0.215 $\pm$ 0.051,  $p$ =<0.001) and BMI ( $\beta$ =-0.151 $\pm$ 0.049,  $p$ =0.003). However, further adjustment for BMI attenuated the association with 2-hour glucose.

Results reported above were unaffected if waist circumference rather than BMI was used in Model 3.

Figure 1 illustrates the associations between sedentary time and 2-hour glucose, HDL-cholesterol and triglycerides when examined as tertiles. Figure 2 shows the association of breaks with waist circumference and BMI.

Interaction analyses indicated a significant effect for study cohort, with the older cohort demonstrating greater associations of MVPA with BMI ( $p$  for interaction  $<0.001$ ) and waist circumference ( $p$  for interaction  $<0.001$ ). For breaks in sedentary time, the same pattern was observed, with the older cohort achieving a stronger association for waist circumference ( $p$  for interaction  $=<0.001$ ) and BMI ( $p$  for interaction  $=<0.001$ ). No other significant interactions were observed for the effect of study group. In addition, there were no significant interactions for sex in the results for sedentary time, MVPA or breaks.

The pattern of results and significance levels were unaffected if data were analysed at 60s epochs. However, standardised beta-coefficients were consistently around 10% lower reflecting the less sensitive nature of the data at longer epochs (data available on request).

## Discussion

This study of those at a high risk of type 2 diabetes mellitus recruited from primary care found that sedentary time was detrimentally associated with 2-hour glucose, triglycerides and HDL-cholesterol, independent of measured confounders. These results remained significant after further adjustment for measures of adiposity. Furthermore, the findings for biochemical factors were consistent across groups with diverse age ranges, providing evidence that the deleterious consequences of excess sedentary time exist across young to old adults. Interestingly, sedentary time was shown to have stronger associations with several important cardio-metabolic markers (2-hour glucose, triglycerides, HDL-cholesterol) than MVPA, after adjustment for each other and other important known confounders. Associations of breaks in sedentary time with markers of health, independent of overall time spent sedentary and in MVPA, were less consistent, although beneficial associations were observed with measures of adiposity. To our knowledge, this is the first study to examine the effect of sedentary time and breaks on markers of cardio-metabolic health in a primary care population at high risk of type 2 diabetes mellitus.

Our study has multiple strengths; most notably it provides novel evidence in a high risk population recruited through primary care using an objective measure of sedentary time, across a wide age range. Furthermore, all participants were from the same geographical location, with similar risk profiles and measurements across both studies were performed by the same team of trained staff, following identical standard operating procedures. In addition, participants were rigorously phenotyped with traditional markers of cardio-metabolic health using standardised biochemical procedures. Limitations include the cross-sectional design which limits inference about the direction of causality between the sedentary variables and markers of cardio-metabolic health; reverse causality remains a possibility. Despite allowing for more robust assessments of sedentary behaviour compared to self-report, accelerometers are not without limitations. For example, they rely on categorising movement (acceleration) strength, rather than directly distinguishing between sitting, lying and standing behaviours.

Our results extend those from other studies that have utilised both self-reported and objective measures of sedentary time and MVPA with cardio-metabolic variables in the general population. Self-reported sedentary behaviour in the form of television viewing time has been positively associated with a multitude of cardio-metabolic risk factors [6, 28-30], including 2-hour glucose [28, 29]. Similarly, recent reviews also report that self-reported sedentary time is consistently associated with an increased risk of diabetes [9] and the metabolic syndrome [31].



Several studies have examined the joint effect of sedentary behaviour and physical activity on health outcomes [5, 32, 33]. In contrast to our observations, most have concluded that physical activity is a stronger predictor of metabolic risk [32] and insulin resistance [33]. This discrepancy in findings may be due to differences in study populations, as our participants had been identified as being at high risk of type 2 diabetes mellitus, were largely obese and spent longer in sedentary pursuits. Our population is representative of those that are likely to be identified as being at high risk of type 2 diabetes mellitus within routine care and referred onto available prevention programmes.

Our study was not able to corroborate previous findings that breaks in sedentary behaviour are independently associated with glucose regulation and triglycerides [8]. However, in contrast to overall sedentary behaviour, strong associations were observed between breaks and measures of adiposity (waist circumference and BMI), even after adjustment for MVPA and total time spent sedentary. Therefore, this study suggests that breaks in sedentary behaviour, rather than total sedentary time *per se* may be an important factor in the regulation of body weight.

The non-significant results observed for FPG and HbA1c are consistent with previous research on physical activity and sedentary behaviour [4, 14, 34] and reflect the different pathophysiological process underlying 2-hour and FPG regulation, with 2-hour glucose largely influenced by peripheral insulin resistance [34, 35]. Our findings, therefore, highlight the importance of using 2-hour glucose as the primary outcome variable when assessing the impact of sedentary time on cardio-metabolic risk.

Animal models have confirmed that distinctive physiological pathways are activated with increased sedentary behaviour, particularly around the metabolism of lipoprotein lipase, which remains largely unaffected by MVPA [36]. Lipoprotein lipase is a key regulator of lipid metabolism and is causally linked to CVD [37]. In addition, sedentary behaviour may also reduce glucose transporter protein content, thus exacerbating insulin resistance [38].

Whilst the mounting epidemiological evidence linking sedentary behaviour with cardio-metabolic variables is supported by plausible mechanistic pathways that have been tested in animal models, published experimental research in humans is largely lacking. To date, only one intervention study focused specifically on sedentary behaviour in adults has been published. Nineteen overweight/obese adults showed large reductions in the area under the glucose and insulin curve when sitting time was regularly punctuated with short periods of both light and moderate intensity activity [39]. Surprisingly, there was no difference between the effect sizes found in the light or moderate intensity profiles. Although encouraging, the findings from this study need to be confirmed in different populations in order to establish a causal link between sedentary behaviour and cardio-metabolic dysfunction.

In conclusion, the findings from this study may have important methodological and public health implications. This study provides novel objective evidence that in individuals at high risk of type 2 diabetes mellitus, sedentary time may be a more important indicator of cardio-metabolic health than MVPA. Therefore, diabetes and cardiovascular prevention programmes concentrating solely on MVPA may overlook an area that is of fundamental importance to cardio-metabolic health. Along with messages around accumulating at least 150 minutes per week of MVPA, which form the corner-stone of diabetes prevention programmes [40], such interventions may be more effective if individuals are further encouraged to reduce their sedentary time and to break up periods of prolonged sitting. Nevertheless, given the

limitations, this study should not be used to confirm a direct link between sitting time and metabolic health, but should act as a stimulus for tightly controlled intervention studies in different populations in order to influence future physical activity and sedentary behaviour interventions and public health initiatives aimed at disease prevention.

### **Acknowledgements**

The research was supported by the National Institute for Health Research (NIHR) Leicester-Loughborough Diet, Lifestyle and Physical Activity Biomedical Research Unit based at the University Hospitals of Leicester, University of Leicester and Loughborough University. The research (was also partly funded by) and took place at the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care based at the University of Leicester. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

### **Funding**

Project STAND is funded by the Medical Research Council and National Prevention Research Initiative funding partners (MRC Project no.91409).

### **Duality of interest**

The authors declare that there is no duality of interest associated with this manuscript.

**Author Contributions:** JH and TY had original idea for the analysis. JH wrote the first draft of the article. JH, TY, SB, KK, EW, MN and MJD made substantial contributions to conception and design. JH and CE carried out acquisition of data. JH, CE and TG processed raw accelerometer files. JH, TY and LG carried out analysis and interpretation of data. All authors reviewed/edited the manuscript and gave final approval of the version to be published.

## References

1. Pate RR, O'Neill JR, Lobelo F (2008) The evolving definition of "sedentary". *Exerc Sport Sci Rev* 36:173-178
2. Tremblay MS, Colley RC, Saunders TJ, Healy GN, Owen N (2010) Physiological and health implications of a sedentary lifestyle. *Appl Physiol Nutr Metab* 35:725-740
3. Sedentary Behaviour Research Network (2012) Letter to the Editor: Standardized use of the terms "sedentary" and "sedentary behaviours". *Appl Physiol Nutr Metab* 37:540-542
4. Healy GN, Matthews CE, Dunstan DW, Winkler EA, Owen N (2011) Sedentary time and cardio-metabolic biomarkers in US adults: NHANES 2003-06. *Eur Heart J* 32:590-597
5. Healy GN, Wijndaele K, Dunstan DW, et al. (2008) Objectively measured sedentary time, physical activity, and metabolic risk: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care* 31:369-371
6. Hu FB (2003) Sedentary lifestyle and risk of obesity and type 2 diabetes. *Lipids* 38:103-108
7. Katzmarzyk PT, Church TS, Craig CL, Bouchard C (2009) Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc* 41:998-1005
8. Healy GN, Dunstan DW, Salmon J, et al. (2008) Breaks in sedentary time: beneficial associations with metabolic risk. *Diabetes Care* 31:661-666
9. Thorp AA, Owen N, Neuhaus M, Dunstan DW (2011) Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med* 41:207-215
10. Mummery WK, Kolt G, Schofield G, McLean G (2007) Associations between physical activity and other lifestyle behaviors in older New Zealanders. *J Phys Act Health* 4:411-422
11. Sugiyama T, Healy GN, Dunstan DW, Salmon J, Owen N (2008) Is television viewing time a marker of a broader pattern of sedentary behavior? *Ann Behav Med* 35:245-250
12. Besson H, Brage S, Jakes RW, Ekelund U, Wareham NJ (2010) Estimating physical activity energy expenditure, sedentary time, and physical activity intensity by self-report in adults. *Am J Clin Nutr* 91:106-114
13. Rikli RE (2000) Reliability, validity, and methodological issues in assessing physical activity in older adults. *Res Q Exerc Sport* 71:S89-96
14. Healy GN, Dunstan DW, Salmon J, et al. (2007) Objectively measured light-intensity physical activity is independently associated with 2-h plasma glucose. *Diabetes Care* 30:1384-1389
15. Bankoski A, Harris TB, McClain JJ, et al. (2011) Sedentary activity associated with metabolic syndrome independent of physical activity. *Diabetes Care* 34:497-503
16. Chatterton H, Younger T, Fischer A, Khunti K, Programme Development Group (2012) Risk identification and interventions to prevent type 2 diabetes in adults at high risk: summary of NICE guidance. *BMJ* 345:e4624

17. Ceriello A, Colagiuri S (2008) International Diabetes Federation guideline for management of postmeal glucose: a review of recommendations. *Diabet Med* 25:1151-1156
18. Paulweber B, Valensi P, Lindstrom J, et al. (2010) A European evidence-based guideline for the prevention of type 2 diabetes. *Horm Metab Res* 42 Suppl 1:S3-36
19. Tremblay MS, LeBlanc AG, Kho ME, et al. (2011) Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *Int J Behav Nutr Phys Act* 8:98
20. Ekelund U, Luan J, Sherar LB, et al. (2012) Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA* 307:704-712
21. Yates T, Davies MJ, Henson J, et al. (2012) Walking away from type 2 diabetes: trial protocol of a cluster randomized controlled trial evaluating a structured education programme in those at high risk of developing type 2 diabetes. *BMC Fam Pract* 13:46
22. Wilmot EG, Davies MJ, Edwardson CL, et al. (2011) Rationale and study design for a randomised controlled trial to reduce sedentary time in adults at risk of type 2 diabetes mellitus: project stand (Sedentary Time AND diabetes). *BMC Public Health* 11:908
23. Gray LJ, Davies MJ, Hiles S, et al. (2012) Detection of impaired glucose regulation and/or type 2 diabetes mellitus, using primary care electronic data, in a multiethnic UK community setting. *Diabetologia* 55:959-966
24. World Health Organisation (2006) Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia. Available from [http://www.who.int/diabetes/publications/diagnosis\\_diabetes2006/en/index.html](http://www.who.int/diabetes/publications/diagnosis_diabetes2006/en/index.html), accessed 13 December 2011
25. The English Indices of Deprivation: Summary (2007). Available from <http://www.communities.gov.uk/documents/communities/pdf/576659.pdf>, accessed 11 January, 2012
26. Freedson PS, Melanson E, Sirard J (1998) Calibration of the Computer Science and Applications, Inc. accelerometer. *Med Sci Sports Exerc* 30:777-781
27. Trost SG, McIver KL, Pate RR (2005) Conducting accelerometer-based activity assessments in field-based research. *Med Sci Sports Exerc* 37:S531-43
28. Dunstan DW, Salmon J, Healy GN, et al. (2007) Association of television viewing with fasting and 2-h postchallenge plasma glucose levels in adults without diagnosed diabetes. *Diabetes Care* 30:516-522
29. Healy GN, Dunstan DW, Salmon J, Shaw JE, Zimmet PZ, Owen N (2008) Television time and continuous metabolic risk in physically active adults. *Med Sci Sports Exerc* 40:639-645
30. Ford ES, Schulze MB, Kroger J, Pischon T, Bergmann MM, Boeing H (2010) Television watching and incident diabetes: Findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. *J Diabetes* 2:23-27
31. Edwardson CL, Gorely T, Davies MJ, et al. (2012) Association of sedentary behaviour with metabolic syndrome: a meta-analysis. *PLoS One* 7:e34916
32. Ekelund U, Griffin DJ, Wareham NJ (2007) Physical activity and metabolic risk in individuals with a family history of type 2 diabetes. *Diabetes Care* 30:337-342
33. Ekelund U, Brage S, Griffin SJ, Wareham NJ, ProActive UK Research Group (2009) Objectively measured moderate- and vigorous-intensity physical activity

- but not sedentary time predicts insulin resistance in high-risk individuals. *Diabetes Care* 32:1081-1086
34. Balkau B, Mhamdi L, Oppert JM, et al. (2008) Physical activity and insulin sensitivity: the RISC study. *Diabetes* 57:2613-2618
  35. Faerch K, Borch-Johnsen K, Holst JJ, Vaag A (2009) Pathophysiology and aetiology of impaired fasting glycaemia and impaired glucose tolerance: does it matter for prevention and treatment of type 2 diabetes? *Diabetologia* 52:1714-1723
  36. Hamilton MT, Hamilton DG, Zderic TW (2007) Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 56:2655-2667
  37. Mead JR, Irvine SA, Ramji DP (2002) Lipoprotein lipase: structure, function, regulation, and role in disease. *J Mol Med (Berl)* 80:753-769
  38. Hamilton MT, Hamilton DG, Zderic TW (2004) Exercise physiology versus inactivity physiology: an essential concept for understanding lipoprotein lipase regulation. *Exerc Sport Sci Rev* 32:161-166
  39. Dunstan DW, Kingwell BA, Larsen R, et al. (2012) Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care* 35:976-983
  40. World Health Organisation (2007) Global recommendations on physical activity for health. Available from [http://whqlibdoc.who.int/publications/2010/9789241599979\\_eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241599979_eng.pdf), accessed 7 February 2012.

## Tables

**Table 1** Demographics, metabolic, anthropometric and accelerometer characteristics of participants

<i>Characteristics</i>	STAND (N=153)	Walking Away (N=725)	All (N=878)
Age (years)	32.9 ± 5.6	63.7 ± 7.8	58.4 ± 13.8
Male	44 (28.8)	470 (64.8)	514 (58.5)
Current Smokers	57 (37.3)	62 (8.5)	119 (13.6)
Family History of Diabetes (1 <sup>st</sup> degree)	100 (65.4)	261 (35.9)	361 (41.1)
<i>Cardio-metabolic variables</i>			
BMI (kg/m <sup>2</sup> )	34.6 ± 4.8	31.2 ± 5.3	32.5 ± 5.2
Waist circumference (cm)	102.9 ± 13.5	101.3 ± 11.7	101.6 ± 12.0
Weight (kg)	98.3 ± 17.3	91.5 ± 16.5	92.7 ± 16.9
FPG (mmol/L)	4.8 (4.5-5.1)	5.2 (4.9-5.6)	5.1 (4.8-5.5)
2-hour plasma glucose (mmol/L)	5.3 (4.5-6.4)	6.1 (4.9-7.8)	5.9 (4.8-7.5)
Body Fat (%)	40.5 ± 7.2	35.6 ± 8.7	36.5 ± 8.6
Total Cholesterol (mmol/L) <sup>a</sup>	4.8 (4.2-5.4)	5.1 (4.3-5.9)	5.0 (4.3-5.8)
HDL Cholesterol (mmol/L) <sup>a</sup>	1.2 (1.0-1.4)	1.4 (1.1-1.6)	1.3 (1.1-1.6)
Total Cholesterol: HDL Ratio (mmol/L) <sup>a</sup>	3.9 (3.2-4.7)	3.7 (3.0-4.4)	3.8 (3.0-4.5)
Triglycerides (mmol/L) <sup>a</sup>	1.30 (0.90-1.70)	1.30 (1.00-1.80)	1.30 (1.00-1.80)
Lipid Lowering Medication	1 (0.6)	240 (33.1)	241 (27.4)
Beta-blockers	2 (1.3)	127 (15.2)	129 (14.7)
HbA1c (%)	5.6 (5.3-5.8)	5.9 (5.6-6.1)	5.8 (5.6-6.1)
<i>Ethnicity</i>			
White European	128 (83.7)	645 (89.0)	773 (88.0)
South Asian	15 (9.8)	53 (7.3)	68 (7.8)
Other	10 (6.5)	27 (3.7)	37 (4.2)
<i>Diagnosis</i>			
Normal glucose tolerance	137 (89.5)	512 (70.6)	649 (73.9)
Isolated IFG	3 (1.9)	38 (5.2)	41 (4.7)
Isolated IGT	11 (7.2)	124 (17.1)	135 (15.4)
Both	1 (0.7)	31 (4.3)	32 (3.6)
Type 2 diabetes mellitus	1 (0.7)	20 (2.8)	21 (2.4)
All (IGR)	16 (10.4)	214 (29.5)	230 (26.2)
<i>Accelerometer variables (time in hours)</i>			
Time accelerometer worn	14.5 ± 1.4	14.4 ± 1.4	14.4 ± 1.4
Sedentary Time <sup>b</sup>	10.3 ± 1.5	10.3 ± 1.5	10.3 ± 1.5
Light-intensity activity <sup>b</sup>	3.5 ± 0.9	3.5 ± 0.9	3.5 ± 0.9
MVPA <sup>b</sup>	0.7 (0.4-0.9)	0.5 (0.3-0.9)	0.6 (0.3-0.9)
<i>Accelerometer variables (percent at each activity level)</i>			
Sedentary time	70.5 ± 7.6	71.5 ± 7.8	71.0 ± 8.0
Light-intensity activity	24.3 ± 6.5	24.3 ± 6.3	24.3 ± 6.4
MVPA	4.7 (2.8-6.2)	3.7 (2.1-5.9)	3.9 (2.3-6.0)
Breaks per day	297 ± 68	273 ± 60	277 ± 62
Average steps per day	7153 ± 2954	6993 ± 3384	7016 ± 3313

<sup>a</sup>Means adjusted for lipid lowering medication

<sup>b</sup>Means adjusted for the time the accelerometer was worn

Sedentary time = <100 cpm, light intensity activity 100-1,951 cpm, and MVPA ≥1952 cpm  
Continuous parametric results as mean±SD, number (column percentage) and continuous non-parametric results as median (interquartile range)

**Table 2** Multiple linear regression models for sedentary time, moderate-to-vigorous physical activity and breaks in sedentary time with cardio-metabolic variables

Model 1						
	Sedentary Time ( $<100\text{cpm}$ ) $\beta$ (SE)	$p$ value	MVPA ( $\geq 1952\text{cpm}$ ) $\beta$ (SE)	$p$ value	Breaks $\beta$ (SE)	$p$ value
2-hour glucose (mmol/L)	0.238 (0.045)	$<0.001$	-0.121 (0.035)	0.001	-0.180 (0.038)	$<0.001$
Waist circumference (cm)	0.250 (0.043)	$<0.001$	-0.270 (0.033)	$<0.001$	-0.198 (0.037)	$<0.001$
BMI ( $\text{Kg/m}^2$ )	0.210 (0.041)	$<0.001$	-0.241 (0.031)	$<0.001$	-0.148 (0.035)	$<0.001$
Triglycerides (mmol/L)	0.217 (0.045)	$<0.001$	-0.149 (0.036)	$<0.001$	-0.150 (0.040)	$<0.001$
Fasting Glucose (mmol/L)	0.046 (0.045)	0.308	-0.033 (0.036)	0.488	-0.024 (0.038)	0.777
Total Cholesterol:HDL Ratio	0.130 (0.043)	0.003	-0.124 (0.034)	$<0.001$	-0.114 (0.037)	0.003
HDL Cholesterol (mmol/L)	-0.187 (0.042)	$<0.001$	0.150 (0.033)	$<0.001$	0.130 (0.036)	$<0.001$
HbA1c (%)	0.035 (0.050)	0.489	-0.034 (0.046)	0.464	-0.021 (0.038)	0.590
Model 2						
	Sedentary Time ( $<100\text{cpm}$ ) $\beta$ (SE) <sup>a</sup>	$p$ value	MVPA ( $\geq 1952\text{cpm}$ ) $\beta$ (SE) <sup>b</sup>	$p$ value	Breaks $\beta$ (SE) <sup>c</sup>	$p$ value
2-hour glucose (mmol/L)	0.235 (0.060)	$<0.001$	-0.033 (0.047)	0.473	-0.111 (0.055)	0.046
Waist circumference (cm)	0.091 (0.057)	0.113	-0.228 (0.043)	$<0.001$	-0.215 (0.051)	$<0.001$
BMI ( $\text{Kg/m}^2$ )	0.054 (0.053)	0.327	-0.215 (0.041)	$<0.001$	-0.151 (0.049)	0.003
Triglycerides (mmol/L)	0.214 (0.062)	0.001	-0.042 (0.048)	0.385	-0.046 (0.056)	0.418
Fasting Glucose (mmol/L)	0.023 (0.062)	0.714	-0.021 (0.048)	0.662	-0.011 (0.038)	0.903
Total Cholesterol:HDL Ratio	0.101 (0.058)	0.085	-0.075 (0.045)	0.096	-0.075 (0.054)	0.167
HDL Cholesterol (mmol/L)	-0.137 (0.056)	0.016	0.083 (0.044)	0.060	0.071 (0.052)	0.175
HbA1c (%)	0.014 (0.051)	0.836	-0.013 (0.036)	0.725	-0.035 (0.056)	0.537
Model 3						
	Sedentary Time ( $<100\text{cpm}$ ) $\beta$ (SE) <sup>a</sup>	$p$ value	MVPA ( $\geq 1952\text{cpm}$ ) $\beta$ (SE) <sup>b</sup>	$p$ value	Breaks $\beta$ (SE) <sup>c</sup>	$p$ value
2-hour glucose (mmol/L)	0.220 (0.060)	$<0.001$	-0.019 (0.055)	0.678	-0.095 (0.056)	0.091
Triglycerides (mmol/L)	0.206 (0.061)	0.001	-0.011 (0.050)	0.826	-0.019 (0.056)	0.736
Fasting Glucose (mmol/L)	0.011 (0.063)	0.857	-0.009 (0.047)	0.850	0.000 (0.050)	0.993
Total Cholesterol:HDL Ratio	0.090 (0.057)	0.120	-0.037 (0.045)	0.412	-0.044 (0.053)	0.408
HDL Cholesterol (mmol/L)	-0.123 (0.056)	0.029	0.041 (0.043)	0.344	0.035 (0.051)	0.495
HbA1c (%)	0.008 (0.062)	0.898	-0.010 (0.046)	0.828	-0.022 (0.054)	0.689


Model 1 was adjusted for age, gender, smoking status, ethnicity, social deprivation, family history, beta blockers, lipid lowering medication and time accelerometer worn

Model 2 was adjusted for the above covariates and <sup>a</sup>MVPA, <sup>b</sup>sedentary time or <sup>c</sup>sedentary time and MVPA


Model 3 was adjusted for the same covariates as Model 2 and BMI

## Figures


(a)



(b)









**Figure 1** Tertiles of sedentary time with 2-hour glucose (**a**), log-triglycerides (**b**) and HDL-cholesterol (**c**). Estimated marginal means are adjusted for age, gender, ethnicity, IMD score, smoking status, family history of type 2 diabetes mellitus, lipid lowering medication, beta blockers, time accelerometer worn, days accelerometer worn, time spent in MVPA and BMI. Tertile cut-points for sedentary time were 9.6 and 10.9hours/day. Medians and ranges for tertile 1 = 8.7hours (2.9-9.5); tertile 2= 10.3hours (9.6-10.9); tertile 3= 11.7hours (11.0-15.8).  $p < 0.001$  for trend (**a**),  $p < 0.05$  for trend (**b**, **c**).

**(a)**



**(b)**



**Figure 2** Tertiles of breaks in sedentary time with waist circumference (**a**) and BMI (**b**). Estimated marginal means are adjusted for age, gender, ethnicity, IMD score, smoking status, family history of type 2 diabetes mellitus, lipid lowering medication, beta blockers, time accelerometer worn, days accelerometer worn and time spent in sedentary and MVPA. Cut points for daily breaks in sedentary time were 234 and 285. Medians and ranges for tertile 1 = 215 (33-234); tertile 2= 268 (235-284); tertile 3= 329 (285-487).  $p < 0.001$  for trend (**a**),  $p < 0.01$  for trend (**b**)