Effect of abdominal binding on respiratory mechanics during exercise in athletes with cervical spinal cord injury

Christopher R. West,¹ Victoria L. Goosey-Tolfrey,² Ian G. Campbell,¹ and Lee M. Romer¹

¹Centre for Sports Medicine and Human Performance, Brunel University, United Kingdom; and ²School of Sport, Exercise & Health Sciences, The Peter Harrison Centre for Disability Sport, Loughborough University, United Kingdom

Submitted 6 March 2014; accepted in final form 15 May 2014

West CR, Goosey-Tolfrey VL, Campbell IG, Romer LM. Effect of abdominal binding on respiratory mechanics during exercise in athletes with cervical spinal cord injury. J Appl Physiol 117: 36-45, 2014. First published May 22, 2014; doi:10.1152/japplphysiol.00218.2014.-We asked whether elastic binding of the abdomen influences respiratory mechanics during wheelchair propulsion in athletes with cervical spinal cord injury (SCI). Eight Paralympic wheelchair rugby players with motor-complete SCI (C5-C7) performed submaximal and maximal incremental exercise tests on a treadmill, both with and without abdominal binding. Measurements included pulmonary function, pressure-derived indices of respiratory mechanics, operating lung volumes, tidal flow-volume data, gas exchange, blood lactate, and symptoms. Residual volume and functional residual capacity were reduced with binding (77 \pm 18 and 81 \pm 11% of unbound, P < 0.05), vital capacity was increased (114 \pm 9%, P < 0.05), whereas total lung capacity was relatively well preserved (99 \pm 5%). During exercise, binding introduced a passive increase in transdiaphragmatic pressure, due primarily to an increase in gastric pressure. Active pressures during inspiration were similar across conditions. A sudden, sustained rise in operating lung volumes was evident in the unbound condition, and these volumes were shifted downward with binding. Expiratory flow limitation did not occur in any subject and there was substantial reserve to increase flow and volume in both conditions. $\dot{V}o_2$ was elevated with binding during the final stages of exercise (8-12%, P < 0.05), whereas blood lactate concentration was reduced (16–19%, P < 0.05). Vo₂/heart rate slopes were less steep with binding (62 \pm 35 vs. 47 \pm 24 ml/beat, P < 0.05). Ventilation, symptoms, and work rates were similar across conditions. The results suggest that abdominal binding shifts tidal breathing to lower lung volumes without influencing flow limitation, symptoms, or exercise tolerance. Changes in respiratory mechanics with binding may benefit O2 transport capacity by an improvement in central circulatory function.

diaphragm; respiratory muscles; tetraplegia; upper-body exercise; wheelchair exercise

INDIVIDUALS WITH CERVICAL spinal cord injury (SCI) exhibit restrictive pulmonary dysfunction, characterized by a significant reduction in lung volumes (51, 53). This restrictive defect has been attributed to weakened respiratory muscles (30), reduced compliance of the lung and chest wall (37), and reduced expanding effect of the diaphragm on the lower rib cage owing to increased abdominal wall compliance (44). During exercise, individuals with cervical SCI demonstrate an immediate and sustained rise in end-expiratory and end-inspiratory lung volumes (i.e., dynamic hyperinflation) (41). This rise in operating lung volumes would be expected to increase the elastic work of breathing, impair the capacity of the

inspiratory muscles to generate pressure, and reduce the relative contribution of the diaphragm to inspiration (41). Cervical SCI also leads to alterations in cardiovascular function during exercise. With complete cervical SCI, maximal heart rate is usually limited to ~120-130 beats/min owing to a lack of supraspinal sympathetic drive to the heart (19). Furthermore, vasomotor tone is impaired owing to a lack of descending sympathetic vascular control (26) and low catecholamine spillover (38). Consequently, blood cannot be redistributed effectively during exercise. This has been associated with venous pooling in nonactive vascular beds (43) and may, in turn, restrict O₂ transport to working muscles by compromising venous return and stroke volume (22). The aforementioned increase in abdominal compliance may further compromise venous return and stroke volume by reducing the abdominothoracic pressure gradient (3, 4).

Previous studies have attempted to increase O₂ transport in individuals with cervical SCI by using a supine position during arm exercise (18, 21), electrical stimulation of lower-limb muscles (14, 17), and application of lower-body positive pressure by means of an antigravity suit (20, 21, 34). An alternative method has been to apply external compression to the abdomen using an elastic binder. This latter approach has been shown to confer multiple benefits at rest, including increases in vital capacity, expiratory flow, respiratory muscle strength, blood pressure, and stroke volume (47, 52). The effects of abdominal binding on exercise responses have been variable (20, 21). These inconsistencies may have stemmed from differences in exercise protocols, exercise modalities, and subject characteristics. In athletes with cervical SCI, we recently showed that abdominal binding increases the distance covered during a field-based endurance test (50). On the basis of a significant positive correlation between distance covered in the field and peak O₂ uptake assessed in the laboratory (r = 0.75, P < 0.05), we considered that the ergogenic effect of binding on endurance performance might have been attributable to an improvement in central circulatory function (50).

The purpose of the present study, therefore, was to better understand the influence of abdominal binding on the acute physiological responses to exercise in athletes with cervical SCI. The specific objective was to determine the effect of abdominal binding on respiratory mechanics during graded wheelchair exercise. Our hypothesis was that abdominal binding would increase intra-abdominal pressure, reduce operating lung volumes, and improve diaphragm function during exercise. We reasoned that these binding-induced changes would improve the circulatory function of the diaphragm, thereby enhancing the overall exercise response through an increase in venous return, cardiac output, and O₂ transport.

Address for reprint requests and other correspondence: L. M. Romer, Centre for Sports Medicine and Human Performance, Brunel Univ., UB8 3PH, UK (e-mail: lee.romer@brunel.ac.uk).

METHODS

Subjects

After providing written informed consent, 8 members of the Great Britain wheelchair rugby squad (1 woman) participated in the study. The subjects had traumatic SCI (2 C_5 , 5 C_6 , 1 C_7) and motor-complete lesions [American Spinal Injury Association Impairment Scale A (n =7) or B (n = 1)]. Subject characteristics (mean \pm SD) were: age 29 \pm 2 yr, stature 1.79 \pm 0.10 m, body mass 67 \pm 15 kg, and time postinjury 9 ± 3 yr. None of the subjects smoked, had a history of cardiopulmonary disease, or were taking medications known to influence the exercise response. At the time of study the subjects were performing at least 15 h/wk of endurance, resistance, and sport-specific training. All of the subjects had taken part in our previous binding studies (50, 52) and were familiar with treadmill exercise testing. The primary outcome measures in the current study do not overlap with previous analyses. Subjects were required to refrain from strenuous exercise for 48 h before testing. Caffeine and alcohol were prohibited for 12 and 24 h, respectively, and no food was allowed within 2 h before testing. Upon arrival at the laboratory, the subjects emptied their bladders to reduce the likelihood of autonomic dysreflexia (12).

Study Design

Subjects visited the laboratory on two separate occasions over a period of ~ 1 wk. Visit 1 included an evaluation of pulmonary function and static respiratory pressures. Visit 2 included submaximal and maximal exercise tests on a treadmill (Fig. 1). The assignment of conditions (unbound and bound) was randomized and counterbalanced. The order of exercise tests was sequential (i.e., submaximal exercise in both conditions, maximal exercise in both conditions). The subjects rested for 30 min between conditions and 60 min between tests. The conditions could not be blinded, but the participants were unaware of the experimental hypotheses and expected outcomes of the study. Cardiopulmonary, metabolic, and perceptual responses were assessed during the submaximal and maximal exercise tests. Because of the invasiveness of the procedures (balloon catheters) and the duration of the experimental visit (~4 h) it was neither feasible nor ethical to measure intrathoracic pressures in both tests; therefore, respiratory mechanics and ventilatory constraint were assessed during submaximal exercise only. The subjects performed all tests in their own sports wheelchair. Gloves were worn for the exercise tests and leg/chest straps if needed. The study procedures received institutional ethical approval and conformed to the Declaration of Helsinki.

Procedures

Abdominal binding. The binder (493R Universal Back Support; McDavid, Woodridge, IL) incorporated a semirigid neoprene back panel with six plastic stays (100% neoprene rubber), flexible sidepanels (90% nylon, 10% Lycra), and a flexible neoprene front panel with double Velcro fastening. The binder was individually sized and fitted in the upright position with the upper edge just beneath the costal margin so that the binder interfered minimally with rib cage movement. An inflatable rubber reservoir with a known volume of air was connected to a digital manometer (C9553; JMW, Harlow, UK) and placed between the binder and the anterior abdominal wall. The binder tightness was adjusted until end-expiratory gastric pressure was approximately twice that in the unbound condition; this level of binding has been shown to optimize resting cardiopulmonary function (52) and improve field-based endurance performance (50). The corresponding abdominal-wall pressure was used to set the binder tightness for the maximal exercise test.

Pulmonary function and static respiratory pressures. Pulmonary volumes, capacities, and flows were assessed using spirometry and body plethysmography (Zan 530; nSpire Health, Oberthulba, Würzburg, Germany) (24, 32, 48). Maximum static inspiratory and expiratory pressures were measured at the mouth (MicroRPM; CareFusion, Basingstoke, UK) from functional residual capacity and total lung capacity, respectively (15).

Exercise tests. Exercise tests were performed on a motorized treadmill with a moving rail to prevent falls (Saturn 300/125r; HP Cosmos, Nussdorf-Traunstein, Germany). The submaximal test consisted of a steady-state resting period followed by four stages, starting at 1.6 m/s and incrementing by 0.4 m/s every 4 min with a 30-s break between stages (27). The maximal test consisted of a fixed speed, chosen according to the responses elicited during the submaximal test, and an increase in gradient of 0.2% every 40 s. The maximal test was terminated when subjects were unable to maintain the treadmill speed, i.e., when they touched the spring of the safety rail for a third time. Standardized verbal encouragement was given throughout the tests, but no information was provided regarding speed, time, or physiological response. Push rate was freely chosen and assessed based on the number of hand-to-rim contacts recorded during the final minute of each stage. After the maximal test the subjects rested for 2 min and then performed an active recovery at low exercise intensity for 5 min. Pretest values were not different at baseline, indicating that the time between tests ensured a full recovery. Power output for each subjectwheelchair combination was determined prior to exercise using a separate drag-test (45).

Cardiopulmonary, metabolic, and perceptual responses. Ventilatory and pulmonary gas exchange variables were assessed breath-bybreath using an online system (Oxycon Pro; Jaeger, Höchberg, Germany). Arterial O₂ saturation was estimated using a pulse oximeter with earlobe sensor (PalmSAT 2500; Nonin Medical, Minneapolis, MN). Heart rate was assessed beat-by-beat via telemetry (Vantage NV; Polar Electro Oy, Kempele, Finland). Earlobe capillary blood was sampled immediately before each test and after each submaximal stage for the determination of lactate concentration in hemolyzed whole-blood (1500 SPORT; YSI Inc, Yellow Springs, OH). After the maximal test, blood was sampled at 0.5, 2, 4, 6, and 8 min and peak lactate concentration was defined as the highest value. Ratings of dyspnea (respiratory discomfort) and arm discomfort were obtained immediately after each stage using Borg's modified 0–10 categoryratio scale (8).

Respiratory mechanics and ventilatory constraint. Gastric pressure (P_{ga}) and esophageal pressure (P_{es}) were measured continuously using previously described procedures (41). Transdiaphragmatic pressure (P_{di}) was obtained by electronic subtraction of P_{es} from P_{ga} . An



Fig. 1. Experimental overview for submaximal and maximal exercise tests.

Abdominal Binding and Respiratory Mechanics In Cervical SCI · West CR et al.

analog airflow signal from the online gas analysis system was simultaneously input into the data acquisition system and aligned to the pressure signals based on the sampling delay for flow. Maximal static inspiratory efforts from functional residual capacity were performed at resting baseline to obtain maximum values for P_{di}, P_{ga}, and P_{es}. To evaluate the passive increase in pressures introduced by application of the binder we report end-expiratory and end-inspiratory values for $P_{\rm di},\,$ Pga, and Pes. To permit comparison of the active pressures generated in both conditions we report inspiratory pressure swings from endexpiratory values, calculated as peak-to-peak (Pdi,tidal, Pga,tidal, Pes,tidal) and integrated pressure-time product (PTP_{di}, PTP_{ga}, PTP_{es}). Dynamic lung compliance during inspiration was calculated as the ratio of tidal volume to Pes,tidal (36). To determine the likelihood of inspiratory muscle fatigue, the tension-time index of the diaphragm (TTI_{di}) was calculated as $P_{di}/P_{di,max}$. T_I/T_{TOT} , where \overline{P}_{di} is mean transdiaphragmatic pressure integrated over inspiration with reference to the endexpiratory level, P_{di,max} is maximum transdiaphragmatic pressure, T_I is inspiratory time, and T_{TOT} is total breath time (7).

The degree of ventilatory constraint was assessed by measuring changes in operating lung volumes, expiratory flow limitation, inspiratory flow reserve, and the ratio of minute ventilation (V_E) to the maximal estimated ventilation for a given breathing pattern (V_{ECAP}), as described previously (5, 23). Briefly, changes in operating lung volumes [end-expiratory lung volume (EELV) and end-inspiratory lung volume (EILV)] were assessed by measuring inspiratory capacity (IC) relative to total lung capacity (TLC), immediately before exercise and during the final 30 s of each submaximal exercise stage [EELV =TLC - IC; EILV = (TLC - IC) + tidal volume]. Peak inspiratory Pes during the IC maneuver was not significantly different across exercise stages in either condition, indicating good reproducibility of maximal efforts for assessment of operating lung volumes. The degree of expiratory flow limitation, if present, was defined as the percent of the tidal flow-volume loop that met or exceeded the expiratory portion of the largest maximal flow-volume loop obtained before or <2 min after exercise (highest sum of FEV_1 and FVC). Inspiratory flow reserve (IFR) was expressed as the peak inspiratory flow generated during tidal breathing relative to that achieved during the maximal flow-volume maneuver at the same lung volume. The level of ventilation relative to a theoretical maximal ventilatory capacity $(\dot{V}_{\rm F})$ V_{ECAP}) was also determined, where V_{ECAP} represents the total area under the expiratory flow curve between EILV and EELV.

Data Analysis

Cardiopulmonary data at rest and during submaximal exercise were averaged over 30-s epochs. To avoid breath contamination from paired IC measurements, the first 30 s of every 4th min of submaximal exercise was analyzed. The 30 s of data used for analysis was filtered to remove outlying breaths, defined as any breath deviating by more than three standard deviations from the mean T_{TOT} during the preceding 5 breaths. Peak cardiopulmonary responses are reported as the highest 30-s average. To determine the degree of expiratory flow limitation, an average breath was constructed for the selected 30-s period by splitting each breath into equal time segments. The number of time segments was based on the mean T_{TOT} with a resolution of 0.01 s. A flow-volume loop was then constructed from the average breath and placed at EELV inside the maximal flow-volume loop for the subsequent assessment of ventilatory constraint.

Statistics

Analyses were performed using SPSS 16.0 for Windows (IBM, Chicago, IL). Data were checked for normality using the Kolmogorov-Smirnov test and homogeneity of variance using Levene's statistic. None of the assumptions underlying parametric testing was violated. Submaximal exercise data were assessed for differences using two-factor (condition \times time) repeated-measures ANOVA. Where a significant interaction effect was detected, post hoc analysis was carried out using Bonferroni-corrected pairwise comparisons. Pulmonary function and maximal exercise data were assessed for differences using two-tailed paired *t*-tests. Pearson's correlation coefficient was calculated to establish correlations between heart rate (dependent variable) and O₂ uptake by subject. The slope and intercept of the equations describing each of these correlations were assessed using linear regression analysis. Critical significance level α was set at 0.05. Values are presented as means \pm SD unless stated otherwise.

RESULTS

Pulmonary Function and Static Respiratory Pressures

Pulmonary function and static respiratory pressures are summarized in Table 1. Abdominal binding increased vital capacity, whereas decreases were noted for functional residual capacity and residual volume. Total lung capacity was relatively well preserved. Forced expiratory volume in 1 s was increased with binding. Maximum inspiratory mouth pressure was not affected by binding, whereas maximum expiratory mouth pressure was increased.

Table 1. Effect of abdominal binding on pulmonary function and static respiratory pressures

	Unbound	Bound	$\%\Delta$
TLC, liters	5.40 ± 1.15	5.38 ± 1.29	-1 ± 5
	(77 ± 9)	(76 ± 10)	
FRC, liters	3.25 ± 0.92	$2.68 \pm 1.01*$	-19 ± 11
	(98 ± 23)	(81 ± 27)	
RV, liters	1.83 ± 1.01	$1.42 \pm 0.99*$	-23 ± 32
	(109 ± 59)	(83 ± 57)	
IC, liters	2.42 ± 0.61	$2.91 \pm 0.69*$	21 ± 7
	(65 ± 7)	(78 ± 9)	
IRV, liters	1.70 ± 0.53	$2.20 \pm 0.58*$	32 ± 14
ERV, liters	1.08 ± 0.38	1.03 ± 0.30	-2 ± 16
	(67 ± 21)	(64 ± 16)	
VC, liters	3.49 ± 0.97	$3.93 \pm 0.94*$	14 ± 9
	(65 ± 10)	(74 ± 10)	
FEV ₁ , liters	2.96 ± 0.81	$3.33 \pm 0.72*$	15 ± 14
	(68 ± 12)	(77 ± 9)	
FEV ₁ /VC, %	84.0 ± 9.8	86.4 ± 7.6	3 ± 7
	(102 ± 11)	(105 ± 9)	
PEF, 1/s	5.8 ± 1.5	6.2 ± 1.6	7 ± 13
	(60 ± 11)	(64 ± 12)	
MEF ₂₅₋₇₅ , 1/s	3.18 ± 1.05	3.81 ± 1.00	28 ± 40
	(65 ± 21)	(78 ± 19)	
MVV ₁₂ , l/min	109 ± 29	111 ± 28	3 ± 13
	(68 ± 17)	(69 ± 18)	
P _{Imax} , cmH ₂ O	-98 ± 45	-103 ± 43	9 ± 20
	(86 ± 33)	(91 ± 32)	
P _{Emax} , cmH ₂ O	59 ± 18	$73 \pm 21*$	26 ± 34
	(43 ± 9)	(53 ± 12)	

Values are means \pm SD for 8 subjects. TLC, total lung capacity; FRC, functional residual capacity; RV, residual volume; IC, inspiratory capacity; IRV, inspiratory reserve volume; ERV, expiratory reserve volume; VC, vital capacity; FEV₁, forced expiratory volume in 1 s; PEF, peak expiratory flow; MEF₂₅₋₇₅, midexpiratory flow between 25 and 75% of VC; MVV₁₂, maximal voluntary ventilation in 12 s; P_{Imax}, maximum static inspiratory pressure from FRC; P_{Emax}, maximum static expiratory pressure from TLC. Values in parentheses are percent of able-bodied predicted values for pulmonary volumes, capacities, and flows (35); MVV (16); and respiratory pressures (10). Predicted values for ERV and IC were derived from differences between corresponding predicted values for FRC and RV, and between TLC and FRC, respectively (35). **P* < 0.05.

Cardiopulmonary, Metabolic, and Perceptual Responses

Responses during the submaximal exercise test are summarized in Table 2. In the unbound condition, the test elicited a wide range of values relative to peak: $\dot{V}o_2$ (64–95%), \dot{V}_E (46–83%), and heart rate (69–90%). There were no differences in ventilation or breathing pattern across conditions. The timing (T_I/T_{TOT}) and drive (V_T/T_I) components of ventilation were also not different across conditions. There was a significant interaction effect between condition and time for $\dot{V}o_2$ (P = 0.002) and blood lactate concentration (P = 0.010), whereby $\dot{V}o_2$ was elevated (8%) and lactate was reduced (19%) in the bound condition during the final stage of the test. The O₂ pulse ($\dot{V}o_2$ /heart rate) was also elevated in the bound condition during the final stage (13.4 ± 2.3 vs. 12.3 ± 2.3 ml/beat, P =0.04). The $\dot{V}o_2$ /heart rate relationship for measurements during submaximal exercise are shown in Fig. 2. The relationships were linear, with high correlations in the unbound and bound condition ($r = 0.933 \pm 0.069$ and 0.967 ± 0.032 , respectively; both P < 0.05). The slopes were less steep in the bound condition (47 ± 24 vs. 62 ± 35 ml/beat, P = 0.022), whereas the intercepts were not different (42 ± 20 vs. 34 ± 22 beat/min, P = 0.149). Perceptual intensities were similar across conditions.

Responses during the maximal exercise test are summarized in Table 3. Peak power output and push rate were not different across conditions. Peak $\dot{V}o_2$ was increased by 12% with binding (P = 0.001), yet peak values for heart rate and minute ventilation were similar across conditions. Thus, peak O_2 pulse was also significantly elevated in the bound condition, whereas, in general, ventilatory equivalents for O_2 (and CO_2) were lower. Peak blood lactate concentration was reduced by 16% in the bound condition (P = 0.052). Perceptual intensities were similar across conditions.

Table 2. Effect of abdominal binding on cardiopulmonary, metabolic, and perceptual responses at rest and during submaximal incremental wheelchair propulsion

Effect	Baseline	Stage 1	Stage 2	Stage 3	Stage 4
Power output, W					
UB	0	20.2 ± 4.5	25.1 ± 5.6	30.1 ± 6.7	35.9 ± 7.9
В	0	20.2 ± 4.5	25.1 ± 5.6	30.1 ± 6.7	35.9 ± 7.9
Push rate, /min					
UB	0	51 ± 11	53 ± 11	63 ± 16	61 ± 14
В	0	49 ± 10	53 ± 11	61 ± 14	60 ± 14
UB	0.32 ± 0.07	0.82 ± 0.17	0.92 ± 0.15	1.07 ± 0.21	1.22 ± 0.26
В	0.27 ± 0.07	0.78 ± 0.17	0.95 ± 0.18	1.13 ± 0.22	$1.39 \pm 0.26^{\circ}$
VCO ₂ , l/min					
UB	0.27 ± 0.06	0.72 ± 0.16	0.85 ± 0.16	1.01 ± 0.18	1.29 ± 0.23
В	0.25 ± 0.08	0.68 ± 0.17	0.85 ± 0.15	1.06 ± 0.22	1.29 ± 0.27
Ϋ́E, l/min					
UB	9.3 ± 2.3	21.2 ± 4.5	25.6 ± 4.9	30.0 ± 5.9	38.4 ± 7.7
В	9.5 ± 3.5	20.8 ± 4.4	26.0 ± 4.7	32.2 ± 7.4	37.3 ± 10.3
$f_{\rm R}$, breaths/min					
UB	15.5 ± 3.3	28.2 ± 5.6	34.9 ± 7.0	37.4 ± 9.1	38.6 ± 8.2
В	14.0 ± 2.8	31.0 ± 6.9	35.9 ± 6.6	40.7 ± 7.2	40.2 ± 10.0
V _T , liters					
UB	0.61 ± 0.16	0.87 ± 0.19	0.85 ± 0.17	0.93 ± 0.18	1.01 ± 0.17
В	0.72 ± 0.32	0.84 ± 0.31	0.84 ± 0.24	0.90 ± 0.22	0.99 ± 0.21
T _I /T _{TOT}					
UB	0.45 ± 0.03	0.48 ± 0.05	0.45 ± 0.04	0.46 ± 0.05	0.48 ± 0.02
В	0.44 ± 0.06	0.45 ± 0.04	0.47 ± 0.04	0.51 ± 0.06	0.47 ± 0.03
V _T /T _i , l/s					
UB	0.31 ± 0.09	0.66 ± 0.15	0.88 ± 0.20	0.99 ± 0.30	1.23 ± 0.23
В	0.30 ± 0.09	0.73 ± 0.16	0.86 ± 0.18	0.94 ± 0.53	1.23 ± 0.32
SpO ₂ , %					
UB	97 ± 1	97 ± 2	98 ± 2	97 ± 3	96 ± 3
В	97 ± 1	97 ± 2	97 ± 2	96 ± 3	97 ± 3
Heart rate, beats/min					
UB	60 ± 9	83 ± 11	92 ± 9	102 ± 10	108 ± 10
В	58 ± 11	78 ± 11	88 ± 9	99 ± 8	104 ± 6
[La ⁻] _B , mmol/l‡					
UB	0.7 ± 0.2	0.6 ± 0.2	0.8 ± 0.3	1.4 ± 0.6	2.1 ± 1.2
В	0.7 ± 0.2	0.6 ± 0.1	0.6 ± 0.2	1.0 ± 0.3	$1.5 \pm 0.8*$
RPE (dyspnea)					
UB	0	1.1 ± 0.9	2.2 ± 0.8	3.3 ± 1.4	3.7 ± 0.8
В	0	1.3 ± 0.9	2.3 ± 0.9	3.3 ± 1.3	3.4 ± 1.3
RPE (arm discomfort)					
UB	0	1.4 ± 0.9	2.3 ± 0.6	4.1 ± 1.1	4.9 ± 1.9
B	0	14 ± 08	23 ± 0.6	37 ± 07	44 ± 10

Values are means \pm SD for 8 subjects.UB, unbound; B, bound; \dot{V}_{02} , O_2 uptake; \dot{V}_{C02} , CO_2 output; \dot{V}_E , minute ventilation; f_R , respiratory frequency; V_T , tidal volume; T_i/T_{tot} , inspiratory duty cycle; V_T/T_i , mean inspiratory flow; SpO₂, arterial O₂ saturation; [La⁻]_B, blood lactate concentration; RPE, ratings of perceived exertion. \pm Significant interaction effect (P < 0.05). *Significant post hoc pairwise comparison (P < 0.05).



Fig. 2. O₂ uptake/heart rate slopes in the bound (dashed line) and unbound condition (solid line) for measurements during each stage of the submaximal exercise test. Slopes were less steep in the bound condition (P < 0.05); see text for details. Data are means \pm SE for 8 subjects.

Respiratory Mechanics and Ventilatory Constraint

Pressure-derived measurements of respiratory mechanics and ventilatory constraint are reported for seven subjects, because one subject could not tolerate the balloon catheters. End-expiratory and end-inspiratory pressures during the submaximal exercise test are shown in Fig. 3. In the unbound condition, end-expiratory and end-inspiratory P_{di} increased sharply from baseline to the first stage of exercise. Endinspiratory P_{di} continued to increase throughout exercise, whereas end-expiratory P_{di} increased initially and leveled-off thereafter. Both pressures were significantly elevated with application of the binder, primarily because of an increase in the P_{ga} contribution.

Additional indices of respiratory mechanics and ventilatory constraint are summarized in Table 4. Dynamic inspiratory pressures (peak-to-peak and integrated) increased progres-

Table 3. Effect of abdominal binding on peakcardiopulmonary, metabolic, and perceptual responses

	Unbound	Bound	P value
Power output, W	49 ± 12	50 ± 13	0.980
Push rate, /min	61 ± 13	60 ± 13	0.918
VO ₂ , l/min	1.29 ± 0.33	1.43 ± 0.35	0.001*
VO ₂ , ml/kg/min	19.0 ± 2.1	21.2 ± 2.8	0.001*
VCO ₂ , l/min	1.38 ± 0.36	1.54 ± 0.35	0.155
RER	1.08 ± 0.12	1.08 ± 0.13	0.985
Ϋ́E, l/min	48.9 ± 14.1	46.1 ± 8.7	0.528
$f_{\rm R}$, breaths/min	54 ± 14	53 ± 15	0.838
V _T , 1	0.94 ± 0.21	0.92 ± 0.24	0.709
T _i /T _{tot}	0.48 ± 0.04	0.52 ± 0.06	0.074
V _T /T _i , l/s	1.70 ± 0.67	1.70 ± 0.70	0.978
$\dot{V}_{E}/\dot{V}O_{2}$	39.0 ± 10.2	33.0 ± 6.0	0.067
$\dot{V}_{E}/\dot{V}_{CO_{2}}$	35.6 ± 6.1	30.6 ± 4.4	0.012*
P _{ET} CO ₂ , mmHg	35.5 ± 5.8	37.5 ± 8.0	0.232
SpO ₂ , %	95 ± 3	95 ± 3	0.949
Heart rate, beats/min	120 ± 12	122 ± 13	0.534
[.] VO₂/heart rate, ml/beat	10.7 ± 3.1	12.4 ± 3.2	0.001*
[La ⁻] _B , mmol/l	4.6 ± 1.2	3.8 ± 1.0	0.052
RPE (dyspnea)	7.0 ± 2.7	7.1 ± 2.9	0.917
RPE (arm discomfort)	7.5 ± 2.0	7.4 ± 2.0	0.919

Values are means \pm SD for 8 subjects. *Significant difference between conditions (P < 0.05).



Fig. 3. End-expiratory (squares) and end-inspiratory (circles) transdiaphragmatic pressure (*A*), gastric pressure (*B*), and esophageal pressure (*C*) at rest and during submaximal wheelchair propulsion in the bound (dashed lines) and unbound condition (solid lines). Note that end-expiratory and end-inspiratory transdiaphragmatic and gastric pressures were elevated throughout exercise in the bound condition. Data are means \pm SE for 7 subjects. \dagger Significant main effect for condition (P < 0.05). *Significant post hoc pairwise comparison (P < 0.05).

sively throughout exercise, but were not different across conditions. Dynamic lung compliance fell from baseline to the first stage of exercise then remained stable through to the final stage. Dynamic lung compliance was slightly higher in the bound condition during the latter stages of exercise but did not reach statistical significance. In the unbound condition, TTI_{di} increased progressively throughout exercise due almost entirely to the aforementioned increase in tidal transdiaphragmatic pressure. There was no effect of binding on breath timing, but a slight increase in the maximum pressure-generating capacity of the diaphragm (unbound 125 ± 49 vs. bound $138 \pm 32 \text{ cmH}_2\text{O}$, P = 0.207) resulted in a trend toward a binding-induced reduction in TTI_{di} (0.20 vs. 0.16 for final stage).

Operating lung volumes at rest and during exercise are shown in Fig. 4. In the unbound condition, there was a sharp rise in EELV and EILV from rest to the first stage of exercise and a more gradual increase through to the final stage. Both volumes were shifted to a lower percentage of total lung capacity in the bound condition $(-7 \pm 2\%)$ for EELV, P = 0.017; $-8 \pm 2\%$ for EILV, P = 0.035), and the rates of rise

Effect	Baseline	Stage 1	Stage 2	Stage 3	Stage 4
P _{di.tidal} , cmH ₂ O					
UB	16.0 ± 6.9	29.9 ± 12.8	33.5 ± 11.5	40.9 ± 12.8	44.4 ± 10.1
В	20.5 ± 6.0	35.3 ± 16.7	39.0 ± 17.3	39.7 ± 14.2	43.9 ± 14.4
$P_{ga,tidal}, cmH_2O$					
UB	13.3 ± 6.6	22.7 ± 10.8	25.4 ± 10.0	31.9 ± 10.6	35.5 ± 7.6
В	16.7 ± 5.6	27.5 ± 14.5	29.5 ± 13.7	31.5 ± 11.5	35.4 ± 11.8
Pes,tidal, cmH ₂ O*					
UB	-2.7 ± 0.7	-7.2 ± 2.5	-8.1 ± 2.6	-8.9 ± 3.0	-10.4 ± 4.3
В	-3.8 ± 1.8	-7.9 ± 3.1	-9.5 ± 5.4	-8.9 ± 4.0	-9.9 ± 5.3
PTP _{di} , cmH ₂ O·s/min					
UB	225 ± 123	347 ± 148	419 ± 137	467 ± 224	461 ± 231
В	287 ± 137	420 ± 244	470 ± 242	515 ± 136	514 ± 204
PTP ₅₀ , cmH ₂ O·s/min					
UB	187 ± 123	249 ± 132	304 ± 132	328 ± 183	304 ± 211
В	232 ± 131	314 ± 189	316 ± 165	375 ± 109	454 ± 127
PTP _{es} , cmH ₂ O·s/min					
UB	-38 ± 12	-98 ± 53	-115 ± 48	-135 ± 65	-157 ± 44
В	-55 ± 24	-106 ± 79	-135 ± 65	-140 ± 72	-161 ± 86
C _{L,dyn} , ml/cmH ₂ O					
UB	184 ± 47	118 ± 58	98 ± 46	103 ± 51	98 ± 54
В	162 ± 45	113 ± 57	114 ± 51	117 ± 79	123 ± 79
TTI _{di}					
UB	0.070 ± 0.029	0.109 ± 0.044	0.137 ± 0.063	0.147 ± 0.059	0.203 ± 0.115
В	0.071 ± 0.029	0.074 ± 0.037	0.111 ± 0.043	0.128 ± 0.055	0.159 ± 0.085
IRV/TLC, %					
UB	39 ± 9	27 ± 3	28 ± 5	21 ± 5	17 ± 10
В	43 ± 13	33 ± 9	31 ± 7	29 ± 6	26 ± 4
IFR, % capacity					
UB	6 ± 2	20 ± 13	27 ± 15	26 ± 11	34 ± 18
В	10 ± 2	20 ± 7	28 ± 15	29 ± 11	34 ± 14
$\dot{V}_{E}/\dot{V}_{ECAP}, \%$					
UB	10 ± 6	17 ± 5	26 ± 9	24 ± 7	40 ± 26
В	15 ± 10	18 ± 9	28 ± 16	23 ± 8	31 ± 16

Table 4. Effect of abdominal binding on respiratory mechanics and ventilatory constraint at rest and during submaximal incremental wheelchair propulsion

Values are means \pm SD for 7 subjects. UB, unbound; B, bound; P_{di,tidal}, inspiratory tidal transdiaphragmatic pressure; P_{ga,tidal}, inspiratory tidal gastric pressure; P_{es,tidal}, inspiratory tidal esophageal pressure; PTP_{di}, diaphragm pressure-time product; PTP_{ga}, gastric pressure-time product; PTP_{es}, esophageal pressure-time product; C_{L,dyn}, dynamic lung compliance; TTI_{di}, inspiratory diaphragm tension-time index; IRV/TLC, index of change in end-inspiratory lung volume; IFR, inspiratory flow reserve; V_E/V_{ECAP}, ventilatory capacity calculated from a theoretical maximal exercise ventilation based on the maximal available expiratory airflow over the range of the tidal breath placed at the measured end-expiratory lung volume. *Significant main effect for condition (P < 0.05).

were reduced. During the final stage in the unbound condition, EILV averaged 83% of total lung capacity with three subjects exceeding 90%. With binding, EILV was reduced to less than



Fig. 4. End-expiratory (squares) and end-inspiratory (circles) lung volume at rest and in response to submaximal wheelchair propulsion in the bound (dashed lines, closed symbols) and unbound condition (solid lines, open symbols). Note the immediate and progressive increase from resting values in operating lung volumes (i.e., dynamic hyperinflation) and the downward shift in lung volumes in response to abdominal binding. Data are means \pm SE for 7 subjects. \pm Significant main effect for condition (P < 0.05). \pm Significant interaction effect (P < 0.05). \pm Significant post hoc pairwise comparison (P < 0.05).

80% of total lung capacity in all subjects. There was no encroachment of the tidal flow-volume curves on the maximum flow-volume envelope in any subject (e.g., Fig. 5). Furthermore, there was substantial reserve for increasing flow and volume as indicated by the low values for IFR and \dot{V}_E/\dot{V}_{ECAP} , respectively (Table 4).

DISCUSSION

This study investigated the influence of abdominal binding on respiratory mechanics during wheelchair exercise in highly trained athletes with cervical SCI. The main finding was that binding induced passive increases in intra-abdominal pressure that resulted in a shift of tidal breathing to lower lung volumes with no effect on expiratory flow limitation, symptoms, or exercise tolerance. The binding-induced changes in intra-abdominal pressure were accompanied by increases in whole body O₂ uptake and decreases in systemic blood lactate at high relative intensities of exercise ($\geq 95\%$ peak O₂ uptake). These latter findings suggest that abdominal binding influences the overall exercise response by an increase in O₂ transport capacity.

To our knowledge, this is the first report of respiratory mechanics during wheelchair exercise in individuals with SCI



Fig. 5. Maximal and tidal flow-volume curves at rest and during the submaximal exercise test for a single subject in the unbound and bound condition. Each tidal flow-volume curve is ensemble-averaged over 30 s of resting baseline (R) and over the first 30 s of the final minute of each exercise stage (1-4). Note the leftward shift of the tidal flow-volume curves as exercise progresses, the rightward shift of the tidal flow-volume curves and concomitant increases in inspiratory reserve volume with binding, and the increase in vital capacity and maximal midexpiratory flows. Vertical dotted lines indicate the binding-induced changes in total lung capacity (*left*) and residual volume (*right*).

and the first to assess the effect of abdominal binding on exercise responses in cervical SCI. A novel finding was the sudden and sustained rise in end-expiratory lung volume (i.e., dynamic hyperinflation), despite no evidence of expiratory flow limitation. This finding is consistent with our previous observation for cervical SCI during arm-crank ergometry (41), but in contrast to that reported for able-bodied subjects during lower-body exercise whereby end-expiratory lung volume only increases above relaxation volume when subjects approach their mechanical limits to generate expiratory flow (2). It is not entirely clear whether the rise in end-expiratory lung volume is a consequence of expiratory muscle weakness (40) or merely the "normal" response to upper-body exercise (11). The expiratory muscle paralysis that accompanies cervical SCI leads to an increased recruitment of nontypical accessory muscles of expiration (e.g., pectoralis major) to expire below functional residual capacity (13). However, many of these accessory muscles are also involved as prime movers during wheelchair propulsion (28, 46). It is perhaps, therefore, unsurprising that hyperinflation prevails from the onset of exercise. The increased elastic recoil characteristics of the lung and chest wall at high lung volumes may be a mechanism by which individuals with cervical SCI are able to "passively" increase expiratory flow during exercise. Although abdominal binding did not prevent dynamic hyperinflation, it did cause a parallel downward shift in operating lung volumes at rest and throughout exercise. An increase in elastic recoil pressure with binding might be expected to increase driving pressure for tidal expiratory flow (9), albeit at a lower operating lung volume. Importantly, the downward shift in operating lung volumes did not appear to impose mechanical ventilatory constraints (see Fig. 5). Indeed, ventilatory reserve as a fraction of capacity was similar in both conditions, presumably owing to the bindinginduced increases in vital capacity and maximal midexpiratory flows.

We reason that abdominal binding improves the ventilatory response to exercise through several interrelated factors. First, the binding-induced increase in intra-abdominal pressure during inspiration would be expected to increase expansion of the lower rib cage (31, 44). This, in turn, may improve overall gas exchange consequent to an increase in the ventilation-perfusion ratio of lung units (55). Indeed, the ventilatory equivalent for CO_2 at peak exercise was reduced with binding (Table 3). Moreover, the physiological dead space ventilation estimated using the alveolar ventilation equation and an assumed anatomic dead space of 150 ml was more than halved (3.6 bound vs. 7.5 l/min unbound). Although the presumed increase in lower rib cage expansion has been attributed to an increase in appositional forces (31, 44), more recent evidence suggests that binding may enable the diaphragm to operate on a more effective portion of its length-tension relationship and thereby exert greater insertional force (57). This increase in mechanical advantage might be expected to decrease the propensity for diaphragm fatigue. In the unbound condition, the product of $\overline{P}_{di}/P_{di,max}$ and T_I/T_{TOT} (TTI_{di}) during the final stage of submaximal exercise (0.20) exceeded "critical" values that have been proposed to elicit diaphragm fatigue in healthy, nondisabled individuals (>0.15) (7) and individuals with cervical SCI (>0.10) (33). With binding, however, there was a reduction in TTI_{di} (0.16) consequent to a slight increase in the capacity of the diaphragm to generate inspiratory pressure (P_{di,max}). Although we acknowledge that the critical TTI_{di} concept may not apply directly to the hyperpnea of exercise (41), other factors known to influence energy demands, namely respiratory frequency and velocity of diaphragm shortening (V_T/T_I) , were unaffected by binding. Thus the potential benefits of binding may revolve around an increase in the capacity and/or efficiency of the inspiratory muscles, which, in turn, would be expected to improve the overall energetics of these muscles. Despite the aforementioned changes in respiratory mechanics, dyspnea intensity ratings were essentially the same at any given power output and ventilation in both conditions. This latter finding suggests that binding-induced alterations in respiratory mechanics do not contribute importantly to exertional dyspnea in highly fit individuals with cervical SCI.

The changes in respiratory mechanics with binding were accompanied by significant changes in O_2 uptake (8-12%) and blood lactate concentration (-16-19%) at high relative power outputs. By using a similar exercise protocol and subject population, Leicht et al. (27) reported within-day coefficients of variation of <6% for peak O_2 uptake and <14% for peak lactate concentration. Thus the relatively large changes noted in the current study were likely to be "true" differences. The findings are an extension of our recent field-based study in which the distance covered during a 4-min maximal push test was significantly increased with binding and the blood lactate response was significantly reduced (50). In the only other study

to investigate the influence of abdominal binding in athletes with SCI, Kerk et al. (25) found no change in O₂ uptake during submaximal or maximal wheelchair exercise. The discrepancy may be because Kerk et al. (25) set the degree of abdominal compression based on a change in abdominal girth, whereas we adjusted the binder so that end-expiratory gastric pressure reached a level known to optimize resting cardiopulmonary function (52). Furthermore, Kerk et al. (25) studied athletes with high-thoracic SCI (\geq T₆), who, because of partial or full descending sympathetic control of the myocardium and upperbody vasculature, would be less likely to exhibit cardiovascular limitation during exercise and therefore benefit from binding.

The reason for the binding-induced increase in O₂ uptake is not entirely clear. Power outputs were matched and push rates were similar across conditions. Moreover, we have shown that propulsion kinematics are not significantly altered with binding (50). It seems unlikely, therefore, that the greater increase in O_2 uptake could be accounted for by an increase in the amount of active musculature and/or a decrease in mechanical efficiency. A potential explanation relates to an increase in work (and O_2) cost) of breathing, as suggested by the slightly elevated tidal swings in transdiaphragmatic pressure with binding. In healthy nondisabled subjects, who would be expected to achieve much higher levels of ventilation than individuals with cervical SCI, the O_2 cost of breathing during maximal whole body exercise averages 8-10% of total O₂ uptake (1). Thus, although an increase in respiratory muscle work might have accounted for a small proportion of the increase in total O₂ uptake with binding, we doubt whether this could have contributed a significant amount to the 12% increase at peak exercise.

A more likely explanation for the binding-induced increase in O2 uptake relates to an improvement in central hemodynamics. Although our study was not specifically designed to address this issue, our observations do merit discussion. The increases in abdominal pressure due to application of the binder (Fig. 3) may be expected to decrease vascular compliance, increase mean vascular pressure, and therefore increase stroke volume. The increase in end-expiratory and end-inspiratory abdominal pressures might also be expected to increase the degree of driving pressure for venous return during tidal breathing. In this regard, Aliverti et al. (3, 4) showed that the circulatory function of the diaphragm in nondisabled subjects is greatly enhanced by the action of the abdominal muscles. Increases in abdominal pressure with quiet diaphragmatic breathing were shown to expel blood from the splanchnic vascular bed (3, 4). Moreover, increases in abdominal pressure resulting from expulsive maneuvers performed by simultaneous contractions of the diaphragm and abdominal muscles were shown to augment the circulatory function of the diaphragm (3, 4). These findings are relevant in so far as individuals with cervical SCI lack central sympathetic control (42). As a result, blood pooling occurs in nonactive vascular beds, including the splanchnic region (43). This, in turn, may limit O₂ transport capacity by restricting the ability to increase venous return and stroke volume (22). In the current study, the increase in heart rate for a given increase in O_2 uptake was reduced by ${\sim}20\%$ with binding (Fig. 2) and the O_2 pulse at high relative exercise intensities was increased by $\sim 16\%$. These latter findings are consistent with our observation of an improvement in left-ventricular function at rest (52) and are highly suggestive of a binding-induced increase in stroke volume during exercise (56).

Another potential mechanism for the proposed increase in stroke volume with binding relates to the downward shift in operating lung volumes. In the unbound condition, end-inspiratory lung volume averaged 83% of total lung capacity, and three subjects achieved >90% (see Fig. 5). Conceivably, this severe level of dynamic hyperinflation may place a constraint on ventricular preload during inspiration by a compressive effect of the lung on the cardiac fossa and the inferior and superior vena cava (29, 39). In turn, the decrease in endinspiratory lung volume with binding may have reduced mechanical compression of the heart and great vessels, thereby resulting in an elevation of cardiac filling and stroke volume. An effect of changing operating lung volumes on cardiac function might be particularly relevant for individuals with cervical SCI because lung compliance is reduced in this population (37). Thus binding may exert a cardiogenic benefit, both directly via an abdominothoracic translocation of blood and indirectly via an attenuation of dynamic hyperinflation. The consequent increase in blood flow to working muscles may explain the modest but consistent reductions in blood lactate concentration at high exercise intensities. This effect of increasing blood flow may be attributed to alterations in metabolism resulting from increases in O₂ delivery and metabolite removal (6).

Despite a greater peak O₂ uptake with binding, peak power output was similar across conditions. This appears to suggest that exercise tolerance was limited more by the ability of the muscles to use O₂ (i.e., peripheral factors) than the capacity to transport O_2 (i.e., central factors). Alternatively, the exercise protocol (i.e., rapid increases in gradient with a constant speed) may have been suboptimal for eliciting a true peak response, therefore masking our ability to detect a binding-induced increase in exercise tolerance. We recently showed that peak heart rate is significantly higher during a field-based endurance test compared with a laboratory-based incremental treadmill test (54). Moreover, when the subjects in the current study were tested using the field-based test, every subject demonstrated a binderinduced improvement in endurance performance (50). Further support for our postulate that rapid increases in gradient may not be suitable for detecting changes in exercise tolerance stems from the finding that elite hand-cyclists with cervical SCI perform worse against their counterparts with thoracic SCI during uphill pushing vs. on the flat (49).

In conclusion, abdominal binding shifts tidal breathing to lower lung volumes with no effect on flow limitation, symptom intensities, or exercise tolerance. Changes in respiratory mechanics with binding may raise muscle blood flow and O_2 delivery during maximal exercise by an increase in cardiac filling and output. Potential mechanisms include a translocation of blood from the abdomen to the heart and a decrease in mechanical compression of the heart and great vessels via a shift of tidal breathing to lower lung volumes. The physiological relevance of the findings is that O₂ transport capacity in cervical SCI may be limited by an inability of the cardiovascular system to further increase cardiac output. From a practical perspective, binder-induced improvements in central circulatory function may enable individuals with cervical SCI to achieve greater cardiovascular adaptations to exercise training. Future studies should include direct measurements of central and peripheral hemodynamics to fully characterize the acute

and chronic effects of abdominal binding on O_2 delivery and utilization during exercise.

ACKNOWLEDGMENTS

We acknowledge the support provided by Great Britain Wheelchair Rugby and ParalympicsGB as well as the technical assistance of Christof Leicht and John Lenton. We are also grateful for the advice and critique provided by William Sheel and Paolo Dominelli.

Present addresses: C. West, ICORD, 818 West 10th Ave, Vancouver, BC, V5Z 1M9, Canada; I. Campbell, University of Hertfordshire, Hatfield, Hertfordshire, AL10 9AB, UK.

GRANTS

The study was funded by UK Sport through the Ideas4Innovation Programme.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

AUTHOR CONTRIBUTIONS

Author contributions: C.R.W., V.L.G.-T., I.G.C., and L.M.R. conception and design of research; C.R.W. performed experiments; C.R.W. and L.M.R. analyzed data; C.R.W. and L.M.R. interpreted results of experiments; C.R.W. prepared figures; C.R.W. and L.M.R. drafted manuscript; C.R.W., V.L.G.-T., I.G.C., and L.M.R. edited and revised manuscript; C.R.W., V.L.G.-T., I.G.C., and L.M.R. approved final version of manuscript.

REFERENCES

- Aaron EA, Seow KC, Johnson BD, Dempsey JA. Oxygen cost of exercise hyperpnea: implications for performance. J Appl Physiol 72: 1818–1825, 1992.
- Alison JA, Regnis JA, Donnelly PM, Adams RD, Sullivan CE, Bye PT. End-expiratory lung volume during arm and leg exercise in normal subjects and patients with cystic fibrosis. *Am J Respir Crit Care Med* 158: 1450–1458, 1998.
- Aliverti A, Bovio D, Fullin I, Dellaca RL, Lo Mauro A, Pedotti A, Macklem PT. The abdominal circulatory pump. *PLoS One* 4: e5550, 2009.
- Aliverti A, Uva B, Laviola M, Bovio D, Lo Mauro A, Tarperi C, Colombo E, Loomas B, Pedotti A, Similowski T, Macklem PT. Concomitant ventilatory and circulatory functions of the diaphragm and abdominal muscles. J Appl Physiol 109: 1432–1440, 2010.
- ATS/ACCP. ATS/ACCP statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med* 167: 211–277, 2003.
- Barclay JK. A delivery-independent blood flow effect on skeletal muscle fatigue. J Appl Physiol 61: 1084–1090, 1986.
- Bellemare F, Grassino A. Effect of pressure and timing of contraction on human diaphragm fatigue. J Appl Physiol 53: 1190–1195, 1982.
- 8. **Borg G.** *Borg's Perceived Exertion and Pain Scales.* Champaign, IL: Human Kinetics, 1998.
- Bradley CA, Anthonisen NR. Rib cage and abdominal restrictions have different effects on lung mechanics. J Appl Physiol 49: 946–952, 1980.
- Bruschi C, Cerveri I, Zoia MC, Fanfulla F, Fiorentini M, Casali L, Grassi M, Grassi C. Reference values of maximal respiratory mouth pressures: a population-based study. *Am Rev Respir Dis* 146: 790–793, 1992.
- Cerny FJ, Ucer C. Arm work interferes with normal ventilation. Appl Ergon 35: 411–415, 2004.
- Cunningham DJ, Guttmann L, Whitteridge D, Wyndham CH. Cardiovascular responses to bladder distension in paraplegic patients. J Physiol 121: 581–592, 1953.
- De Troyer A, Estenne M, Heilporn A. Mechanism of active expiration in tetraplegic subjects. N Engl J Med 314: 740–744, 1986.
- Figoni SF, Rodgers MM, Glaser RM, Hooker SP, Feghri PD, Ezenwa BN, Mathews T, Suryaprasad AG, Gupta SC. Physiologic responses of paraplegics and quadriplegics to passive and active leg cycle ergometry. J Am Paraplegia Soc 13: 33–39, 1990.
- Green M, Road J, Sieck GC, Similowski T. Tests of respiratory muscle strength. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med* 166: 518–624, 2002.
- Grimby G, Sóderholm B. Spirometric studies in normal subjects. Acta Med Scand 173: 199–206, 1963.

- Hooker SP, Figoni SF, Rodgers MM, Glaser RM, Mathews T, Suryaprasad AG, Gupta SC. Physiologic effects of electrical stimulation leg cycle exercise training in spinal cord injured persons. *Arch Phys Med Rehabil* 73: 470–476, 1992.
- Hooker SP, Greenwood JD, Boyd LA, Hodges MR, McCune LD, McKenna GE. Influence of posture on arm exercise tolerance and physiologic responses in persons with spinal cord injured paraplegia. *Eur J Appl Physiol* 67: 563–566, 1993.
- Hooker SP, Greenwood JD, Hatae DT, Husson RP, Matthiesen TL, Waters AR. Oxygen uptake and heart rate relationship in persons with spinal cord injury. *Med Sci Sports Exerc* 25: 1115–1119, 1993.
- Hopman MT, Dueck C, Monroe M, Philips WT, Skinner JS. Limits to maximal performance in individuals with spinal cord injury. *Int J Sports Med* 19: 98–103, 1998.
- Hopman MT, Monroe M, Dueck C, Phillips WT, Skinner JS. Blood redistribution and circulatory responses to submaximal arm exercise in persons with spinal cord injury. *Scand J Rehabil Med* 30: 167–174, 1998.
- Hostettler S, Leuthold L, Brechbuhl J, Mueller G, Illi SK, Spengler CM. Maximal cardiac output during arm exercise in the sitting position after cervical spinal cord injury. *J Rehabil Med* 44: 131–136, 2012.
- Johnson BD, Weisman IM, Zeballos RJ, Beck KC. Emerging concepts in the evaluation of ventilatory limitation during exercise: the exercise tidal flow-volume loop. *Chest* 116: 488–503, 1999.
- Kelley A, Garshick E, Gross ER, Lieberman SL, Tun CG, Brown R. Spirometry testing standards in spinal cord injury. *Chest* 123: 725–730, 2003.
- Kerk JK, Clifford PS, Snyder AC, Prieto TE, O'Hagan KP, Schot PK, Myklebust JB, Myklebust BM. Effect of an abdominal binder during wheelchair exercise. *Med Sci Sports Exerc* 27: 913–919, 1995.
- Krassioukov A. Autonomic function following cervical spinal cord injury. *Respir Physiol Neurobiol* 169: 157–164, 2009.
- Leicht CA, Tolfrey K, Lenton JP, Bishop NC, Goosey-Tolfrey VL. The verification phase and reliability of physiological parameters in peak testing of elite wheelchair athletes. *Eur J Appl Physiol* 113: 337–345, 2013.
- Lin HT, Su FC, Wu HW, An KN. Muscle forces analysis in the shoulder mechanism during wheelchair propulsion. *Proc Inst Mech Eng H* 218: 213–221, 2004.
- 29. Marini JJ, Culver BH, Butler J. Mechanical effect of lung distention with positive pressure on cardiac function. *Am Rev Respir Dis* 124: 382–386, 1981.
- Mateus SR, Beraldo PS, Horan TA. Maximal static mouth respiratory pressure in spinal cord injured patients: correlation with motor level. *Spinal Cord* 45: 569–575, 2007.
- Mead J, Banzett RB, Lehr J, Loring SH, O'Cain CF. Effect of posture on upper and lower rib cage motion and tidal volume during diaphragm pacing. *Am Rev Respir Dis* 130: 320–321, 1984.
- 32. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CP, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J. Standardisation of spirometry. *Eur Respir J* 26: 319–338, 2005.
- Nava S, Rubini F, Zanotti E, Caldiroli D. The tension-time index of the diaphragm revisited in quadriplegic patients with diaphragm pacing. *Am J Respir Crit Care Med* 153: 1322–1327, 1996.
- Pitetti KH, Barrett PJ, Campbell KD, Malzahn DE. The effect of lower body positive pressure on the exercise capacity of individuals with spinal cord injury. *Med Sci Sports Exerc* 26: 463–468, 1994.
- 35. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. *Eur Respir J Suppl* 16: 5–40, 1993.
- Rodarte JR, Rehder K. Dynamics of respiration. In: Handbook of Physiology. The Respiratory System. Mechanics of Breathing. Bethesda, MD: Am Physiol Soc, 1986, p. 131–144.
- 37. Scanlon PD, Loring SH, Pichurko BM, McCool FD, Slutsky AS, Sarkarati M, Brown R. Respiratory mechanics in acute quadriplegia. Lung and chest wall compliance and dimensional changes during respiratory maneuvers. Am Rev Respir Dis 139: 615–620, 1989.
- Schmid A, Huonker M, Barturen JM, Stahl F, Schmidt-Trucksass A, Konig D, Grathwohl D, Lehmann M, Keul J. Catecholamines, heart rate, and oxygen uptake during exercise in persons with spinal cord injury. *J Appl Physiol* 85: 635–641, 1998.
- Takata M, Robotham JL. Ventricular external constraint by the lung and pericardium during positive end-expiratory pressure. *Am Rev Respir Dis* 143: 872–875, 1991.

- Taylor BJ, How SC, Romer LM. Expiratory muscle fatigue does not regulate operating lung volumes during high-intensity exercise in healthy humans. J Appl Physiol 114: 1569–1576, 2013.
- Taylor BJ, West CR, Romer LM. No effect of arm-crank exercise on diaphragmatic fatigue or ventilatory constraint in Paralympic athletes with cervical spinal cord injury. J Appl Physiol 109: 358–366, 2010.
- Theisen D. Cardiovascular determinants of exercise capacity in the Paralympic athlete with spinal cord injury. *Exp Physiol* 97: 319–324, 2012.
- Thijssen DH, Steendijk S, Hopman MT. Blood redistribution during exercise in subjects with spinal cord injury and controls. *Med Sci Sports Exerc* 41: 1249–1254, 2009.
- 44. Urmey W, Loring S, Mead J, Slutsky AS, Sarkarati M, Rossier A, Brown R. Upper and lower rib cage deformation during breathing in quadriplegics. *J Appl Physiol* 60: 618–622, 1986.
- 45. van der Woude LHV, Veeger HEJ, Dallmeijer AJ, Janssen TWJ, Rozendaal LA. Biomechanics and physiology in active manual wheelchair propulsion. *Med Eng Phys* 23: 713–733, 2001.
- Vanlandewijck Y, Theisen D, Daly D. Wheelchair propulsion biomechanics: implications for wheelchair sports. *Sports Med* 31: 339–367, 2001.
- Wadsworth BM, Haines TP, Cornwell PL, Paratz JD. Abdominal binder use in people with spinal cord injuries: a systematic review and meta-analysis. *Spinal Cord* 47: 274–285, 2009.
- Wanger J, Clausen JL, Coates A, Pedersen OF, Brusasco V, Burgos F, Casaburi R, Crapo R, Enright P, van der Grinten CP, Gustafsson P, Hankinson J, Jensen R, Johnson D, Macintyre N, McKay R, Miller MR, Navajas D, Pellegrino R, Viegi G. Standardisation of the measurement of lung volumes. *Eur Respir J* 26: 511–522, 2005.

- 49. Weissland T, Lepretre PM. Are tetraplegic handbikers going to disappear from team relay in para-cycling? *Fron Physiol* 4: 77, 2013.
- West CR, Campbell IG, Goosey-Tolfrey VL, Mason BS, Romer LM. Effects of abdominal binding on field-based exercise responses in Paralympic athletes with cervical spinal cord injury. *J Sci Med Sport*. First published July 20, 2013; doi:10.1016/j.jsams.2013.06.001.
- West CR, Campbell IG, Romer LM. Assessment of pulmonary restriction in cervical spinal cord injury: a preliminary report. *Arch Phys Med Rehabil* 93: 1463–1465, 2012.
- West CR, Campbell IG, Shave RE, Romer LM. Effects of abdominal binding on cardiorespiratory function in cervical spinal cord injury. *Respir Physiol Neurobiol* 180: 275–282, 2012.
- West CR, Campbell IG, Shave RE, Romer LM. Resting cardiopulmonary function in Paralympic athletes with cervical spinal cord injury. *Med Sci Sports Exerc* 44: 323–329, 2012.
- West CR, Romer LM, Krassioukov A. Autonomic function and exercise performance in elite athletes with cervical spinal cord injury. *Med Sci Sports Exerc* 45: 261–267, 2013.
- West JB. Ventilation-perfusion relationship. In: *Respiratory Physiology: The Essentials*, edited by Duffy N. Baltimore, MD: Lippincott Williams & Wilkins, 2012, p. 56–76.
- Whipp BJ, Higgenbotham MB, Cobb FC. Estimating exercise stroke volume from asymptotic oxygen pulse in humans. J Appl Physiol 81: 2674–2679, 1996.
- Wilson TA, De Troyer A. Effects of the insertional and appositional forces of the canine diaphragm on the lower ribs. *J Physiol* 591: 3539–3548, 2013.

