Loughborough University

This item was submitted to Loughborough's Research Repository by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Endurance performance is influenced by perceptions of pain and temperature: theory, applications and safety considerations

PLEASE CITE THE PUBLISHED VERSION

https://doi.org/10.1007/s40279-017-0852-6

PUBLISHER

Springer International Publishing

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This is a post-peer-review, pre-copyedit version of an article published in Sports Medicine. The final authenticated version is available online at: https://doi.org/10.1007/s40279-017-0852-6.

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Stevens, Christopher John, Alexis R. Mauger, Peter Hassmèn, and Lee Taylor. 2017. “Endurance Performance Is Influenced by Perceptions of Pain and Temperature: Theory, Applications and Safety Considerations”. figshare. https://hdl.handle.net/2134/11617731.v1.

This is article is published in Sports Medicine, a full copy can be accessed here –
https://link.springer.com/journal/volumesAndIssues/40279

Title: Endurance performance is influenced by perceptions of pain and temperature: Theory, applications and safety considerations.

Running Head: Perceptions of pain and temperature influence performance

Authors: Christopher J. Stevens[1*#], Alexis R. Mauger[2], Peter Hassmèn[1] & Lee Taylor[3,4*#]

*Christopher Stevens and Lee Taylor contributed equally to this manuscript

Affiliations:

[1]School of Health and Human Sciences, Southern Cross University, Coffs Harbour, NSW, Australia

[2]Endurance Research Group, School of Sport and Exercise Sciences, University of Kent, Chatham Maritime, Kent, United Kingdom

[3]ASPETAR, Qatar Orthopaedic and Sports Medicine Hospital, Athlete Health and Performance Research Centre, Doha, Qatar

[4]Loughborough University, School of Sport, Exercise and Health Sciences, Loughborough, UK

[#]**Addresses for Correspondence:**
Dr Christopher John Stevens
School of Health and Human Sciences
Southern Cross University
Hogbin Dr, Coffs Harbour, 2450, NSW, Australia
Email: Christopher.Stevens@scu.edu.au
Ph: +(61) 411 797 245

Dr Lee Taylor
P.O. Box 29222
ASPETAR Orthopaedic and Sports Medicine Hospital
Athlete Health and Performance Research Centre
Aspire Zone, Doha, Qatar, P.O. BOX 29222
Email: Lee.Taylor@aspetar.com
Ph: +(974) 4413 2403

Abstract

Models of endurance performance now recognise input from the brain, including an athlete's ability to cope with various non-pleasurable perceptions during exercise, such as pain and temperature. Exercise training can reduce perceptions of both pain and temperature over time, partly explaining why athletes generally have a higher pain tolerance, despite a similar pain threshold, compared to active controls. Several strategies with varying efficacy may ameliorate the perceptions of pain (e.g. acetaminophen, transcranial direct current stimulation and transcutaneous electrical stimulation) and temperature (e.g. menthol beverages, topical menthol products and other cooling strategies, especially those targeting the head) during exercise to improve athletic performance. This review describes both the theory and practical applications of these interventions in the endurance sport setting, as well as the potentially harmful health consequences of their use.

Key Points

*Athletes generally have a higher pain tolerance, despite a similar pain threshold, compared to active controls.

*Acetaminophen ingestion and transcutaneous electrical stimulation have been demonstrated to significantly decrease pain and improve endurance performance/capacity, but no data exist in elite populations.

*Endurance performance can be influenced by thermal perceptions independently of any changes in physical body temperature.

*Cooling strategies that induce large perceptual cooling effects and have practical application in endurance competition are ergogenic for athletes competing in the heat.

*Practitioners and athletes should deliberate on the ethical and safety considerations of strategies that override homeostatic controls before implementing them in competition.

1.0 Introduction

The mechanisms underpinning endurance capacity and performance have largely been defined in the exercise sciences with physiological concepts: a high maximal oxygen uptake, lactate threshold and exercise economy [1]. Recently, models of endurance performance have attributed greater value to the brain within exercise regulation [2-5], and specifically how athletes interpret what they 'feel' during exercise (hereafter referred to as 'perceptions'). Some describe the rating of perceived exertion (RPE) as the ultimate limiting factor to exercise tolerance [2], as it may be responsible for pacing, fatigue and exhaustion [3]. However, precisely what the RPE measures has been debated [6, 7]. Regardless, RPE likely incorporates aspects of effort, exertion and afferent sensory feedback, including the perceptions of pain and temperature [6].

Perceptions of pain and temperature during exercise have recently received increased research attention. They appear to be attenuated with normal exercise training [8, 9], and strategies manipulating these perceptions to improve endurance performance have also been investigated [10, 11]. While such research may drive advancements in knowledge around the relationship between psychology and physiology for endurance athletes, some aspects of the topic have been at the centre of intense international debate about the ethical and health considerations of manipulating the senses to override homeostatic controls during exercise. Therefore, the purpose of the current review is to summarise recent advancements in research surrounding the influences of pain and

temperature perceptions during exercise, while also reiterating the message of athlete safety in relation to their applications.

2.0 Literature Search Methods

Searching was carried out within a range of databases including Web of Science, Scopus and MEDLINE, up to 9th October, 2017. Searching was conducted in English and included the terms: perception, sensation, pain, temperature, heat, exercise, endurance and performance. Various combinations of search clusters were used, and search results were limited to peer-reviewed content. Additional works from key authors were also included in the study. Inclusion criteria stipulated that investigations must have been peer reviewed and measured perceptions of pain and/or temperature. This review was focussed on endurance exercise, which reflects the majority of the literature available. Participants of all abilities were included within both the sport and occupational health settings.

3.0 Perceptions of Pain and Endurance Performance

3.1 Theory

Pain is defined by the International Association for the Study of Pain as "an unpleasant sensory and emotional experience that is associated with actual or potential tissue damage or described in such terms" [12]. The meaning and implications of pain are learnt in the early years of life, where the sensation is related to injury or tissue damage [13]. However, pain is always subjective and the perception of the sensation is not always proportional to the size of the nociceptive signal and given pain is always unpleasant, the perception of pain will also partly be an emotional experience [13].

These sensory and subjective principles are critical for understanding the role of pain in exercise and athletic performance [14].

During exercise, pain arises as a natural consequence of intramuscular pressure, muscle distortion and a build-up of deleterious metabolites in the muscle [15]. This 'exercise-induced pain' (EIP) starts as a localised perception in the primary exercising muscle or muscles, but spreads to other locations over time, with significantly increased levels of pain experienced in areas as central as the chest at the end of exhaustive running [16]. In cycling, EIP has been shown to increase linearly with both power output and physiological intensity [17], whilst in self-paced exercise EIP appears to increase linearly as a function of distance completed or time elapsed, with near maximal levels experienced at the end of exercise [10]. Consequently, an individual exerting an intense and sustained physiological intensity will experience a level of pain and discomfort that is extremely unpleasant [17]. This is important because pain is ultimately a protective function that serves as a warning of actual or impending tissue damage and therefore a stimulus to disengage with the action or behaviour that is causing it [18]. Therefore, the drive to disengage with the exercise task, or reduce intensity to reduce the level of pain, will become increasingly stronger as the exercise continues [14, 19].

This association between pain and exercise no doubt provides the basis of long-held beliefs that pain tolerance is an important pre-requisite for athletic performance (i.e. no pain, no gain). It is important to note that pain tolerance (i.e. the maximum level of perceived pain that someone is able to tolerate, or the duration someone is willing to be exposed to a given pain intensity) is very different to pain threshold (the level at which a stimulus is initially perceived as pain), and the distinction between these should be

emphasised and assessed with appropriate methods so that the role of pain in exercise performance is better understood. However, even when this distinction is made, the impact of pain on exercise performance can be difficult to empirically test, so that despite the clear reference to its significance by competitors [19, 20], the experimental evidence supporting this notion is surprisingly limited and often inconsistent. A meta-analysis examining pain perception differences between athletes and active controls showed that in most well-controlled studies, athletes exhibited a higher pain tolerance, despite similar pain thresholds [21]. This is logical, given that EIP tends to reach only moderate or strong levels, but must be tolerated for the duration of the exercise. A recent investigation demonstrated that specific tolerance to EIP was not only strongly correlated with endurance performance, but could also predict the completion time of a 10 mile cycling time trial [22]. Even when EIP tolerance was combined with traditional measures of endurance performance (maximal oxygen uptake, gas exchange threshold and peak power output), it was still able to explain some of the variance (7.5% variance, $p=0.002$) in time trial completion time [22]. It is likely that this increased tolerance of pain arises from particular physiological or psychological adaptations that occur as a result of exercise training [8, 21]. Indeed, an improvement in pain tolerance has been an outcome of aerobic training programs in previously untrained individuals [23-25], which the authors attributed predominantly to psychological adaptations to training (although this was not measured). The increased pain tolerance could have arisen as a result of diminished signalling in response to the noxious stimulus [26-28], but this desensitisation of the nociceptors is hard to measure and unlikely given that these studies also observed no change in pain threshold. A more likely explanation is that exposure to frequent and unpleasant sensory experiences during aerobic training permits the participants to develop a means of coping and a tolerance to these

123 sensations. Indeed, coping skills can increase pain control [29] and the attitude of
124 athletes towards pain has been shown to differ from that of normally active controls
125 [30]. Recent evidence suggests that highly trained individuals (>10 h per week training
126 and competing at, or above the top tier of their local competition) demonstrate
127 significantly enhanced conditioned pain modulation compared with non-athletic
128 controls [8]. This suggests that highly trained athletes have a higher endogenous pain
129 inhibition ability, and that this may reduce the level of pain experienced at a given
130 exercise intensity. However, it is not clear whether this is something that develops in
131 response to training, or is an inherent characteristic that naturally predisposes these
132 individuals to engaging in competitive sport.
133
134 Whilst the relationship between perception of pain and athletic performance is
135 undoubtedly complex, the general consensus is that EIP forms an important part of
136 endurance exercise [14]. At the very least, EIP likely provides important information
137 for the exerciser regarding the current level of fatigue, and this information can
138 subsequently be used to effectively pace an exercise bout [20]. Thoughts of pain and
139 discomfort were shown to be prominent at the start of a 10 mile TT, but reduced towards
140 the final sections of the race [20]. A similar observation was shown for thoughts relating
141 to the monitoring or altering of pace, which may be indicative of the riders using the
142 conscious feeling of pain and discomfort as a means to guide decisions on the
143 appropriateness of their pacing strategy. This suggestion is supported by studies that
144 have examined the effect of analgesia during exercise [10, 31, 32], where a decreased
145 sensation of pain usually results in an increased power output being sustained and a
146 significantly improved endurance performance (i.e. pacing strategy is adjusted when
147 pain perception is moderated). When pain was completely blocked during exercise

through the injection of intrathecal fentanyl, participants adopted an extremely positive pacing strategy, which resulted in very high levels of peripheral fatigue and a sub-optimal performance [33]. Whilst the purpose of this study was not to examine the effect of pain during exercise, it does support the notion that EIP is an important sensation in determining beneficial pacing during self-paced exercise.

3.2. Applications

Use of analgesics and/or their abuse in pursuit of athletic performance enhancement, or to mask injury to facilitate participation/competition, is not uncommon. The ethics and motivation for their use by athletes is beyond the scope of this review and has been discussed elsewhere [34-38], however their use and abuse appears rife within elite and recreational sport [39-43].

Although analgesia use appears widespread, empirical evidence supporting their use to augment athletic performance is sparse, particularly in elite populations (i.e. maximal aerobic power of 350-500 W and maximal aerobic capacity of 72-80 $mL \cdot kg^{-1} \cdot min^{-1}$ [44]). Trained but non-elite cyclists improved their 10 mile (16.1 km) time-trial completion time when ingesting 1.5 g of acetaminophen (paracetamol) compared to placebo [10]. Faster completion time was accompanied by elevated heart rate and blood lactate (increased intensity), yet perceived pain and exertion were not different to the placebo condition [10]. Indeed, augmented mean power output was seen across repeated anaerobic Wingate tests in response to the same dose of acetaminophen within a recreationally active population, again attributed to reduced EIP [32]. However, data from elite populations regarding analgesia use and their effects on exercise performance is lacking, despite their widespread use in sport [34, 35, 40-42].

Transcranial direct current stimulation (tDCS) has shown some efficacy to reduce perceived pain (without the reliance on pharmacological analgesia), albeit in a cold pressor test within healthy recreationally active individuals [45]. This reduction in perceived pain was not seen in subsequent time to exhaustion (TTE) cycling exercise, i.e. EIP was unchanged as were performance metrics [45]. Similar findings of increased pain inhibitory capacity from high definition tDCS without enhancement of subsequent force production or attenuation of muscular fatigue have been reported elsewhere [46]. Indeed, the efficacy of tDCS to augment exercise performance, whether through favourable EIP modulation or otherwise, across various exercise modalities is currently ambiguous and the available evidence is often limited by poor ecological validity and experimental design [47-50]. Transcutaneous electrical nerve stimulation (TENS) has shown greater promise, whereby reduced EIP translated to increased TTE during isometric muscular contractions in healthy recreationally active participants [31]. However, the evidence pertaining to tDCS/TENS and EIP outlined above, relative to exercise performance, should be interpreted with caution. This is particularly the case when considering implementation of these findings in elite athletes (which the evidence above does not examine), given that such athletes have altered pain perception compared to their non-elite or recreational counterparts [8]. Indeed, evidence from such elite populations is required to inform practice, as is further discussion of any ethical/doping issues by germane authorities and medical support staff relative to tDCS/TENS use in an attempt to enhance athletic performance.

Although limited and from unfit [23] or healthy but relatively untrained populations [8, 25], evidence suggests that pain tolerance can be favourably enhanced by aerobic

exercise training, whereas pain threshold remains unchanged. Triathletes have been shown to have greater tolerance and more efficient pain modulation than non-athletic controls [51]. But whether practitioners can evoke performance improvements in well-trained individuals by increasing their pain tolerance, in lieu of physiological adaptations, requires further evidence from studies with robust experimental designs. Practitioners should also be aware that acute psychological stress might remove such advantages in pain modulation [52]. Indeed, a dose response was seen between athlete susceptibility to psychological stress, magnitude of perceived stress and any reduction in their advantageous pain modulation capacity [52]. Practitioners may therefore wish to deploy psychological skills training to develop robust stress resilience and coping strategies in their athletes; this will in turn help them retain optimal pain modulation/resilience capacity [29], which has been shown to be conducive to optimal exercise performance [22, 31]. The research determining the effects of interventions acting on pain perception and exercise performance is summarised in Table 1.

Insert Table 1 near here

While there is evidence of acetaminophen being beneficial during exercise, this is within a well-controlled research setting where interactions with other medications and supplements (anti-inflammatory drugs, caffeine, etc.), other common conditions (redistribution of blood flow, hydration status, diet, etc.) and the intensity of exercise are explicitly controlled to ensure the safety of participants. Acetaminophen use during exercise where such robust controls are not in place must not be advocated; the risk of interaction with any of the stated factors is not known and may increase the danger of injury/damage to an individual.

4.0. Perceptions of Temperature and Endurance Performance

4.1 Theory

The perception of temperature (as measured by ratings of 'thermal sensation') and the perception of comfort associated with that temperature (as measured by ratings of 'thermal comfort') have been studied in exercising humans through psychometric scales since the late 1960s [53]. Warm and uncomfortable perceptions of temperature develop during prolonged exercise [54], and become exacerbated when exercise is completed in hot compared to cool environments [53, 55]. Thermal sensation and thermal comfort ratings should be treated separately however, as they are measured with separate scales, and do not always act in unison [56]. Original investigations demonstrated that at the onset of exercise, the increasingly hotter thermal sensation and thermal discomfort ratings are both associated with changes in the mean body temperature [53]. During steady state exercise of 30-40 minutes duration however, thermal sensation ratings can be attributed to skin and ambient air temperatures, while the thermal discomfort ratings can be attributed to the skin blood flow and the sweating response [53]. Thus, the psychophysiological basis of thermal sensation and discomfort ratings are likely governed by sensory and effector mechanisms, respectively. Detailed molecular, neuroanatomical, and neurophysiological mechanisms that allow humans to sense temperature have been described recently [57] and therefore this section will focus on relationships between ratings of thermal sensation, thermal discomfort and exercise performance. Lastly, it should be noted that the term 'sensation' (i.e. the process of a sensory receptor being stimulated) has been used incorrectly in this context, and this has become ingrained within the thermoregulation literature. The term thermal 'sensation' as it is described across the literature, is in fact a 'perception' (i.e. how one

interprets how they feel). Considering that the term ‘thermal sensation’ is used as a measure of thermal perception (i.e. thermal sensation ratings) frequently in the current literature, we will continue to use it throughout this review (albeit erroneously) when discussing research that has measured these ratings of thermal sensation.

The exercise, heat stress, and cooling literature has demonstrated associations between thermal perceptions, body temperatures and endurance exercise capacity or performance [56, 58-62]. However, reviews and models based on early thermoregulation research predominately implicated an elevated core body temperature as the direct cause of fatigue and impaired performance in the heat [63-66], while the role of thermal perceptions in pacing, fatigue and performance received little attention in comparison. However, Cheung [67] presented three forms of evidence to describe a possible relationship between thermal perceptions and exercise performance. Evidence included: a) alterations in pacing during exercise with heating and cooling interventions [68, 69]; b) the observation of reduced thermal perceptions following aerobic conditioning [9] and; c) the capacity for psychological skills training to improve running performance in the heat [70]. Nevertheless, the clarity of any relationship between thermal perception and exercise performance was still limited by the inability to separate thermal perception from thermal state (i.e. body temperatures) during exercise. Recently however, the use of the cooling compound menthol (which acts on cutaneous thermoreceptors to induce perceptions of coolness) has allowed the separation of thermal perceptions and thermal state within research, although the core temperature can be altered by menthol in some instances where it is applied to a large body surface area [71, 72]. Pre-cooling (i.e. cold water immersion, ice slurry ingestion, etc.) cannot reduce thermal perceptions without simultaneously reducing the thermal

state (reductions in physical body temperatures e.g. various tissue temperatures). Hence, placebo controlled research that shows performance improvement with menthol application alone can be attributed to thermal perception only, as body temperatures generally remain unchanged [73-75].

In the first study to use menthol to separate thermal perceptions and thermal state, Schlader et al. [60] demonstrated that under heat stress, cooling of the face via both thermal (forced convection) and non-thermal (menthol gel) methods significantly increased power output in a RPE-clamp cycling protocol compared to a control. Hence, both the facial temperature and the perception of that temperature were demonstrated to be involved in the regulation of exercise (via alterations in the RPE). Another comparison between a menthol mouth rinse (that lowered thermal sensation ratings) and an ice slurry beverage (that lowered core body temperature) revealed that only the menthol intervention significantly improved running performance in the heat [76]. Hence, not only does thermal perception play a role in endurance pacing and performance, but this role may in fact be more influential than the physical temperature of the body in some circumstances of mild-moderate heat stress.

Thermal perceptions are strongly influenced by the temperature of the face and head [77, 78], especially during heat exposure, where facial warming is most uncomfortable and facial cooling is most comfortable compared to other body segments [78]. As such, focused cooling of the face and/or head during exercise has significantly increased cycling time to exhaustion [79], cycling work rate at a fixed RPE [60] and 5 km running performance time [80], all alongside a significantly cooler thermal sensation and/or increased thermal comfort rating. In an alternative model that required cyclists to

298 control their work rate to maintain thermal comfort in cold conditions, facial warming
299 and cooling significantly decreased and increased work rate, respectively [81]. Such
300 changes may be explained by the combination of a greater density of thermal afferents
301 in the face along with processing mechanisms in the central nervous system [77, 78].
302 Considering these interventions did not influence core body temperature, it is evidence
303 for peripheral temperatures modulating endurance exercise performance, likely via
304 mechanisms involving thermal perception [82].
305
306 Adjustments in pacing during exercise in the heat have also been observed prior to a
307 physical change in the core body temperature [69, 83, 84], which suggests the
308 involvement of afferent feedback and psychological factors in the regulation of
309 endurance exercise performance in the heat. Thermal perception is not the ultimate
310 afferent modulator of exercise performance however, since when the skin temperature
311 was altered from cold to hot and vice-versa during a 60 minute cycling time trial,
312 changes in the work-rate did not mirror changes in thermal perception across the trial
313 [85]. Instead, the exercise intensity was more closely associated with the RPE, which
314 is likely influenced by thermal perception [86, 87], but to a greater extent by the
315 combination of whole-body feedback [88], exercise end-point [89] and feed-forward
316 processes in the brain (corollary discharge) [7]. Hence, a previous review of the this
317 evidence [82] proposed that during exercise in the heat, elevated skin temperature and
318 the associated thermal perceptions are responsible for a self-selected reduction in work-
319 rate in order to maintain the RPE and allow the completion of an exercise task without
320 hyperthermia.
321

Studies whereby exercise performance was altered following application of menthol interventions allow confirmation that exercise performance is influenced by temperature perception [60, 76]. Such influence is maximised when cooling or heating interventions are targeted to the face and/or head [60, 79-81, 85], due to the high alliesthesial thermosensitivity of these areas [77]. Such afferent feedback is but one piece of the puzzle with an influence on the overall RPE, pacing and endurance performance, particularly relevant during exercise heat-stress. Lastly, it should be noted that this cooling research is limited by both the dominance of research participants being male (data on females are needed), as well as the difficulty in implementing a robust placebo condition relative to the experimental designs, and hence, future researchers should address these limitations. Another interesting avenue for future research would be to discriminate the relative contributions of thermal sensation and comfort ratings to endurance performance.

4.2 Applications

Thermal perception and comfort, as outlined above, can be important relative to exercise pacing and performance [60, 69, 80]. Heat adaptation (HA) through acclimation/acclimatisation, provides robust protection [90] to thermally mediated reductions in endurance [91] and repeated sprint [92] exercise performance, and against heat illnesses [93]. Relevant to thermal perception and this review article, appropriate HA has efficacy to improve ratings of thermal sensation/comfort in team sport and endurance athletes during subsequent training and competition in the heat [90, 94-96]. However, these HA-mediated thermal sensation rating changes are not seen in resting values but rather at comparative time-points or as a mean value across an exercise/training bout, when comparisons are made pre- to post-HA [90]. An

empirically rationalised [90] attainment of a heat acclimated phenotype may be practically challenging however, as only 13% of long distance athletes at the 2015 World Athletics Championships (where hot conditions were expected) followed a HA regime [97]. Therefore, while HA is recommended to favourably alter thermal sensation ratings during heat-strain, it may not be practically viable for all athletes.

Exercise training per se and increasing 'fitness' has been shown to dissociate physiological and perceptual (i.e. thermal sensation ratings) exercise heat-stress, specifically a dampening of perceptual relative to physiological heat-strain [67]. Thus attainment of high-levels of fitness appears an important counter-measure to reduce perceived temperature during exercise. However, practitioners should exercise caution with highly-trained athletes given their dampened interpretation of thermal perception and ability to produce sustained highly elevated core temperatures [67]; this potentially places these populations at heat illness risk, especially when appropriate HA has not been performed [91]. Moreover, high body mass-index with low running ability also predisposed Marine Corps recruits to heat illness [98], indicating that practitioners should adopt individualised practice across populations of all fitness levels engaging in exercise heat-stress.

Aside from increasing 'fitness' and procurement of a HA phenotype, there are acute strategies which can positively alter ratings of thermal sensation/comfort prior to [99] and during exercise [11, 100] with subsequent ergogenic performance effects. Challenges whereby delineation of changes in thermal sensation rating without concomitant core body temperature rise have been overcome, demonstrating that a pre-exercise change in the rating of thermal sensation alone (no change in core temperature, only skin temperature) can beneficially alter subsequent athletic performance [101]. A

pre-exercise cooling strategy with efficacy to reduce body temperatures can create a 'heat-sink' [102], i.e. a larger heat storage capacity [99] which delays the attainment of an individualised high body temperature at which performance decrements begin to occur. Similarly, a mid-exercise alteration in the rating of thermal sensation via facial water spraying (without change in core body temperature) has been demonstrated to be equally as ergogenic to running performance as pre-cooling by cold water immersion [80]. Therefore, evidence for strategies that reduce the rating of thermal sensation, rather than the core body temperature only, to enhance endurance performance may drive their use by practitioners and athletes [11, 76, 80, 99, 100, 102].

Practitioners should employ evidence-based practice regarding their choice of thermal perception orientated mid-cooling strategies and the tissue to be targeted, relative to attempts to improve exercise performance. This evidence must be considered carefully, given that few laboratory investigations have applied realistic facing wind speeds, without which, artificially high body temperatures are seen along with over-estimates of the beneficial effects of a cooling intervention (thermal perception or body temperature orientated), thus translation from the laboratory to the 'field' may be lacking [100]. The following recommendations are therefore based on the criteria that an appropriate facing winding speed was employed, a time-trial rather than time-to-exhaustion performance measure was used and the mid-cooling strategy employed was highly practical (i.e. frozen items were not consumed or applied to the body). Menthol mouth rinse and/or facial water spray mid-cooling has proven ergogenic to 5 km running time trial performance at ~33^{o}C and ~46 % relative humidity [76, 80]. Indeed, lowering rating of thermal sensation via menthol mouth rinse during exercise improved running performance whereas pre-cooling via ice slurry (which lowered core temperature) did not [76]. Even when pre-cooling was administered through cold water

396 immersion (which lowered core temperature), its ergogenic performance effect was
397 matched by mid-cooling (which lowered rating of thermal sensation rather than core
398 temperature) via facial water spray [80]. Practitioners should therefore seek to either
399 utilise menthol mouth rinse and/or facial water spray in an attempt to improve
400 endurance performance via thermal perception. Practitioners should avoid the
401 temptation to use menthol as a gel or spray to the skin/clothing, as exercise performance
402 is not enhanced concomitantly alongside ratings of thermal sensation [73, 103, 104],
403 likely due to the capacity of menthol to reduce sweat rate and vasoconstrict blood
404 vessels, ultimately promoting heat storage when applied to a large surface area of the
405 body [11]. Alterations of thermal sensation rating through menthol use during exercise
406 appear most beneficial to performance when applied internally via a beverage
407 containing menthol or through mouth rinsing [11]. Relative to mid-cooling via facial
408 water spray, it is important for practitioners to consider that the skin on the torso
409 demonstrates lower alliesthesial thermosensitivity compared to the face [57]. Indeed,
410 cooling of the face can be up to five times more effective than cooling of other body
411 surfaces relative to maintenance of thermal comfort [77]. However, it must be noted
412 that further research is required in this regard, for example, currently there is no
413 evidence whether menthol mouth rinse can be combined with facial water spray to elicit
414 an additive performance effect. Furthermore, optimised and field compatible mid-
415 cooling strategies will evidently be discipline specific according to what might be
416 practical and within the rules of the different sports.

417 A high variance in the efficacy of psychological skills training has been demonstrated
418 relative to several interrelated agendas appropriate to exercise performance
419 optimisation, including tolerance to environmental extremes and pain [70, 105, 106].
420 However, there is evidence [70] that psychological skills training (including goal

setting, arousal regulation, mental imagery and positive self-talk) can augment distance covered during a 90-min treadmill run in the heat (30°C; 40% relative humidity). Recently, a specific form of psychological skills training, motivational self-talk, also enhanced endurance and cognitive function during cycling based exercise heat stress [105]. Positive performance effects were attributable to the two week motivational self-talk intervention allowing participants to tolerate a longer duration near or at a maximal intensity (i.e. ≥ RPE 19) and a higher terminating core temperature without difference in heart rate or oxygen uptake (i.e. greater psychological tolerance to thermophysiological strain) compared to the control group without motivational self-talk, demonstrating that even in well-trained individuals, enhanced psychological tolerance to thermal or exercise discomfort is trainable and pliable via motivational self-talk [105]. Practitioners may consider adding appropriate exercise heat-stress orientated psychological skills training to their athlete's training regime, but whether or not such psychological skills training could provide an additive effect to the above outlined ergogenic thermal perception orientated interventions/approaches requires further research. The research determining the effects of interventions acting on temperature perception and exercise performance is summarised in Table 2.

Insert Table 2 near here

5.0. Interconnections Between Pain and Temperature Perception

While acetaminophen is best known as a pain reliever, it can also be used to treat fever [107] and hence, it may play a role in thermoregulation and temperature perception. In athletes, an acute dose of acetaminophen can delay the increase in body temperature during exercise in the heat [108, 109] and has also been demonstrated to significantly

lower thermal sensation rating and improve cycling time to exhaustion in the heat [109]. However, no such changes were observed following ingestion of acetaminophen when cycling at a fixed rate of metabolic heat production [110].

Another interconnection between pain and temperature perception is evident when considering the use of a cold stimulus as a widespread treatment for pain [111]. Research on menthol application may also allow separation between the physical and perceptual effects of a cold stimulus when considering pain responses. Menthol has been added to commercially available medical creams targeting musculoskeletal pain as it may have an analgesic effect for sports injuries, delayed onset muscle soreness and arthritis [112, 113]. Menthol has been demonstrated to cool not only through the transient receptor potential cation channel subfamily M member 8 (TRPM8), but also to inhibit the transient receptor potential cation channel, subfamily A, member 1 (TRPA1), which is a mediator of inflammatory pain [114]. While topical application of menthol decreased perceived pain and improved physical function in patients with knee osteoarthritis [115], research to date has not investigated the analgesic effects of menthol during exercise in athletes.

6.0. Safety Considerations

Athletes who attempt to override the perceptions of pain during intense endurance exercise may be at risk of serious injury. Given that pain serves as a protective function for the body, it could be suggested that moderating the naturally occurring EIP signals during exercise poses a danger to the individual. While no participants suffered long-term damage as a result of a complete afferent (and consequently pain) block [33], suggesting any regulation of intensity arising from EIP is likely minimal, the risks of

such methods in all athletic situations are largely unexplored. It should also be noted that harmful side effects have been reported with analgesic use during and relative to sport, including gastrointestinal damage, as well as liver and kidney failure [39, 116-118]. Therefore, this article does not condone the use of analgesics to mask the pain from injury during competition, nor in the pursuit of performance enhancement. Use of analgesic strategies must be considered carefully by the athlete and under guidance from their medical support team/physician.

The perception of temperature may also be a protective mechanism to preserve against exertional heat illness and hyperthermia. The ability to perceive the temperature of the environment and adjust behaviour accordingly is undoubtedly a vital attribute in humans that has ensured long-lasting survival [57]. These perceptions may also play a similar role during endurance exercise (especially in the heat) to encourage the exerciser to slow down or stop when it may be otherwise dangerous to continue [82, 119]. Therefore, application of menthol or cooling methods that target perception should be avoided close to the onset of hyperthermia, to allow perception of symptoms associated with high levels of heat stress, adjustment to self-selected exercise intensity and the prevention of heat injury.

While no specific evidence of serious health complications from the use of these strategies has been described within the literature to date, it is wise to exercise the cautions described above. Indeed, the absence of such evidence is likely due to a lack of specific designs to test such hypotheses, which would be mostly unethical in humans.

7.0. Conclusion

New research developments have demonstrated that the perceptions of pain and temperature can both influence endurance exercise performance. These findings highlight the importance of the brain in the development of endurance training adaptations, and in the development of fatigue during acute exercise. While interventions are available to modify the perceptions of pain and temperature to improve performance in endurance-trained populations, these interventions may have serious health consequences when used during intense exercise and/or in thermally challenging environments and as such, a medical practitioner should supervise any use of such interventions closely to ensure the safety of the athlete.

Compliance with Ethical Standards

Funding No sources of funding were used to assist with the preparation of this article.

Conflict of interest Christopher Stevens, Alexis Mauger, Peter Hassmen and Lee Taylor declare that they have no conflicts of interest relevant to the content of this review.

References

1. Joyner MJ, Coyle EF. Endurance exercise performance: The physiology of champions. J Physiol. 2008 Jan 1;586(1):35-44.

2. Marcora SM, Staiano W. The limit to exercise tolerance in humans: Mind over muscle? Eur J Appl Physiol. 2010 Jul;109(4):763-70.

3. Noakes TD. Time to move beyond a brainless exercise physiology: The evidence for complex regulation of human exercise performance. Appl Physiol Nutr Metab. 2011 Feb;36(1):23-35.

4. St Clair Gibson A, Noakes TD. Evidence for complex system integration and dynamic neural regulation of skeletal muscle recruitment during exercise in humans. Br J Sports Med. 2004 Dec;38(6):797-806.

5. St Clair Gibson A, Swart J, Tucker R. The interaction of psychological and physiological homeostatic drives and role of general control principles in the regulation of physiological systems, exercise and the fatigue process - The Integrative Governor theory. Eur J Sport Sci. In Press. doi: 10.1080/17461391.2017.1321688:1-12.

6. Abbiss CR, Peiffer JJ, Meeusen R, et al. Role of ratings of perceived exertion during self-paced exercise: What are we actually measuring? Sports Med. 2015 Sep;45(9):1235-43.

7. Marcora S. Counterpoint: Afferent feedback from fatigued locomotor muscles is not an important determinant of endurance exercise performance. J Appl Physiol. 2010 Feb;108(2):454-7.

8. Flood A, Waddington G, Thompson K, et al. Increased conditioned pain modulation in athletes. J Sports Sci. 2017 Jun;35(11):1066-72.

9. Tikuisis P, McLellan TM, Selkirk G. Perceptual versus physiological heat strain during exercise-heat stress. Med Sci Sports Exerc. 2002 Sep;34(9):1454-61.

10. Mauger AR, Jones AM, Williams CA. Influence of acetaminophen on performance during time trial cycling. J Appl Physiol (1985). 2010 Jan;108(1):98-104.

11. Stevens CJ, Best R. Menthol: A fresh ergogenic aid for athletic performance. Sports Med. 2016 Nov;47(6):1035-42.

12. Merskey H, Spear FG. The concept of pain. J Psychosom Res. 1967 Jun;11(1):59-67.

13. Hadjistavropoulos T, Craig KD. A theoretical framework for understanding self-report and observational measures of pain: A communications model. Behav Res Ther. 2002 May;40(5):551-70.

14. Mauger AR. Factors affecting the regulation of pacing: Current perspectives. Open Access J Sports Med. 2014;5:209-14.

15. O'Connor PJ, Cook DB. Exercise and pain: The neurobiology, measurement, and laboratory study of pain in relation to exercise in humans. Exerc Sport Sci Rev. 1999;27:119-66.

16. Slapsinskaite A, Razon S, Balague Serre N, et al. Local pain dynamics during constant exhaustive exercise. PLoS One. 2015;10(9):e0137895.

17. Cook DB, O'Connor PJ, Eubanks SA, et al. Naturally occurring muscle pain during exercise: Assessment and experimental evidence. Med Sci Sports Exerc. 1997 Aug;29(8):999-1012.

18. Navratilova E, Porreca F. Reward and motivation in pain and pain relief. Nat Neurosci. 2014 Oct;17(10):1304-12.

19. Kress J, Statler T. A Naturalistic investigation of former olympic cyclists' cognitive strategies for coping with exertion pain during performance. J Sport Behav. 2007;30(4):428-52.

20. Whitehead AE, Jones HS, Williams EL, et al. Changes in cognition over a 16.1 km cycling time trial using Think Aloud protocol: Preliminary evidence. Int J Sport Exerc Psychol. In Press. doi: 10.1080/1612197x.2017.1292302:1-9.

21. Tesarz J, Schuster AK, Hartmann M, et al. Pain perception in athletes compared to normally active controls: A systematic review with meta-analysis. Pain. 2012 Jun;153(6):1253-62.

22. Astokorki AH, Mauger AR. Tolerance of exercise-induced pain at a fixed rating of perceived exertion predicts time trial cycling performance. Scand J Med Sci Sports. 2017 Mar;27(3):309-17.

23. Anshel MH, Russell KG. Effect of aerobic and strength training on pain tolerance, pain appraisal and mood of unfit males as a function of pain location. J Sports Sci. 1994;12(6):535-47.

24. O'Leary TJ, Collett J, Howells K, et al. High but not moderate-intensity endurance training increases pain tolerance: A randomised trial. Eur J Appl Physiol. 2017 Nov;117(11): 2201-10.

25. Jones MD, Booth J, Taylor JL, et al. Aerobic training increases pain tolerance in healthy individuals. Med Sci Sports Exerc. 2014 Aug;46(8):1640-7.

26. Micalos PS. Perspectives on biochemical and neurosensory mechanisms for exercise-induced pain inhibition. Fatigue. 2014 Sep;2(4):219-30.

27. Micalos PS, Drinkwater EJ, Cannon J, et al. Reliability of the nociceptive flexor reflex (RIII) threshold and association with pain threshold. Eur J Appl Physiol. 2009 Jan;105(1):55-62.

28. Micalos PS, Korgaonkar MS, Drinkwater EJ, et al. Cerebral responses to innocuous somatic pressure stimulation following aerobic exercise rehabilitation in chronic pain patients: A functional magnetic resonance imaging study. Int J Gen Med. 2014 Aug;7:425-32.

29. Birrer D, Morgan G. Psychological skills training as a way to enhance an athlete's performance in high-intensity sports. Scand J Med Sci Sports. 2010 Oct;20:78-87.

30. Nicholls AR, Polman RCJ. Coping in sport: A systematic review. J Sports Sci. 2007 Jan;25(1):11-31.

31. Astokorki AHY, Mauger AR. Transcutaneous electrical nerve stimulation reduces exercise-induced perceived pain and improves endurance exercise performance. Eur J Appl Physiol. 2017 Mar;117(3):483-92.

32. Foster J, Taylor L, Chrismas BC, et al. The influence of acetaminophen on repeated sprint cycling performance. Eur J Appl Physiol. 2014 Jan;114(1):41-8.

33. Amann M, Proctor LT, Sebranek JJ, et al. Opioid-mediated muscle afferents inhibit central motor drive and limit peripheral muscle fatigue development in humans. J Physiol. 2009 Jan;587(1):271-83.

34. Dietz P, Dalaker R, Letzel S, et al. Analgesics use in competitive triathletes: Its relationship to doping and on predicting its usage. J Sports Sci. 2016 Oct;34(20):1965-9.

35. Matava MJ. Ethical considerations for analgesic use in sports medicine. Clin Sports Med. 2016 Apr;35(2):227-43.

36. Morente-Sanchez J, Zabala M. Doping in sport: A review of elite athletes' attitudes, beliefs, and knowledge. Sports Med. 2013 Jun;43(6):395-411.

37. Lippi G, Sanchis-Gomar F. Acetaminophen and sport performance: Doping or what? Eur J Appl Physiol. 2014 Apr;114(4):881-2.

38. Mauger AR, Taylor L, Chrismas BCR, et al. Reply to letter: Acetaminophen and sport performance: Doping or what? Eur J Appl Physiol. 2014 Apr;114(4):883-4.

39. Kuster M, Renner B, Oppel P, et al. Consumption of analgesics before a marathon and the incidence of cardiovascular, gastrointestinal and renal problems: A cohort study. BMJ Open. 2013;3(4).

40. Tscholl PM, Dvorak J. Abuse of medication during international football competition in 2010-lesson not learned. Br J Sports Med. 2012 Dec;46(16):1140-1.

41. Tscholl PM, Vaso M, Weber A, et al. High prevalence of medication use in professional football tournaments including the World Cups between 2002 and 2014: A narrative review with a focus on NSAIDs. Br J Sports Med. 2015 May;49(9):580-U31.

42. Vaso M, Weber A, Tscholl PM, et al. Use and abuse of medication during 2014 FIFA World Cup Brazil: A retrospective survey. BMJ Open. 2015;5(9): e007608.

43. Brewer CB, Bentley JP, Hallam JS, et al. Use of analgesics for exercise-associated pain: Prevalence and predictors of use in recreationally trained college-aged students. J Strength Cond Res. 2014 Jan;28(1):74-81.

44. Jeukendrup AE, Craig NP, Hawley JA. The bioenergetics of world class cycling. J Sci Med Sport. 2000 Dec;3(4):414-33.

45. Angius L, Hopker JG, Marcora SM, et al. The effect of transcranial direct current stimulation of the motor cortex on exercise-induced pain. Eur Appl Physiol. 2015 Nov;115(11):2311-9.

46. Flood A, Waddington G, Keegan RJ, et al. The effects of elevated pain inhibition on endurance exercise performance. PeerJ. 2017 Mar;5:e3028.

47. Angius L, Hopker J, Mauger AR. The ergogenic effects of transcranial direct current stimulation on exercise performance. Front Physiol. 2017 Feb;8:90. doi: 10.3389/fphys.2017.00090

48. Edwards DJ, Cortes M, Wortman-Jutt S, et al. Transcranial direct current stimulation and sports performance. Front Human Neurosci. 2017 May;11:243. doi: 10.3389/fnhum.2017.00243

49. Barwood MJ, Butterworth J, Goodall S, et al. The effects of direct current stimulation on exercise performance, pacing and perception in temperate and hot environments. Brain Stimul. 2016 Dec;9(6):842-9.

50. Cunningham DA. Noninvasive brain stimulation enhances sustained muscle contractions by reducing neuromuscular fatigue: Implications for rehabilitation. J Neurophysiol. 2017 Mar;117(3):1215-7.

51. Geva N, Defrin R. Enhanced pain modulation among triathletes: A possible explanation for their exceptional capabilities. Pain. 2013 Nov;154(11):2317-23.

52. Geva N, Pruessner J, Defrin R. Triathletes lose their advantageous pain modulation under acute psychosocial stress. Med Sci Sports Exerc. 2017 Feb;49(2):333-41.

53. Gagge AP, Stolwijk JA, Saltin B. Comfort and thermal sensations and associated physiological responses during exercise at various ambient temperatures. Environ Res. 1969 Apr;2(3):209-29.

54. Pandolf KB, Cafarelli E, Noble BJ, et al. Perceptual responses during prolonged work. Percept Mot Skills. 1972 Dec;35(3):975-85.

55. Kamon E, Pandolf K, Cafarelli E. The relationship between perceptual information and physiological responses to exercise in the heat. J Hum Ergol (Tokyo). 1974 Sep;3(1):45-54.

56. Schulze E, Daanen HA, Levels K, et al. Effect of thermal state and thermal comfort on cycling performance in the heat. Int J Sports Physiol Perform. 2015 Jan 22;10(5):655-63.

57. Filingeri D. Neurophysiology of skin thermal sensations. Compr Physiol. 2016;6(3):1429.

58. Abbiss CR, Burnett A, Nosaka K, et al. Effect of hot versus cold climates on power output, muscle activation, and perceived fatigue during a dynamic 100-km cycling trial. J Sports Sci. 2010 Jan;28(2):117-25.

665 59. Peiffer JJ, Abbiss CR. Influence of environmental temperature on 40 km cycling
666 time-trial performance. Int J Sports Physiol Perform. 2011 Jun;6(2):208-20.
667 60. Schlader ZJ, Simmons SE, Stannard SR, et al. The independent roles of
668 temperature and thermal perception in the control of human thermoregulatory behavior.
669 Physiol Behav. 2011 May 3;103(2):217-24.
670 61. Kenny GP, Schissler AR, Stapleton J, et al. Ice cooling vest on tolerance for
671 exercise under uncompensable heat stress. J Occup Environ Hyg. 2011 Aug;8(8):484-
672 91.
673 62. Luomala MJ, Oksa J, Salmi JA, et al. Adding a cooling vest during cycling
674 improves performance in warm and humid conditions. J Therm Biol. 2012
675 Jan;37(1):47-55.
676 63. Cheung SS. Hyperthermia and voluntary exhaustion: Integrating models and
677 future challenges. Appl Physiol Nutr Metab. 2007 Aug;32(4):808-17.
678 64. Cheuvront SN, Kenefick RW, Montain SJ, et al. Mechanisms of aerobic
679 performance impairment with heat stress and dehydration. J Appl Physiol. 2010
680 Dec;109(6):1989-95.
681 65. Maughan RJ. Distance running in hot environments: A thermal challenge to the
682 elite runner. Scand J Med Sci Sports. 2010 Oct;20 Suppl 3:95-102.
683 66. Nybo L. Cycling in the heat: Performance perspectives and cerebral challenges.
684 Scand J Med Sci Sports. 2010 Oct;20 Suppl 3:71-9.
685 67. Cheung SS. Interconnections between thermal perception and exercise capacity
686 in the heat. Scand J Med Sci Sports. 2010 Oct;20 Suppl 3:53-9.
687 68. Booth J, Marino F, Ward JJ. Improved running performance in hot humid
688 conditions following whole body precooling. Med Sci Sports Exerc. 1997
689 Jul;29(7):943-9.

690 69. Tucker R, Marle T, Lambert EV, et al. The rate of heat storage mediates an
691 anticipatory reduction in exercise intensity during cycling at a fixed rating of perceived
692 exertion. J Physiol. 2006 Aug 1;574(Pt 3):905-15.
693 70. Barwood MJ, Thelwell RC, Tipton MJ. Psychological skills training improves
694 exercise performance in the heat. Med Sci Sports Exerc. 2008 Feb;40(2):387-96.
695 71. Gillis DJ, Barwood MJ, Newton PS, et al. The influence of a menthol and
696 ethanol soaked garment on human temperature regulation and perception during
697 exercise and rest in warm, humid conditions. J Therm Biol. 2016 May;58:99-105.
698 72. Kounalakis SN, Botonis PG, Koskolou MD, et al. The effect of menthol
699 application to the skin on sweating rate response during exercise in swimmers and
700 controls. Eur J Appl Physiol. 2010 May;109(2):183-9.
701 73. Barwood MJ, Corbett J, Thomas K, et al. Relieving thermal discomfort: Effects
702 of sprayed L-menthol on perception, performance, and time trial cycling in the heat.
703 Scand J Med Sci Sports. 2015 Jun;25 Suppl 1:211-8.
704 74. Flood TR, Waldron M, Jeffries O. Oral L-menthol reduces thermal sensation,
705 increases work-rate and extends time to exhaustion, in the heat at a fixed rating of
706 perceived exertion. Euro J Appl Physiol. 2017 Jul;117(7):1501-12.
707 75. Mundel T, Jones DA. The effects of swilling an L(-)-menthol solution during
708 exercise in the heat. Eur J Appl Physiol. 2010 May;109(1):59-65.
709 76. Stevens CJ, Thoseby B, Sculley DV, et al. Running performance and thermal
710 sensation in the heat are improved with menthol mouth rinse but not ice slurry
711 ingestion. Scand J Med Sci Sports. 2016 Sep 26;26(10):1209-16.
712 77. Cotter JD, Taylor NA. The distribution of cutaneous sudomotor and alliesthesial
713 thermosensitivity in mildly heat-stressed humans: An open-loop approach. J Physiol.
714 2005 May 15;565(1):335-45.

78. Nakamura M, Yoda T, Crawshaw LI, et al. Regional differences in temperature sensation and thermal comfort in humans. J Appl Physiol (1985). 2008 Dec;105(6):1897-906.

79. Ansley L, Marvin G, Sharma A, et al. The effects of head cooling on endurance and neuroendocrine responses to exercise in warm conditions. Physiol Res. 2008;57(6):863-72.

80. Stevens C, Kittel A, Sculley D, et al. Running performance in the heat is improved by similar magnitude with pre-exercise cold-water immersion and mid-exercise facial water spray. J Sports Sci. 2017;35(8):798-805.

81. Mundel T, Raman A, Schlader ZJ. Head temperature modulates thermal behavior in the cold in humans. Temperature. 2016 Apr-Jun;3(2):298-306.

82. Schlader ZJ, Stannard SR, Mundel T. Human thermoregulatory behavior during rest and exercise - a prospective review. Physiol Behav. 2010 Mar 3;99(3):269-75.

83. Tatterson AJ, Hahn AG, Martin DT, et al. Effects of heat stress on physiological responses and exercise performance in elite cyclists. J Sci Med Sport. 2000 Jun;3(2):186-93.

84. Tucker R, Rauch L, Harley YX, et al. Impaired exercise performance in the heat is associated with an anticipatory reduction in skeletal muscle recruitment. Pflügers Archiv. 2004 Jul;448(4):422-30.

85. Schlader ZJ, Simmons SE, Stannard SR, et al. Skin temperature as a thermal controller of exercise intensity. Eur J Appl Physiol. 2011 Jan 1;111(8):1631-9.

86. Bergh U, Danielsson U, Wennberg L, et al. Blood lactate and perceived exertion during heat stress. Acta Physiol Scand. 1986 Apr;126(4):617-8.

738 87. Pivarnik JM, Grafner TR, Elkins ES. Metabolic, thermoregulatory, and
739 psychophysiological responses during arm and leg exercise. Med Sci Sports Exerc.
740 1988 Feb;20(1):1-5.
741 88. Ulmer HV. Concept of an extracellular regulation of muscular metabolic rate
742 during heavy exercise in humans by psychophysiological feedback. Experientia. 1996
743 May 15;52(5):416-20.
744 89. Crewe H, Tucker R, Noakes TD. The rate of increase in rating of perceived
745 exertion predicts the duration of exercise to fatigue at a fixed power output in different
746 environmental conditions. Eur J Appl Physiol. 2008 Jul;103(5):569-77.
747 90. Tyler CJ, Reeve T, Hodges GJ, et al. The effects of heat adaptation on
748 physiology, perception and exercise performance in the heat: A meta-analysis. Sports
749 Med. 2016 Nov;46(11):1699-724.
750 91. Racinais S, Alonso JM, Coutts AJ, et al. Consensus recommendations on
751 training and competing in the heat. Br J Sports Med. 2015 Sep;49(18):1164-73.
752 92. Girard O, Brocherie F, Bishop DJ. Sprint performance under heat stress: A
753 review. Scand J Med Sci Sports. 2015 Jun;25 Suppl 1:79-89.
754 93. Taylor NA. Human heat adaptation. Compr Physiol. 2014 Jan;4(1):325-65.
755 94. Sunderland C, Morris JG, Nevill ME. A heat acclimation protocol for team
756 sports. Br J Sports Med. 2008 May;42(5):327-33.
757 95. Kelly M, Gastin PB, Dwyer DB, et al. Short duration heat acclimation in
758 Australian football players. J Sports Sci Med. 2016 Mar;15(1):118-25.
759 96. James CA, Richardson AJ, Watt PW, et al. Short-term heat acclimation
760 improves the determinants of endurance performance and 5-km running performance
761 in the heat. Appl Physiol Nutri Metab. 2017 Mar;42(3):285-94.

762 97. Périard JD, Racinais S, Timpka T, et al. Strategies and factors associated with
763 preparing for competing in the heat: A cohort study at the 2015 IAAF World Athletics
764 Championships. Br J Sports Med. 2017;51(4):264-71.
765 98. Gardner JW, Kark JA, Karnei K, et al. Risk factors predicting exertional heat
766 illness in male Marine Corps recruits. Med Sci Sports Exerc. 1996 Aug;28(8):939-44.
767 99. Bongers CC, Hopman MT, Eijsvogels TM. Cooling interventions for athletes:
768 An overview of effectiveness, physiological mechanisms, and practical considerations.
769 Temperature. 2017;4(1):60-78.
770 100. Stevens CJ, Taylor L, Dascombe BJ. Cooling during exercise: An overlooked
771 strategy for enhancing endurance performance in the heat. Sports Med. 2017
772 May;47(5):829-41.
773 101. Kay D, Taaffe DR, Marino FE. Whole-body pre-cooling and heat storage during
774 self-paced cycling performance in warm humid conditions. J Sports Sci. 1999
775 Dec;17(12):937-44.
776 102. Bongers C, Thijssen DHJ, Veltmeijer MTW, et al. Precooling and percooling
777 (cooling during exercise) both improve performance in the heat: A meta-analytical
778 review. Br J Sports Med. 2015 Mar;49(6):377-84.
779 103. Barwood MJ, Corbett J, White D, et al. Early change in thermal perception is
780 not a driver of anticipatory exercise pacing in the heat. Br J Sports Med. 2012
781 Oct;46(13):936-42.
782 104. Barwood MJ, Corbett J, White DK. Spraying with 0.20% L-Menthol does not
783 enhance 5k running performance in the heat in untrained runners. J Sports Med Phys
784 Fitness. 2014 May 20;54(5):595-604.

105. Wallace PJ, McKinlay BJ, Coletta NA, et al. Effects of motivational self-talk on endurance and cognitive performance in the heat. Med Sci Sports Exerc. 2017 Jan;49(1):191-9.

106. McCormick A, Meijen C, Marcora S. Psychological determinants of whole-body endurance performance. Sports Med. 2015 Jul;45(7):997-1015.

107. Feldberg W, Gupta KP, Milton AS, et al. Effect of bacterial pyrogen and antipyretics on prostaglandin activity in cerebrospinal fluid of unanaesthetized cats. Br J Pharmacol. 1972 Nov;46(3):550p-1p.

108. Burtscher M, Gatterer H, Philippe M, et al. Effects of a single low-dose acetaminophen on body temperature and running performance in the heat: A pilot project. Int J Physiol Pathophysiol Pharmacol. 2013;5(3):190-3.

109. Mauger AR, Taylor L, Harding C, et al. Acute acetaminophen (paracetamol) ingestion improves time to exhaustion during exercise in the heat. Exp Physiol. 2014 Jan;99(1):164-71.

110. Coombs GB, Cramer MN, Ravanelli NM, et al. Acute acetaminophen ingestion does not alter core temperature or sweating during exercise in hot-humid conditions. Scand J Med Sci Sports. 2015 Jun;25 Suppl 1:96-103.

111. Swenson C, Sward L, Karlsson J. Cryotherapy in sports medicine. Scand J Med Sci Sports. 1996 Aug;6(4):193-200.

112. Eccles R. Menthol and related cooling compounds. J Pharm Pharmacol. 1994 Aug;46(8):618-30.

113. Johar P, Grover V, Topp R, et al. A comparison of topical menthol to ice on pain, evoked tetanic and voluntary force during delayed onset muscle soreness. Int J Sports Phys Ther. 2012 Jun;7(3):314-22.

114. Macpherson LJ, Hwang SW, Miyamoto T, et al. More than cool: Promiscuous relationships of menthol and other sensory compounds. Mol Cell Neurosci. 2006 Aug;32(4):335-43.

115. Topp R, Brosky JA, Jr., Pieschel D. The effect of either topical menthol or a placebo on functioning and knee pain among patients with knee OA. J Geriatr Phys Ther. 2013 Apr-Jun;36(2):92-9.

116. Alaranta A, Alaranta H, Helenius I. Use of prescription drugs in athletes. Sports Med. 2008;38(6):449-63.

117. Gorski T, Cadore EL, Pinto SS, et al. Use of NSAIDs in triathletes: prevalence, level of awareness and reasons for use. Br J Sports Med. 2011 Feb;45(2):85-90.

118. Van Wijck K, Lenaerts K, Van Bijnen AA, et al. Aggravation of exercise-induced intestinal injury by Ibuprofen in athletes. Med Sci Sports Exerc. 2012 Dec;44(12):2257-62.

119. Marino FE. The evolutionary basis of thermoregulation and exercise performance. Med Sport Sci. 2008;53:1-13.

Table 1: Research determining the effects of interventions acting on pain perception and exercise performance.

Study	Participants	Intervention	Testing Protocol	Testing Conditions	Outcomes
Mauger et al. 2010 [10]	13 males, VO_2max = 65 ± 5 $mL \cdot kg^{-1} \cdot min^{-1}$	Oral ingestion of 1.5 g acetaminophen, 60 min prior to TT	16.1 km cycling TT	Not stated	↓ TT time by 30 s (2%) ↑ Mean PO by 10 W (4%) ↔ EIP, ↔ RPE, ↑ HR, ↑ BLa
Astokorki and Mauger, 2017 [31]	Part 1: 11 males, 7 females. Part 2: 14 males, 8 females, VO_2max = 53 ± 7 $mL \cdot kg^{-1} \cdot min^{-1}$	Biphasic TENS applied unilaterally to BF muscle (Part 1) or bilaterally to VL muscle (Part 2) in a continuous pattern with pulse width of 300 μs and frequency of 100 Hz	Part 1: 20% MVC of biceps flexion held at 90° isometrically until exhaustion. Part 2: 16.1 km cycling TT	Not stated	Part 1: ↑ TTE by 3 min (38%) ↓ EIP, ↔ RPE Part 2: ↓ TT time by 33 s (2%) ↓ EIP, ↔ RPE, ↑ HR, ↑ BLa
Foster et al. 2014 [32]	9 males, VO_2max = 47 ± 6 $mL \cdot kg^{-1} \cdot min^{-1}$	Oral ingestion of 1.5 g acetaminophen, 30 min prior to test	8 x 30 s Wingate tests, with 2 min active rest intervals	Not stated	↑ Mean PO by 19 W (5%) ↔ EIP, ↔ Peak PO, ↔ HR
Angius et al. 2015 [45]	Part A: 9 males, VO_2max = 48 ± 7 $mL \cdot kg^{-1} \cdot min^{-1}$ Part B: 7 males	tDCS applied to the left motor cortex at an intensity of 2 mA for a period of 10 min.	Part A: Cycling TTE at 70% peak PO Part B: CPT with right hand submerged in iced water (0-1°C)	20°C, 50% RH	Part A: ↔ TTE, ↔ EIP ↔ RPE, ↔ HR, ↔ VO_2, ↔ VE ↔ BLa Part B: ↔ pain tolerance, ↓ pain intensity
Flood et al. 2017 [46]	12 males, recreationally active	High definition tDCS applied to the hand motor cortex at an intensity of 2 mA for a period of 20 min.	30% MVC of knee extensor flexion held at 90° isometrically until exhaustion	Not stated	↔ TTE or maximal force production ↑ conditioned pain modulation

↔ = no change, BLa = blood lactate concentration, BF = biceps femoris, CPT = cold pressor test, EIP = exercise induced pain, HR = heart rate, MVC = maximal voluntary contraction, PO = power output, RH = relative humidity, RPE = rating of perceived exertion, tDCS = transcranial direct current stimulation, TENS = transcutaneous electrical nerve stimulation, TT = time trial, TTE = time to exhaustion, VE = volume of expired air, VL = vastus lateralis, VO_2max = maximal oxygen uptake.

Table 2: Research determining the effects of interventions acting on temperature perception and exercise performance.

Study	Participants	Intervention	Testing Protocol	Testing Conditions	Outcomes
Schulze et al. 2015 [56]	7 males, VO_2max = 62 ± 3 mL·kg^{-1}·min^{-1}	Ingestion of 15 g·kg^{-1} ice slurry + ice towels on legs and torso in 30 min + *ad-libitum* ingestion of ice slurry during TT	20 km cycling TT	34°C, 80% RH	↔ Mean power during TT ↑ Mean power by 9 W (1%) during pre-load ↓ TS, ↑ TC, ↓ T_{CORE}, ↓ T_{MS}
Schlader et al. 2011 [60]	12 males, untrained	Topical application of menthol gel on the face (0.5 g·100 cm^2 at 8% prior to protocol)	Cycling TTE RPE clamp protocol at 16 'hard-very hard' in a WPS to ↑ heat stress	20°C, 48% RH	↑ Total work by 39 kJ (21%) ↓ TS, ↑ TC
Kenny et al. 2011 [61]	10 males, untrained	Ice vest worn under NBC	Walking TTE (or 120 min) at 3 mi·h^{-1} in NBC	35°C, 65% RH	↑ TTE by 12 min (12%) ↓ TS, ↓ T_{CORE}, ↓ T_{MS}, ↓ HR, ↓ RPE
Luomala et al. 2012 [62]	7 males, VO_2max = 56 ± 3 mL·kg^{-1}·min^{-1}	Ice vest worn after 30 min of cycling	Cycling TTE (9 min at 60%, 1 min at 80% VO_2max)	30°C, 40% RH	↑ TTE by 13 min (22%) ↓ TS, ↑ TC, ↓ T_{CH}, ↓ T_{UB}
Booth et al. 1997 [68]	5 males, 3 females, VO_2peak = 63 mL·kg^{-1}·min^{-1}	Cold-water immersion for 30 min at 24-29°C	30 min running TT	32°C, 60% RH	↑ TT distance by 304 m (4.2%) ↑ TC, ↓ T_{CORE}, ↓ T_{MS}, ↓ HR
Barwood et al. 2008 [70]	18 males, untrained	Psychological skills training involving goal setting, arousal regulation, mental imagery and positive self talk	90 min running TT	30°C, 40% RH	↑ TT distance by 1.2 km (8%) ↔ T_{AU}, ↔ T_{MS}, ↔ SR, ↔ IL-6, ↔ PRL
Barwood et al., 2015 [71]	8 males, untrained	Menthol sprayed on the cycling jersey (100 mL at 0.2% after 10 km of TT)	16.1 km cycling TT	34°C, 33% RH	↔ TT time ↓ TS, ↑ TC, ↓ RPE
Flood et al. 2017 [72]	8 males, VO_2peak = 55.4 mL·kg^{-1}·min^{-1}	Menthol mouth rinse (25 mL at 0.01% every 10 min)	Cycling TTE RPE clamp protocol at 16 'hard-very hard'	35°C, 48% RH	↑ TTE by 99 s (7.6%) and ↑ Power by 6 W (3.6%) ↓ TS, ↔ TC, ↔ T_{CORE}, ↔ T_{MS}, ↔ SR
Stevens et al. 2016 [75]	11 males, 5 km run time of 18-22 min	Menthol mouth rinse (25 mL at 0.01% every 1 km)	5 km running TT	33°C, 46% RH	↓ TT time by 0.7 min (3%) ↓ TS, ↑ VE, ↑ PRL, ↔ SR
Stevens et al. 2016 [79]	9 males, 5 km run time of 18-22 min	Facial water spray (every 1 km)	5 km running TT	33°C, 46% RH	↓ TT time by 0.6 min (3%) ↓TS, ↓ T_F, ↑ iEMG, ↔ SR
Sunderland et al. 2008 [93]	17 females, trained in team sports	4x30-45 running training sessions (30°C, 27% RH)	Loughborough Intermittent Shuttle Test	30°C, 27% RH	↑ distance by 2.5 km (33%) ↑ TC, ↓ T_{CORE}, ↔ progesterone, ↔ aldosterone, ↔ cortisol

Kelly et al. 2016 [94]	14 males, professional AF players	5x27 min cycling training sessions (39°C, 34% RH)	30 min cycling at 60% VO_2peak	38°C, 29% RH	↑ TC, ↓ RPE, ↓ BLa ↔ VO_2, ↔ T_{CORE}, ↔ T_{MS}
James et al. 2017 [95]	17 males, mean 5 km perf time 20:51	5x90 min running training sessions (36.6°C, 59% RH) to target T_{CORE} 38.5°C	5 km running TT	32°C, 60% RH	↓ TT time by 98 s (6.2%) ↓ TS, ↓ T_{CORE}
Wallace et al. 2017 [104]	14 males and 4 females, trained	2-weeks of motivational self-talk training	30 min cycling at 60%, then TTE at 80% peak power output	35°C, 50% RH	↑ TTE by 192 s (40%) ↑ executive function ↔ HR, ↔ VO_2, ↔ RPE

↔ = no change, AF = Australian Football, BLa = blood lactate concentration, HR = heart rate, iEMG = integrated electromyography, IL = interleukin, NBC = nuclear biological chemical suit, PRL = blood prolactin concentration, RH = relative humidity, RPE = rating of perceived exertion, SR = sweat rate, T_{AU} = aural temperature, TC = thermal comfort, T_{CORE} = core temperature, T_{CH} = chest skin temperature, T_F = forehead skin temperature, T_{MS} = mean skin temperature, T_{UB} = upper back skin temperature, TS = thermal sensation, TT = time trial, TTE = time to exhaustion, VE = volume of expired air, VO_2max = maximal oxygen uptake, VO_2peak = peak oxygen uptake, WPS = water perfused suit.